

Division of Coleoptera  
Department of Entomology  
U. S. National Museum

Edward A. Chapin

from

Dick B. Blackwelder

1940










584  
C2 F19  
ENT

LIST  
OF THE  
Coleoptera  
OF  
Southern California,  
WITH  
Notes on Habits and Distribution  
AND  
Descriptions of New Species  
BY  
H. C. FALL

*Issued November 11, 1901*

SAN FRANCISCO  
CALIFORNIA ACADEMY OF SCIENCES  
1901





LIST OF THE COLEOPTERA OF SOUTHERN CALIFORNIA, WITH NOTES ON HABITS AND DISTRIBUTION AND DESCRIPTIONS OF NEW SPECIES.

BY H. C. FALL.

---

INTRODUCTION.

THE Coleoptera of California, or of any considerable part thereof, have never been made the subject of exclusive treatment by any of our entomologists. Portions of the State have, it is true, formed a part of much more extensive areas treated in some of the earlier faunal papers of Le Conte, but in none of importance have the Californian species been so numerous as to constitute a majority of those listed. Of these papers, two are especially worthy of mention, viz: The Report on the Insects collected on the Pacific R. R. Survey adjacent to the 47th parallel (1857), and the Catalogue of the Coleoptera of the Region adjacent to the Mexican boundary line (1858). In the former, 1,173 species are listed from Northern California, Oregon, Washington, and Alaska, of which 520 are Californian; in the latter, of about 1,000 species mentioned, 300 are credited to Southern California. To both of these, short supplements were added in 1858 and 1859. In an earlier paper (1851), Le Conte had published descriptions of 335 species from various parts of California, all of which are incorporated in the two lists above mentioned.

Previous to the time of Le Conte our knowledge of the Coleoptera of the Pacific Coast was due entirely to the writings of Eschscholtz (1829) and Mannerheim (1843-1853). Both of these authors confined their attention to the fauna of the northern coast region—more especially of Alaska—and, as is remarked by Le Conte, at the close of Mannerheim's labors, the fauna of that region was more completely developed than that of any other part of the continent.

In 1859 Le Conte published a list of 147 species collected by Xantus at Fort Tejon. The number is insignificant, but the list is interesting in the present connection, since Fort Tejon lies at the southern end of the San Joaquin Valley, on the northern slope of the mountains which separate the valley from Southern California. Of the 147 species named, 120 are now known to occur south of the mountains.

From 1859 to 1876 several papers of a faunal nature were contributed by Le Conte, Horn, and Ulke, none of which, however, are so important as to require special mention. In these, as in the two first named, the references to Californian species are for the most part contained in lists covering a much wider territory. Since 1876 only a few local lists have appeared, the most important being that contained in the report of the Death Valley Expedition;\* the Tenebrionidæ, Cicindelidæ, Histeridæ, and Coccinellidæ observed in San Diego County by Dr. F. E. Blaisdell† and two lists presented by the author, one of a collection made in the San Bernardino Mts.,‡ and the other of the Coleoptera known from the Southern California islands.§

---

\* North American Fauna, No. 7, 1893. U. S. Department of Agriculture.

† *Zoe*, Vol. III, 1892; *Entomological News*, Vol. III, 1892.

‡ *Entomological News*, Vol. V, 1894.

§ *Canadian Entomologist*, 1897.

It were desirable for many reasons that the scope of the present work be enlarged to cover the fauna of the entire State. I do not, however, at present feel sufficiently familiar with the northern fauna to warrant so great an undertaking. Indeed, I have many times been tempted to indefinitely postpone the present paper, because of the very meager knowledge which we possess of the great majority of even the commoner species of our district. With a view to completeness, however, and in order to increase the usefulness of the present list, I have in the "Notes" added at the end of each genus the names—nearly always with localities, but usually without further comment—of all other species known to have occurred within the State. It is not unlikely that a certain number of these, recorded by the older writers simply from "California," may have been found within the limits of Southern California; but I have very rarely ventured to include a name in the list without definite knowledge of its occurrence within the limits of the territory here treated.

Briefly described, the term Southern California, as here used, includes that part of the State lying to the south and east of a line drawn from Point Conception eastward along the Santa Inez Mountains, then curving to the north and east around the southern end of the San Joaquin Valley, and along the desert slopes of the Sierra Nevada Mountains through Kern and Inyo counties, to the Nevada state line. The region thus defined is by no means a distinct zoological district, but its northern boundary line is as nearly an interfaunal one as it is possible to draw across the State. Its area is approximately one-third that of the entire State, and includes San Diego, Riverside, San Bernardino, Orange, Los Angeles, and Ventura counties; the southern part

of Santa Barbara, the eastern part of Kern, and the greater portion of Inyo counties.

To the student who is unacquainted with the topography and climatic conditions of California, the following brief sketch of the natural features, more especially of the region under consideration, will be of service.

The State lies approximately between the parallels of  $33^{\circ}$  and  $42^{\circ}$  of latitude, and is crossed by two great mountain systems, lying along and in general parallel with its eastern and western borders. The former—the Sierra Nevada—extends from near the northern border to about the 35th parallel. Its average elevation is from 5,000 to 8,000 feet, but there are occasional ridges of 9,000 and 10,000 feet altitude, and individual peaks that exceed 14,000 feet. The eastern slope descends quite abruptly to the Great Basin, itself 4,000 to 5,000 feet above the sea level; the western slope is much more gradual.

The Coast Range consists, like the Sierra Nevada, of numerous nearly parallel ridges, and extends the entire length of the State. Its general altitude is from 2,000 to 6,000 feet. These two systems are united near Mt. Shasta in the north by a series of cross ranges, and again in the south near the northern boundary of Los Angeles County, thus inclosing the great central valley, some 400 miles long and 60 miles wide, drained by the Sacramento and San Joaquin rivers, which, after their junction, find their way to the ocean through a gap in the Coast Range at the Golden Gate. South of this junction of the Sierra Nevada and Coast Range, the two systems are continued as a single one, running in a generally southeasterly direction toward the Colorado River, sending off, however, at about the middle of its length, a branch to the south, known as the San Jacinto



Range. The principal chain, extending from the Tejon to the Colorado River, has received various local names, the central and highest portion being known as the San Bernardino Mountains. Taken as a whole, it is for convenience frequently referred to in the following pages as the Southern Sierras.

To the east of these mountains lie the Colorado and Mojave deserts, while to the west the country is broken, especially toward the coast, by numerous short ranges of no great altitude, between which lie fertile valleys.

The climate of Southern California varies much according to locality, but it is in general mild and dry. Along the coast the mean temperatures of January and July differ by but little more than  $10^{\circ}$ , but toward the interior the difference between summer and winter temperature becomes rapidly more marked. Except in the higher parts of the mountains, the mercury rarely drops below  $28^{\circ}$  Fahrenheit in winter, but not infrequently rises to or even exceeds  $100^{\circ}$  in the summer months in the interior valleys. The summers in the desert region of the southeast are excessively hot, a temperature of  $130^{\circ}$  being at times experienced at Yuma and other points along the lower Colorado River.

In all parts of our territory the year is divided into two well defined seasons, a rainy and a dry. The former usually begins toward the end of October and lasts till April. During this season the fair days greatly outnumber the rainy ones, and the precipitation is on the average less than that during any period of equal length on the Atlantic slope. The average rainfall at Yuma is less than 4 inches; at San Diego 10 inches; at Los Angeles 20 inches, and becomes somewhat greater as we ascend the western slopes of the mountains. Snow falls every winter at altitudes of 3,000 feet and over, but

lasts only a short time, except on the highest peaks, where, in sheltered spots on northern slopes, it may be seen in midsummer. At intervals of many years the ground is whitened for a few hours in the valleys.

The locally diverse conditions of temperature and rainfall are of necessity accompanied by a variation in the character and abundance of the vegetation, and consequently of the insect fauna. Only along the crests or upper slopes of the higher ranges—notably of the San Bernardino Mountains—is there anything approaching forest growth. At these altitudes—5,000-8,000 feet—there are considerable areas covered sparsely with pine, cedar, fir, and scattered oaks. Alders line the streams in the mountain cañons, while willows, sycamores and cottonwoods grow along the water courses at lower levels wherever the supply of moisture is sufficiently permanent. Oaks of several species are more or less common throughout the lower foothills, and occur here and there in park-like groves in the valleys. The foothills are everywhere covered with a mixed growth of scrub-oak, *Ceanothus*, greasewood, manzanita, and a variety of other bushes and low trees, which, in some places, form almost impenetrable thickets. The valleys are green in the rainy season and in early spring are bright with a profusion of wild flowers, but are burned dry and brown by the sun in summer and autumn. The vegetation of the Colorado and Mojave deserts is naturally limited in amount and peculiar in kind, consisting mainly of *Yucca*, cactus, and mesquite, with the usual “sagebrush”.

From the preceding remarks it may be seen at once that Southern California is separable into two principal faunal districts—one the desert region to the eastward, occupying more than half the entire area and extending

far to the north and east beyond the State line; the other, the more copiously watered country lying between the principal mountain range and the coast. To these two we may add two others of very limited area, but quite as well defined, viz, the timber belt of the higher ranges; and the beaches, salt flats, and sand dunes immediately adjoining the seashore.

While it is true that a large proportion of the species occurring in each of these districts is to be found in none of the others, yet our acquaintance with many parts of the territory here considered is still too imperfect to permit the expression of these facts in figures.

In the preparation of the following list I have carefully searched all the literature at my command, and I believe that omissions of this sort will not be numerous; there doubtless are, however, a considerable number of species in collections, which are not here included, for the simple reason that they have not been recorded. Of such as these, and of omissions or corrections of any kind, I would be very glad to be notified in order that they may be incorporated in any revision or supplement which may appear in the future.

Notwithstanding the inevitable shortcomings incident to any preliminary list—and this can scarcely claim to be more—the number of species here credited to Southern California is larger than that of any other faunal list yet published in North America.\* The Southern California list proper contains 2,197 species and 17 varieties, and to these may be added 1,220 species and 4 varieties from other parts of the State, to which reference is made in the notes, making a total of 3,417 species and 21 varieties for the entire State.

---

\* A catalogue of the insects of New Jersey by John B. Smith has appeared since the above was written and contains the names of 2,845 species of Coleoptera—a number greater than that here recorded for Southern California, but considerably smaller than that here credited to the whole State.

Following the notes on distribution, etc., there will be found descriptions of a considerable number of the new species mentioned in the list. There are still many others left undescribed, chiefly for the reason that their proper definition would require a more extended investigation than is now possible. A few allied species from other parts of the country have also been described.

Finally, it is a pleasure to acknowledge the kind and ready assistance of entomological friends in the preparation of the following pages, more particularly of my fellow Californians, Messrs. Fényes, Daggett, Fuchs, Van Dyke, Rivers and Ricksecker.

To Mr. Fuchs I am especially indebted for a careful examination of the manuscript and the consequent detection of a few errors and the addition of a number of species based upon his own collection.

## LIST OF THE COLEOPTERA OF SOUTHERN CALIFORNIA.

## CICINDELIDÆ.

<i>Tetracha carolina</i> Linn.	<i>Cicindela hirticollis</i> Say.
<i>Cicindela latesignata</i> Lec.	<i>tenuisignata</i> Lec.
<i>senilis</i> Horn.	<i>trifasciata</i> var. <i>sigmoidea</i> Lec.
<i>vulgaris</i> Say.	<i>sperata</i> Lec.
var. <i>obliquata</i> Kirby.	<i>gabbii</i> Horn.
var. <i>vibex</i> Horn.	<i>lemniscata</i> Lec.
<i>12-guttata</i> Dej.	<i>circumpicta</i> var. <i>prætextata</i> Lec.
var. <i>oregona</i> Lec.	<i>hæmorrhagica</i> Lec.
var. <i>guttifera</i> Lec.	var. <i>pacifica</i> Schaupp.
<i>pusilla</i> Say.	<i>sommeri</i> Mann.

## CARABIDÆ.

<i>Omophron dentatum</i> Lec.	<i>Dyschirius truncatus</i> Lec. (?).
<i>Cychrus obliquus</i> Lec.	<i>marinus</i> Lec.
<i>striatus</i> Lec.	<i>aratus</i> Lec.
<i>minus</i> Horn.	<i>unipunctatus</i> sp. nov.
<i>punctatus</i> Lec.	<i>Clivina dentipes</i> Dej.
<i>subtilis</i> Schaum.	<i>punctulata</i> Lec.
<i>Calosoma angulatum</i> Chev.	<i>Schizogenius crenulatus</i> Lec.
<i>prominens</i> Lec.	<i>depressus</i> Lec.
<i>peregrinator</i> Guer.	<i>seticollis</i> sp. nov.
<i>parviceps</i> Csy.	<i>pluripunctatus</i> Lec.
<i>triste</i> Lec.	<i>Bembidium nitidum</i> Kby.
<i>semilæve</i> Lec.	<i>erasum</i> Lec.
<i>simplex</i> Lec.	<i>carinatum</i> Lec.
<i>cancellatum</i> Esch.	<i>bifossulatum</i> Lec.
<i>latipenne</i> Horn.	<i>brevistriatum</i> Haywd.
<i>Loricera californica</i> Lec.	<i>longulum</i> Lec.
<i>Notiophilus semiopacus</i> Esch.	<i>recticolle</i> Lec.
obscurus sp. nov.	<i>transversale</i> Dej.
<i>Nebria eschscholtzii</i> Mén.	<i>consanguineum</i> Haywd.
<i>Metrius contractus</i> Esch.	<i>striola</i> Lec.
<i>Promecognathus lævissimus</i> Dej.	<i>platynoides</i> Haywd.
<i>Scarites subterraneus</i> Fab.	<i>lucidum</i> Lec.
<i>Dyschirius tridentatus</i> Lec.	<i>californicum</i> Haywd.
<i>patruelis</i> Lec.	<i>henshawi</i> Haywd.
<i>basalis</i> Lec.	<i>scudderi</i> Haywd.
<i>æneus</i> Dej.	<i>tigrinum</i> Lec.
<i>gibbipennis</i> Lec.	<i>insulatum</i> Lec.
<i>analís</i> Lec.	<i>nubiculosum</i> Chd.

- Bembidium indistinctum Dej.*  
*variegatum Say.*  
*intermedium Kby.*  
*timidium Lec.*  
*versicolor Lec.*  
*grandicollis Lec.*  
*ephippiger Lec.*  
*vile Lec.*  
*horni Haywd.*  
*bifasciatum Mots.*  
*dubitans Lec.*  
*acutifrons Lec.*  
*cautum Lec.*  
*connivens Lec.*  
*trechiforme Lec.*  
*iridescens Lec.*  
*laticeps Lec.*
- Tachys vittiger Lec.*  
*mordax Lec.*  
*rufotestaceus Haywd.*  
*virgo Lec.*  
*vorax Lec.*  
*corax Lec.*  
*edax Lec.*  
*nanus Gyll.*  
*anthrax Lec.*  
*audax Lec.*  
*rapax Lec.*  
*sellatus Lec.*
- Pogonus plauatus Horn.*  
*Trechus barbaræ Horn.*  
*pomonæ sp. nov.*
- Pterostichus ater Dej.*  
*vicinus Mann.*  
*californicus Dej.*  
*menetriesii Mots.*  
*inermis sp. nov.*  
*gracilior Lec.*  
*hornii Lec.*  
*isabellæ Lec.*  
*congestus Mén.*  
*subcordatus Lec.*  
*scitulus Lec.*  
*lætulus Lec.*
- Pterostichus occidentalis Dej.*  
*lustrans Lec.*  
*splendidulus Lec.*  
 four species undescribed
- Amara jacobinæ Lec.*  
*scitula Zimm.*  
*longula Zimm.*  
*insignis Dej.*  
*insularis Horn.*  
*impuncticollis Say.*  
*interstitialis Dej.*  
*californica Dej.*  
*remotestriata Dej.*  
*gibba Lec.*  
*aurata Dej.*
- Badister anthracinus Lec.*  
*Calathus ruficollis Dej.*  
*obscurus Lec.*
- Pristonychus complanatus Dej.*  
*Platynus agilis Lec.*  
*brunneomarginatus Mann.*  
*extensicollis var. simplex Lec.*  
*californicus Dej.*  
*funebis Lec.*  
*frater Lec.*  
*maculicollis Dej.*  
*variolatus Lec.*  
*fossiger Dej.*  
*deplanatus Mén.*
- Perigona nigriceps Dej.*  
*Lachnophorus elegantulus Mann.*  
*Euphorticus occidentalis Horn.*  
*Galerita lecontei Dej.*  
*Thalpius hornii Chd.*  
*rufulus Lec.*  
*Ega lætula Lec.*  
*Tetragonoderus fasciatus Hald.*  
*pallidus Horn.*  
*Lebia cyanipennis Dej.*  
*var. ruficollis Lec.*  
*viridis Say.*  
*furcata Lec.*  
*guttula Lec.*  
*bilineata Mots.*

- Dromius piceus Dej.*  
*Apristus laticollis Lec.*  
*Blechnus glabratus Duft.*  
     *lucidus Lec.*  
*Axinopalpus biplagiatus Dej.*  
     *fusciceps Lec.*  
*Tecnophilus croceicollis Mén.*  
*Philophuga castanea Horn.*  
*Pinacodera punctigera Lec.*  
*Cymindis cribricollis Dej.*  
*Brachynus lateralis Dej.*  
     *fidelis Lec.*  
     *tschernikhii Mann.*  
     *carinulatus Mots.*  
     *costipennis Mots.*  
*Chlænius ruficauda Chd.*  
     *viridifrons Esch.*  
     *cursor Chev.*  
     *cumatilis Lec.*  
     *lencoscelis Chev.*  
     *obsoletus Lec.*  
     *variabilipes Esch.*  
     *glaucus Lec.*  
     *tricolor Dej.*
- Chlænius harpalinus Esch.*  
*Oodes elegans Lec.*  
*Agonoderus lineola Fab.*  
     *pallipes Fab.*  
*Discoderus amœnus Lec.*  
*Harpalus sp.*  
*Stenolophus limbalis Lec.*  
     *anceps Lec.*  
     *cincticollis Lec.*  
     *flavipes Lec.*  
*Bradycellus cognatus Gyll.*  
     *rupestris Say.*  
     *rivalis Lec.*  
     *californicus Lec.*  
*Tachycellus nitidus Dej.*  
     two new species  
*Anisodactylus dilatatus Dej.*  
     *piceus Mén.*  
     *semipunctatus Lec.*  
     *consobrinus Lec.*  
     *californicus Dej.*  
     *amaroides Lec.*  
*Anisotarsus flebilis Lec.*

## AMPHIZOIDÆ.

- Amphizoa insolens Lec.*

## HALIPLIDÆ.

- Haliplus concolor Lec.*  
*Cnemidotus callosus Lec.*
- Cnemidotus simplex Lec.*

## DYTISCIDÆ.

- Laccophilus decipiens Lec.*  
     *terminalis Sharp.*  
     *mexicanus Aubé.*  
     *4-lineatus Horn.*  
*Hydrovatus brevipes Sharp.*  
*Desmopachria latissima Lec.*  
*Bidessus cinctellus Lec.*  
     *pictodes Sharp.*  
     *affinis Say.*  
     *subtilis Lec.*  
     *amandus Lec.*
- Hygrotus hydropicus Lec.*  
*Cœlambus medialis Lec.*  
     *pedalis sp nov.*  
     *fraternus Lec.*  
*Deronectes griseostriatus De G.*  
     *striatellus Lec.*  
*Hydroporus addendus Cr.*  
     *subpubescens Lec.*  
     *fortis Lec.*  
     *axillaris Lec.*  
     *vilis Lec.*



- |                                     |                                    |
|-------------------------------------|------------------------------------|
| Hydroporus latebrosus <i>Lec.</i>   | Rhantus binotatus <i>Harr.</i>     |
| Coptotomus interrogatus <i>Fab.</i> | anisonychus <i>Cr.</i>             |
| Hydrotrupes palpalis <i>Sharp.</i>  | Corymbites strigatus <i>Lec.</i>   |
| Ilybiosoma regularis <i>Lec.</i>    | Eretes sticticus <i>Linn.</i>      |
| Agabinus glabrellus <i>Mots.</i>    | Dytiscus marginicollis <i>Lec.</i> |
| Agabus lugens <i>Lec.</i>           | Thermonectes basilaris var. inter- |
| lineellus <i>Lec.</i>               | medius <i>Cr.</i>                  |
| morosus <i>Lec.</i>                 | marmoratus <i>Hope.</i>            |
| obsoletus <i>Lec.</i>               | Cybister ellipticus <i>Lec.</i>    |
| lecontei <i>Cr.</i>                 | explanatus <i>Lec.</i>             |

## GYRINIDÆ.

- |                              |                              |
|------------------------------|------------------------------|
| Gyrinus plicifer <i>Lec.</i> | Gyretes sinuatus <i>Lec.</i> |
| consobrinus <i>Lec.</i>      |                              |

## HYDROPHILIDÆ.

- |                                      |   |
|--------------------------------------|---|
| Helophorus obscurus <i>Lec.</i>      | Berosus infuscatus <i>Lec.</i>          |
| Hydrochus variolatus <i>Lec.</i>     | rugulosus <i>Horn.</i>                  |
| vagus <i>Lec.</i>                    | Chætarthria nigrella <i>Lec.</i>        |
| sp. nov.                             | minor sp. nov.                          |
| Ochthebius rectus <i>Lec.</i>        | pallida <i>Lec.</i>                     |
| costipennis sp. nov.                 | Limnebius piceus <i>Horn.</i>           |
| puncticollis <i>Lec.</i>             | Laccobius ellipticus <i>Lec.</i>        |
| discretus <i>Lec.</i>                | Philydrus carinatus <i>Lec.</i>         |
| nitidus <i>Lec.</i>                  | nebulosus <i>Say.</i>                   |
| interruptus <i>Lec.</i>              | californicus <i>Horn.</i>               |
| lineatus <i>Lec.</i>                 | diffusus <i>Lec.</i>                    |
| sculptus <i>Lec.</i>                 | Helochares normatus <i>Lec.</i>         |
| holmbergi <i>Mükl.</i>               | Cymbiodyta punctatostriata <i>Horn.</i> |
| Hydræna pennsylvanica <i>Kies.</i>   | dorsalis <i>Mots.</i>                   |
| Hydrophilus triangularis <i>Say.</i> | Hydrobius fuscipes <i>Linn.</i>         |
| insularis <i>Lap.</i>                | Creniphilus subcupreus <i>Say.</i>      |
| Tropisternus limbalis <i>Lec.</i>    | elegans sp. nov.                        |
| californicus <i>Lec.</i>             | rufiventris <i>Horn.</i>                |
| salsamentus sp. nov.                 | infuscatus <i>Mots.</i>                 |
| ellipticus <i>Lec.</i>               | Dactylosternum cacti <i>Lec.</i>        |
| Hydrocharis glaucus <i>Lec.</i>      | Cercyon fimbriatus <i>Mann.</i>         |
| obtusatus <i>Say.</i>                | luniger <i>Mann.</i>                    |
| Berosus punctatissimus <i>Lec.</i>   | fulvipennis <i>Mann.</i>                |
| miles <i>Lec.</i>                    | nigriceps <i>Marsh.</i>                 |
| salinus sp. nov.                     | lugubris <i>Payk.</i>                   |
| subsignatus <i>Lec.</i>              | Megasternum posticatum <i>Mann.</i>     |



## SILPHIDÆ.

<i>Necrophorus marginatus</i> <i>Fab.</i>	<i>Anisotoma humeralis</i> <i>Horn.</i>
<i>guttula</i> <i>Mots.</i>	<i>difficilis</i> <i>Horn.</i>
<i>pustulatus</i> var. <i>nigritus</i> <i>Mann.</i>	<i>paludicola</i> <i>Cr.</i>
<i>Silpha lapponica</i> <i>Hbst.</i>	<i>obsoleta</i> <i>Melsh.</i>
<i>ramosa</i> <i>Say.</i>	<i>Cyrtusa picipennis</i> <i>Lec.</i>
<i>Agyrtes longulus</i> <i>Lec.</i>	<i>Agathidium revolvens</i> <i>Lec.</i>
<i>Pinodytes cryptophagoides</i> <i>Mann.</i>	<i>concinnum</i> <i>Mann.</i>
<i>Ptomaphagus consobrinus</i> <i>Lec.</i>	<i>virile</i> sp. nov.
<i>californicus</i> <i>Lec.</i>	<i>pulchrum</i> <i>Lec.</i>
<i>Hydnobius latidens</i> <i>Lec.</i>	<i>Clambus</i> sp.

## SCYDMÆNIDÆ.

<i>Connophron occidens</i> <i>Csy.</i>	<i>Veraphis colon</i> <i>Horn.</i>
<i>digressum</i> <i>Csy.</i>	<i>Ceramphis deformata</i> <i>Horn.</i>
<i>Scydmaenus ovipennis</i> <i>Csy.</i>	<i>Papusus macer</i> <i>Csy.</i>

## PSELAPHIDÆ.

<i>Articerns fuchsii</i> var. <i>californicus</i> <i>Brend.</i>	<i>Reichenbachia turgidicornis</i> <i>Csy.</i>
	<i>tumidicornis</i> <i>Csy.</i>
<i>Biotus formicarius</i> <i>Csy.</i>	<i>sagax</i> <i>Lec.</i>
<i>Ctenisis dispar</i> <i>Sharp.</i>	<i>Bryaxis foveata</i> <i>Lec.</i>
<i>Tychus tenellus</i> <i>Lec.</i>	<i>loripes</i> <i>Csy.</i>
<i>hexagonus</i> <i>Csy.</i>	<i>Oropus</i> sp.
<i>Scalenarthrus hornii</i> <i>Lec.</i>	<i>Sagola isabellæ</i> <i>Lec.</i>
<i>Decarthron brendeli</i> <i>Csy.</i>	<i>Enplectus orbiceps</i> <i>Csy.</i>
<i>Pselaptus belfragei</i> <i>Lec.</i>	<i>californicus</i> <i>Csy.</i>
<i>Reichenbachia deformata</i> <i>Lec.</i>	<i>Actium californicum</i> <i>Lec.</i>
<i>falli</i> <i>Csy.</i>	

## STAPHYLINIDÆ.

<i>Falagria læviuscula</i> <i>Lec.</i>	<i>Tachyusa</i>
<i>cavipennis</i> <i>Lec.</i>	four species
sp. dub.	<i>Tinotus caviceps</i> <i>Csy.</i>
<i>Echidnoglossa</i>	<i>Myrmedonia fauvelii</i> <i>Sharp.</i>
five species	<i>Phloeopora</i> (?)
<i>Hoplandria</i> (?) sp.	two species
<i>Pontomalota opaca</i> <i>Lec.</i>	<i>Polystoma arenaria</i> <i>Csy.</i>
<i>Atheta</i>	<i>pacifica</i> <i>Csy.</i>
twenty-five species	<i>Aleochara sulcicollis</i> <i>Mann.</i>
<i>Lomechusa angusta</i> sp. nov.	<i>puberula</i> <i>Klug.</i>
<i>Tarphiota pallidipes</i> <i>Csy.</i>	four undetermined species
<i>fucicola</i> <i>Mäkl.</i>	<i>Maseochara valida</i> <i>Lec.</i>

- Maseochara puberula* *Csy.*  
*Thiasophila* sp.  
*Isoglossa* sp.  
*Oxypoda*  
     five species  
*Thinusa* sp.  
*Bolitochara californica* *Csy.*  
     sp.  
*Leptusa* sp.  
*Oligota* sp.  
*Bryobiota bicolor* *Csy.*  
*Somatium oviforme* *Csy.*  
*Gyronecha obscura* *Csy.*  
     sp.  
*Bamona falliana* *Csy.*  
*Myllaena* sp.  
*Acylophorus pronus* *Er.*  
*Heterothops fumatus*  
     var. *californicus* *Lec.*  
     *carbonatus* sp. nov.  
     *pusio* *Lec.*  
     *occidentis* *Csy.*  
*Quedius explanatus* *Lec.*  
     *fulgidus* *Fab.*  
     *erythrogaster* *Mann.*  
     *capucinus* *Grav.*  
     *desertus* *Horn.*  
     *limbifer* *Horn.*  
     *prostrans* *Horn.*  
*Thinopinus pictus* *Lec.*  
*Creophilus villosus* *Grav.*  
*Hadrotes crassus* *Mann.*  
*Staphylinus saphyrinus* *Lec.*  
     *luteipes* *Lec.*  
     *tarsalis* *Mann.*  
*Belonuchus ephippiatus* *Say.*  
     sp. nov.  
*Philonthus nitescens* *Horn.*  
     *alutaceus* *Horn.*  
     *semiruber* *Horn.*  
     *hepaticus* *Er.*  
     *flavolimbatus* *Er.*  
     *grandicollis* *Horn.*  
     *longicornis* *Steph.*  
*Philonthus discoidens* *Grav.*  
     *alumnus* *Er.*  
     *pettiti* *Horn.*  
     *triangulum* *Horn.*  
     *ferreipennis* *Horn.*  
     *sordidus* *Grav.*  
     *versutus* *Horn.*  
     *virilis* *Horn.*  
     *nigritulus* *Grav.*  
     *instabilis* *Horn.*  
     *quadrulus* *Horn.*  
     *lecontei* *Horn.*  
*Actobius ocreatus* *Horn.*  
     *sobrinus* *Er.*  
     *puncticeps* *Horn.*  
     *gratus* *Lec.*  
     *pæderoides* *Lec.*  
     *elegantulus* *Horn.*  
     *formosus* sp. nov.  
*Cafius canescens* *Mann.*  
     *seminitens* *Horn.*  
     *lithocharinus* *Lec.*  
     *luteipennis* *Horn.*  
     *sulcicollis* *Lec.*  
     *decipiens* *Lec.*  
     *opacus* *Lec.*  
*Xantholinus cephalus* *Say.*  
     *picipennis* *Lec.*  
     *dimidiatus* *Lec.*  
     *obscurus* *Er.*  
     *nannus* *Lec.*  
     *pusillus* *Sachse.*  
*Leptacinus brunnescens* *Lec.*  
     *pallidulus* *Lec.*  
     three species, probably undescribed  
*Stenus renifer* *Lec.*  
     *sculptilis* *Csy.*  
     *zunicus* *Csy.*  
     *incultus* *Csy.*  
     *costalis* *Csy.*  
     *alveolatus* *Csy.*  
     *pacificus* *Csy.*  
     *insignis* *Csy.*

- Stenus terricola* *Csy.*  
*vestalis* *Csy.*  
*californicus* *Csy.*  
*gilæ* *Csy.*  
*sayi* *Csy.*  
*pinguis* *Csy.*  
*lætulus* *Csy.*  
*lucidus* *Csy.*  
 two species undetermined  
*Cryptobium tumidum* *Lec.*  
*californicum* *Lec.*  
 species undetermined  
*Lathrobium puncticeps* *Lec.*  
*jacobinum* *Lec.*  
*californicum* *Lec.*  
*lituarium* *Lec.*  
 sp. nov.  
*Caloderma rugosa* *Csy.*  
*continens* *Csy.*  
*mobile* *Csy.*  
*reducta* *Csy.*  
 three species undetermined  
*Medon malaca* *Csy.*  
*latiuscula* *Csy.*  
 four species undetermined  
*Lithocharis ochracea* *Grav.*  
*Stilicus occiduus* sp. nov.  
*Pælerus femoralis* *Lec.*  
*compotens* *Lec.*  
*ustus* *Lec.*  
*Sunius californicus* *Aust.*  
*longiusculus* *Mann.*  
*Scopæus truncaticeps* *Csy.*  
*armiger* sp. nov.  
*Scopæodera nitida* *Lec.*  
*Leptorus texanus* *Csy.*  
*californicus* sp. nov.  
*longipennis* sp. nov.  
*Orus punctatus* *Csy.*  
*fraternus* sp. nov.  
*montanus* sp. nov.  
*Pinophilus densus* *Lec.*  
*Palaminus lividus* *Lec.*  
*Tachinus agilis* *Horn.*  
*Tachinus debilis* *Horn.*  
*Tachyporus californicus* *Horn.*  
*nitidulus* *Fab.*  
*Cilea silphoides* *Linn.*  
*Erchomus punctipennis* *Lec.*  
*Conosoma bipustulatum* *Grav.*  
*castaneum* *Horn.*  
*Boletobius cincticollis* *Say.*  
*Bryoporus* sp. dub.  
*Pseudopsis oblitterata* *Lec.*  
*detrita* sp. nov.  
*minuta* sp. nov.  
*Oxyporus* sp. nov.  
*Bledius ferratus* *Lec.*  
*jacobinus* *Lec.*  
*armatus* *Er.*  
*cribricollis* *Lec.*  
*eximius* *Csy.*  
*flavipennis* *Lec.*  
*nitidiceps* *Lec.*  
*relictus* sp. nov.  
*opacifrons* *Lec.*  
*punctatissimus* *Lec.*  
*laticollis* *Lec.*  
*luteipennis* *Lec.*  
*rusticens* sp. nov.  
*ruficornis* *Lec.*  
*clarus* sp. nov.  
*pleuralis* *Lec.*  
*diagonalis* *Lec.*  
*ornatus* *Lec.*  
*phytosinus* *Lec.*  
*forcipatus* *Lec.*  
*Platystethus americanus* *Er.*  
*Oxytelus sculptus* *Grav.*  
*sobrinus* *Lec.*  
*Haploderus cephalotes* *Csy.*  
*flavipennis* *Csy.*  
*Trogophlæus dentiger* *Csy.*  
*gilæ* *Csy.*  
*prominens* *Csy.*  
*pacificus* *Csy.*  
*diffusus* *Csy.*  
*confinis* *Csy.*

<i>Trogophlœus pauperculus</i> <i>Csy.</i>	<i>Homalium repandum</i> <i>Er.</i>
<i>debilis</i> <i>Csy.</i>	<i>plagiatum</i> <i>Mann.</i>
<i>blediinus</i> <i>Csy.</i>	<i>humile</i> <i>Mükl.</i>
<i>flum</i> <i>Csy.</i>	<i>alutaceum</i> <i>Fauv.</i>
<i>tantillus</i> <i>Csy.</i>	<i>theveneti</i> <i>Fauv.</i>
<i>Apocellus analis</i> <i>Lec.</i>	two species undetermined
<i>gracilicornis</i> <i>Csy.</i>	<i>Anthobium atriventre</i> <i>Csy.</i>
<i>sphæricollis</i> <i>Say.</i>	<i>nigerrimum</i> <i>Csy.</i>
<i>Ancyrophorus planus</i> <i>Lec.</i>	two species undetermined
<i>Thinobius oxytelinus</i> <i>Lec. (?)</i>	<i>Orobanus rufipes</i> <i>Csy.</i>
<i>hesperius</i> <i>Csy.</i>	<i>Protinus</i>
<i>sp. nov.</i>	two species
<i>Zalobius serricollis</i> <i>Lec.</i>	<i>Lispinus californicus</i> <i>Lec.</i>
<i>Geodromicus temporalis</i> <i>Csy.</i>	<i>linearis</i> <i>Er.</i>
<i>Amphichroum floribundum</i> <i>Lec.</i>	<i>Glyptoma costale</i> <i>Er.</i>
<i>puberulum</i> <i>Fauv.</i>	<i>Trigonurus edwardsii</i> <i>Sharp.</i>
<i>Lathrimæum subcostatum</i> <i>Mükl.</i>	<i>Micropeplus punctatus</i> <i>Mükl.</i>
<i>Homalium strigipenne</i> <i>Mükl.</i>	<i>sp. nov. (?)</i>

## TRICHOPTERYGIDÆ.

<i>Motschulskium sinuatocolle</i> <i>Matth.</i>	<i>Trichopteryx laticollis</i> <i>Mann.</i>
<i>Ptilium</i> <i>sp.</i>	several species not identified.
<i>Ptenidium pullum</i> <i>Mükl.</i>	<i>Smicrus filicornis</i> <i>Fairm.</i>

## HYDROSCAPHIDÆ.

*Hydroscapha natans* *Lec.*

## SPHÆRIIDÆ.

*Sphærius politus* *Horn.*

## SCAPHIDIIDÆ.

<i>Scaphisoma castaneum</i> <i>Mots.</i>	<i>Scaphisoma rufulum</i> <i>Lec.</i>
--	---------------------------------------

## PHALACRIDÆ.

<i>Phalacrus ovalis</i> <i>Lec.</i>	<i>Acyломus nebulosus</i> <i>Csy.</i>
<i>penicillatus</i> <i>Say.</i>	<i>Eustilbus apicalis</i> <i>Melsh.</i>
<i>conjunctus</i> <i>Csy.</i>	<i>obtusus</i> <i>Lec.</i>
<i>sp. nov. (?)</i>	<i>nanulus</i> <i>Csy.</i>
<i>Olibrus wickhami</i> <i>Csy.</i>	<i>notabilis</i> <i>sp. nov.</i>
<i>sp. nov.</i>	

## CORYLOPHIDÆ.

<i>Sacium amabile</i> Lec.	<i>Orthoperus</i>
<i>decolor</i> Lec.	two species unidentified
<i>scitulum</i> Lec.	<i>Aenigmaticum californicum</i> Csy.
<i>Sericoderus flavidus</i> Lec.	

## COCCINELLIDÆ.

<i>Anisosticta seriata</i> Melsh.	<i>Hyperaspis fimbriolata</i> Melsh.
<i>Megilla maculata</i> De G.	var. <i>dissoluta</i> Cr.
<i>vittigera</i> Mann.	<i>lateralis</i> Muls.
<i>Hippodamia 5-signata</i> Kirby.	<i>tæniata</i> Lec.
<i>ambigua</i> Lec.	<i>excelsa</i> sp. nov.
<i>convergens</i> Guér.	<i>undulata</i> Say. (?)
<i>spuria</i> Lec.	<i>spiculinata</i> sp. nov.
<i>parenthesis</i> Say.	<i>annexa</i> Lec.
<i>Coccinella 9-notata</i> Hbst.	<i>tristis</i> Lec.
var. <i>franciscana</i> Muls.	<i>Hyperaspidium trimaculatus</i> Linn.
<i>transversoguttata</i> var. <i>transver-</i>	<i>arcuatus</i> Lec.
<i>salis</i> Muls.	<i>Scymnus tædatus</i> sp. nov.
var. <i>californica</i> Mann.	<i>sordidus</i> Horn.
<i>Cycloneda sanguinea</i> Linn.	<i>guttulatus</i> Lec.
<i>oculata</i> Fab.	<i>nebulosus</i> Lec.
<i>abdominalis</i> Say.	<i>pallens</i> Lec.
<i>Harmonia picta</i> Rand.	<i>minus</i> sp. nov.
<i>Mysia hornii</i> Cr.	<i>cinctus</i> Lec.
<i>Psyllobora 20-maculata</i> var. <i>tædata</i> Lec.	<i>pacificus</i> Cr.
<i>Chilocorus bivulnerus</i> Muls.	<i>flebilis</i> Horn.
<i>cacti</i> Linn.	<i>cervicalis</i> Muls.
<i>Exochomus pilatii</i> Muls.	<i>marginicollis</i> Mann.
<i>californicus</i> Csy.	<i>ardelio</i> Horn.
<i>fasciatus</i> Csy.	<i>punctum</i> Lec.
<i>childreni</i> Muls.	<i>nanus</i> Lec.
<i>histrio</i> sp. nov.	<i>coniferarum</i> Cr.
<i>Cryptognatha pusilla</i> Lec.	<i>Scymnillus aterrimus</i> Horn.
<i>catalinæ</i> Horn.	<i>Cephaloscymnus occidentalis</i> Horn.
<i>Smilia reversa</i> sp. nov.	<i>ornatus</i> Horn.
<i>ovalis</i> Lec.	<i>Rhizobius lophanthæ</i> Blaisd.

## ENDOMYCHIDÆ.

Gen. et sp. dub.	<i>Aphorista læta</i> Lec.
<i>Aphorista morosa</i> Lec.	

## EROTYLIDÆ.

<i>Languria californica</i> sp. nov.	<i>Tritoma californica</i> Lec.
<i>Dacne californica</i> Horn.	

## COLYDIIDÆ.

<i>Rhagodera tuberculata</i> Mann.	<i>Lasconotus servus</i> Horn.
<i>Anchomma costatum</i> Lec.	pusillus Lec.
<i>Synchita variegata</i> Lec.	<i>Aulonium longum</i> Lec.
<i>Ditoma ornata</i> Lec.	<i>Aglenus brunneus</i> Gyll.
sulcata Lec.	<i>Oxylæmus californicus</i> Cr.
<i>Phlæonemus catenulatus</i> Horn.	<i>Cerylon castaneum</i> Say.
<i>Lasconotus linearis</i> Cr.	

## RHYSSODIDÆ.

*Rhyssodes hamatus* Lec.:

## CUCUJIDÆ.

<i>Silvanus surinamensis</i> Linn.	<i>Læmophlæus nitens</i> Lec.
bidentatus Fab.	hornii Csy.
imbellis Lec.	cephalotes Lec.
<i>Cathartus advena</i> Walth.	ferrugineus Steph.
opaculus Lec.	pusillus Sch.
<i>Nausibius clavicornis</i> Kug.	<i>Lathropus vernalis</i> Lec.
<i>Narthecius grandiceps</i> Lec.	pubescens Csy.
<i>Pediacus depressus</i> Hbst.	<i>Brontes dubius</i> var. <i>truncatus</i> Mots.
<i>Læmophlæus biguttatus</i> Say.	

## CRYPTOPHAGIDÆ.

<i>Henoticus</i> sp.	<i>Cryptophagus</i>
<i>Cryptophagus cellaris</i> Scop.	two species unidentified
debilis Lec.	<i>Atomaria lætula</i> Lec. (?)
lecontei G. & H.	two species unidentified

## MYCETOPHAGIDÆ.

<i>Mycetophagus californicus</i> Horn.	<i>Berginus pumilus</i> Lec.
pluriguttatus Lec.	<i>Myrmechixenis latridioides</i> Cr.
<i>Litargus balteatus</i> Lec.	<i>Diplocælus</i> sp.
<i>Typhœa fumata</i> Linn.	

## DERMESTIDÆ.

<i>Byturus griseus</i> Lec.	<i>Dermestes vulpinus</i> Fab.
<i>Dermestes marmoratus</i> Say.	<i>Perimegatoma variegatum</i> Horn.
mannerheimii Lec.	<i>Attagenus piceus</i> Oliv.
talpinus Mann.	<i>Trogoderma ornatum</i> Say.
tristis Fall.	sternale Jayne.
carnivorus Fab.	<i>Anthrenus scrophulariæ</i> Linn.

<i>Anthrenus verbasci</i> Linn.	<i>Cryptorhopalum ruficorne</i> Lec.
<i>Cryptorhopalum balteatum</i> Lec.	species undescribed
apicale Mann.	<i>Orphilus niger</i> Rossi.

HISTERIDÆ.

<i>Hololepta yucateca</i> Mars.	<i>Saprinus alienus</i> Lec.
cacti Lec.	lugens Er.
vicina Lec.	oregonensis Lec.
populnea Lec.	liticolus sp. nov.
neglecta Blaisd.	scissus Lec.
<i>Hister sellatus</i> Lec.	laridus Lec.
lucanus Lec.	insertus Lec.
simplicipes sp. nov.	ciliatus Lec.
sexstriatus Lec.	vitiosus Lec.
militaris Horn.	lubricus Lec.
bimaculatus Linn.	plenus Lec.
lecontei Mars.	fimbriatus Lec.
punctiger Lec.	cærulescens Lec.
<i>Tribalister marginellus</i> Lec. (?)	intritus Csy.
<i>Tribalus californicus</i> Horn.	consobrinus sp. nov.
<i>Epierus regularis</i> Beauv.	bigemmeus Lec.
planulus Er.	estriatus Lec.
nasutus Horn.	lucidulus Lec.
<i>Onthophilus lecontei</i> Horn.	propensus Csy.
<i>Paromalus difficilis</i> Horn.	gaudens Lec.
<i>Carcinops opuntiae</i> Lec.	serrulatus Lec.
tejonicus Horn.	sulcifrons Lec.
gileusis Lec.	<i>Plegaderus fraternus</i> Horn.
consors Lec.	nitidus Horn.
tenellus Er.	consors Horn.
<i>Anapleus marginatus</i> Lec.	<i>Teretrius placitus</i> Horn.
<i>Saprinus discoidalis</i> Lec.	obliquulus Lec.
interstitialis Lec.	<i>Abræus bolteri</i> Lec.
pectoralis Lec.	<i>Bacanius globulinus</i> Csy.
behrensi Horn.	<i>Acritus maritimus</i> Lec.
obscurus Lec.	volitans sp. nov.
pæminosus Lec.	<i>Æletes basalis</i> Lec.

NITIDULIDÆ.

<i>Brachypterus troglodytes</i> Murr.	<i>Athonæus agavensis</i> Cr.
<i>Cercus sericans</i> Lec.	<i>Carpophilus yuccæ</i> Cr.
<i>Amartus tinctus</i> Mann.	hemipterus Linn.
rufipes Lec.	pallipennis Say.



*Carpophilus dimidiatus* *Fab.*  
*decipiens* *Horn.*  
*discoideus* *Lec.*  
*Colastus truncatus* *Rand.*  
*Conotelus mexicanus* *Murr.*  
*Epuræa*  
*species near rufa* (?)  
*terminalis* *Mann.*  
*avara* *Rand.*

*Epuræa ovata* *Horn.*  
*Nitidula ziezac* *Say.*  
*Omosita discoidea* *Fab.*  
*Perthalyca murrayi* *Horn.*  
*Meligethes brassicæ* *Scop.*  
*Cybocephalus californicus* *Horn.*  
*Cryptarcha concinna* *Melsh.*  
*Smicrips hypocoproides* *Reitt.*  
*Pityophagus rufipennis* *Horn.*

## LATHRIDIIDÆ.

*Holoparamesus kunzei* *Aubé.*  
*pacificus* *Lec.*  
*caularum* *Aubé.*  
*Metophtthalmus rudis* *Fall.*  
*trux* *Fall.*  
*Lathridius armatulus* *Fall.*  
*Coninomus constrictus* *Gyll.*  
*australicus* *Belon.*  
*Enicmus suspectus* *Fall.*  
*desertus* *Fall.*  
*minutus* *Linn.*  
*crenatus* *Lec.*  
*tenuicornis* *Lec.*  
*Cartodere argus* *Reitt.*  
*Revelieria californica* *Fall.*

*Corticaria planula* *Fall.*  
*serrata* *Payk.*  
*occidua* *Fall.*  
*tenuipes* *Fall.*  
*elongata* *Gyll.*  
*ferruginea* *Marsh.*  
*Melanophthalma casta* *Fall.*  
*simplex* *Lec.*  
*insularis* *Fall.*  
*distinguenda* *Com.*  
*gibbosa* *Hbst.*  
*incompta* *Fall.*  
*similata* *Gyll.*  
*americana* *Mann.*  
*Fuchsina occulta* *Fall.*

## TROGOSITIDÆ.

*Alindria teres* *Melsh.*  
*Trogosita virescens* *Fab.*  
*yuccæ* *Cr.*

*Tenebrioides sinuata* *Lec.*  
*Peltis pippingskœldi* *Mann.*  
*Grynocharis pilosula* *Cr.*

## MONOTOMIDÆ.

*Monotoma picipes* *Hbst.*  
*mucida* *Lec.*  
*Phyconomus marinus* *Lec.*

*Hesperobænus abbreviatus* *Mots.*  
*Bactridium striatum* *Lec.*  
*striolatum* *Reitt.* (?)

## BYRRHIDÆ.

*Nosodendron californicum* *Horn.*  
*Amphicyrta dentipes* *Er.*  
*Limnichus californicus* *Lec.*  
*perpolitus* *Csy.*

*Limnichus nebulosus* *Lec.*  
*tenuicornis* *Csy.*  
*naviculatus* *Csy.*  
*Bothriophorus minutus* *Lec.*



## PARNIDÆ.

<i>Psephenus falli</i> Csy.	<i>Elmis divergens</i> Lec.
<i>Throscinus crotchii</i> Lec.	foveatus Lec.
<i>Dryops productus</i> Lec.	<i>Stenelmis nubifer</i> sp. nov.
suturalis Lec.	<i>Macronychus parvulus</i> Horn.

## HETEROCERIDÆ.

<i>Heterocerus gnatho</i> Lec.	<i>Heterocerus collaris</i> Kies.
gemmatus Horn.	pusillus Say.

## DASCYLLIDÆ.

<i>Eurypogon confusus</i> sp. nov.	<i>Helodes</i>
<i>Allopogon villosus</i> Horn.	species undescribed
<i>Anorus piceus</i> Lec.	<i>Cyphon exiguus</i> Horn.
<i>Eucinetus infumatus</i> Lec.	concinuus Lec.
<i>Helodes apicalis</i> Lec.	variabilis Thumb.
	<i>Placonycha edwardsii</i> Lec.

## RHIPICERIDÆ.

*Sandalus californicus* Lec.

## ELATERIDÆ.

<i>Sarpedon scabrosus</i> Bonv.	<i>Hypnoideus pectoralis</i> Say.
<i>Adelocera sparsa</i> Cand.	<i>Anchastus cinereipennis</i> Mann.
<i>Meristhus cristatus</i> Horn.	bicolor var. desertus Horn.
<i>Chalcolepidius webbiai</i> Lec.	<i>Elater hepaticus</i> Melsh.
tartarus Fall.	fastus Lec.
<i>Alaus melanops</i> Lec.	cordifer Lec.
<i>Cardiophorus amplicollis</i> Mots.	ater Lec.
gemmifer Blanch.	longicornis Lec.
luridipes Cand.	carbonicolor Esch. (?)
edwardsii Horn.	dimidiatus Lec.
latiusculus Esch.	<i>Drasterius livens</i> Lec.
tenebrosus Lec.	<i>Megapenthes tartareus</i> Lec.
æneus Horn.	turbulentus Lec.
seniculus Blanch.	aterrimus Horn.
<i>Horistonotus inanus</i> Lec.	stigmatosus Lec.
basalis Horn.	<i>Ludius lecontei</i> Horn.
sufflatus Lec.	ater Cand.
simplex Lec.	<i>Agriotes imperfectus</i> Lec.
flavidus sp. nov.	hispidus Lec.
<i>Esthesopus dispersus</i> Horn.	<i>Dolopius lateralis</i> Esch.
<i>Cryptohypnus squalidus</i> Lec.	<i>Melanotus longulus</i> Lec.
<i>Hypnoideus striatulus</i> Lec.	fissilis Say.
ornatus Lec.	cribricollis Cand.
gradarius Horn.	variolatus Lec.

- |                                 |                                     |
|---------------------------------|-------------------------------------|
| Melanotus sp. nov.              | Corymbites pruininus <i>Horn.</i>   |
| Limoniis mirus <i>Lec.</i>      | leucaspis <i>Germ.</i>              |
| crotchii <i>Horn.</i>           | Asaphes morio <i>Lec.</i>           |
| occidentalis <i>Cand.</i>       | dilaticollis <i>Mots.</i>           |
| ornatulus <i>Lec.</i>           | tumescens <i>Lec.</i>               |
| pilosus <i>Lec.</i>             | Melanactes densus <i>Lec.</i>       |
| californicus <i>Mann.</i>       | Aphricus californicus <i>Lec.</i>   |
| canus <i>Lec.</i>               | Aplastus angusticollis <i>Horn.</i> |
| Athous excavatus <i>Mots.</i>   | corymbitoides <i>Horn.</i>          |
| limbatus <i>Lec.</i>            | speratus <i>Lec.</i>                |
| two undescribed species         | molestus <i>Horn.</i>               |
| Sericosomus debilis <i>Lec.</i> | Plastocerus schaumii <i>Lec.</i>    |
| flavipennis <i>Mots.</i>        | Euthysanius lautus <i>Lec.</i>      |
| Corymbites maurus <i>Say.</i>   | pretiosus <i>Lec.</i>               |
| jaculus <i>Lec.</i>             | Perothops witticki <i>Lec.</i>      |
| fallax <i>Say.</i>              |                                     |

## THROSCIDÆ.

- |                                 |                               |
|---------------------------------|-------------------------------|
| Throscus sejunctus <i>Horn.</i> | Throscus sericeus <i>Lec.</i> |
| parvulus <i>Lec.</i>            | Pactopus hornii <i>Lec.</i>   |

## BUPRESTIDÆ.

- |                                      |  |
|--------------------------------------|--|
| Gyascutus planicosta <i>Lec.</i>     | Chrysobothris femorata <i>Fab.</i>       |
| obliteratus <i>Lec.</i>              | contigua <i>Lec.</i>                     |
| Hippomelas californicus <i>Horn.</i> | cuprascens <i>Lec.</i>                   |
| Dicerca californica <i>Cr.</i>       | speculifer <i>Horn.</i> (?)              |
| hornii <i>Cr.</i>                    | californica <i>Lec.</i>                  |
| pectorosa <i>Lec.</i>                | mali <i>Horn.</i>                        |
| Pæcilonota ferrea <i>Melsh.</i> (?)  | deserta <i>Horn.</i>                     |
| Buprestis gibbsii <i>Lec.</i>        | merkelii <i>Horn.</i>                    |
| læviventris <i>Lec.</i>              | prasina <i>Horn.</i>                     |
| maculiventris <i>Say</i> var.        | lucana <i>Horn.</i>                      |
| fasciata <i>Fab.</i>                 | Actenodes mendax <i>Horn.</i>            |
| aurulenta <i>Linn.</i>               | calcarata <i>Chev.</i>                   |
| Melanophila.                         | Glyptoscelimorpha marmorata <i>Horn.</i> |
| conspua <i>Lec.</i>                  | Dystaxia murrayi <i>Lec.</i>             |
| longipes <i>Say.</i>                 | Schizopus lætus <i>Lec.</i>              |
| acuminata <i>De Geer.</i>            | Polycesta velasco <i>Lap &amp; Gory.</i> |
| gentilis <i>Lec.</i>                 | californica <i>Lec.</i>                  |
| intrusa <i>Horn.</i>                 | Acmæodera flavomarginata <i>Gray.</i>    |
| Anthaxia æneogaster <i>Lap.</i>      | lanata <i>Horn.</i>                      |
| deleta <i>Lec.</i>                   | fenyesi <i>Fall.</i>                     |
| Chrysobothris octocola <i>Lec.</i>   | plagiaticauda <i>Horn.</i>               |
| debilis <i>Lec.</i>                  | jocosa <i>Fall.</i>                      |

*Acmæodera coquilletti Fall.**angelica Fall.**tuta Horn.**hepburnii Lec.**quadriseriata Fall.**morbosa Fall.**flavosticta Horn.**acuta Lec.**labyrinthica Fall.**connexa Lec.**variegata Lec.**vandykei Fall.**prorsa Fall.**dohrni Horn.**dolorosa Fall.**pastica Fall.**alicia Fall.**versuta Horn.**guttifera Lec.**Acmæodera quadrivittata Horn.**gemina Horn.**gibbula Lec.**comata Lec.**alacris Horn.**Ptosima walshii Lec.**Chrysophana placida Lec.**Agrilus angelicus Horn.**niveiventris Horn.**walsinghami Cr.**obolinus Lec.**jacobinus Horn.**politus Say.**blandus Horn.**gibbicollis sp. nov.**illectus sp. nov.**lacustris Lec.**Taphrocerus gracilis Say. (?)*

## LAMPYRIDÆ.

*Eros lætus Mots.**Ellychnia californica Mots.**corrusca Linn.**Pyropyga fenestralis Melsh.**Microphotus angustus Lec.**Pterotus obscuripennis Lec.**Zarhipis integripennis Lec.**Mastinocerus opacus Horn.**Cenophengus debilis Lec.**Podabrus comes Lec.*

species undescribed

*Silis cava Lec.**filigera Lec.**Telephorus consors Lec.**notatus Mann. (?)**ingenuus Lec.**ochropus Lec.**lautus Lec.**Polemios languidus Horn.**Ditemnus obtusus Lec.**Malthodes laticollis Lec.**fuscus Lec.*

## MALACHIDÆ.

*Collops marginicollis Lec.**cribrosus Lec.**argutus sp. nov.**punctulatus Lec.**marginellus Lec.*

species undescribed

*Endeodes abdominalis Lec.**collaris Lec.**Malachius auritus Lec.**inornatus sp. nov.**Malachius pristinus sp. nov.**acutipennis sp. nov.**thevenetii Horn.**directus sp. nov.**nigrinus sp. nov.**prolixicornis sp. nov.**Tanaops abdominalis Lec.**Microlipus laticeps Lec.**longicollis Mots.**Pseudebæus bicolor Lec. (?)*

*Attalus trimaculatus Mots.*

*rufomarginatus Mots.*

*oregonensis Horn.*

*basalis Lec.*

*cinctus Lec.*

*difficilis Lec.*

*lobulatus Lec.*

*transmarinus Fall.*

*Pristoscelis grandiceps Lec.*

*Eudasytes ursinus Csy.*

*Asydates rufiventris Csy.*

*explanatus Csy.*

*Trichochrous compactus Csy.*

*antennatus Mots.*

*griseus Lec.*

*apicalis Csy.*

*umbratus Lec.*

*propinquus Csy.*

*fulvovestitus Csy.*

*suffusus Csy.*

*testaceus Csy.*

*subcalvus Csy.*

*prominens Csy.*

*cuspidatus Csy.*

*fuscus Lec.*

*seriellus Csy.*

*brevicornis Lec.*

*vilis Csy.*

*insignis Csy.*

*suturalis Lec.*

*conspersus Csy.*

*lobatus Csy.*

*barbaræ Csy.*

*sordidus Lec.*

*Trichochrous crinifer Csy.*

*fulvescens Csy.*

*squalidus Lec.*

*pedalis Lec.*

*nigrinus Csy.*

*ænescens Lec.*

*politus Csy.*

*punctipennis Lec.*

ten or twelve doubtful or undescribed species

*Adasytes laciniatus Csy.*

*Listrus interruptus Lec. (?)*

*obscurus Lec.*

*extricatus Csy.*

*difficilis Lec.*

*luteipes Lec.*

*balteellus Csy.*

*famelicus Csy.*

*definitus sp. nov.*

*Dasytellus inconspicuus Csy.*

*Dasytes dissimilis Csy.*

*clementæ sp. nov.*

*macer Csy.*

*pusillus Lec.*

*musculus sp. nov.*

*lineellus Csy.*

*Dasytastes catalinæ Lec.*

*remissus Csy.*

*bicolor Csy.*

*insularis sp. nov.*

*Eschatocrepis constrictus Lec.*

*Allonyx sculptilis Lec.*

*Vectura longiceps Csy.*

*Eurelymis flavipes Lec.*

#### CLERIDÆ.

*Elasmocerus californicus sp. nov.*

*Cymatodera puncticollis Bland.*

*californica Horn.*

*morosa Lec.*

*punctata Lec.*

*fuscula Lec.*

*undulata var. balteata Lec.*

*angustata Spin.*

*Cymatodera ovipennis Lec.*

*var. pilosella Lec.*

*Trogodendron edwardsii Horn.*

*Trichodes ornatus Say.*

*var. tenellus Lec.*

*Clerus quadrisignatus Say.*

*abruptus Lec.*

*eximius Mann.*

*Clerus moestus* Kl.  
*Hydnocera robusta* Horn.  
     *scabra* Lec.  
     *discoidea* Lec.  
     sp. near *pallipennis* Say.  
     *bicolor* Lec.

*Chariessa elegans* Horn.  
     *dichroa* Lec.  
*Cregya fasciata* Lec.  
*Lebasiella maculicollis* Lec.  
*Corynetes rufipes* Fab.  
     *ruficollis* Fab.

## PTINIDÆ.

*Ptinus verticalis* Lec.  
     *interruptus* Lec.  
*Hedobia granosa* Lec.  
*Ernobius debilis* Lec.  
     undescribed species  
*Ozognathus cornutus* Lec.  
     *misellus* Lec.  
*Xestobium*  
     undescribed species  
*Oligomerus*  
     three undescribed species  
*Sitodrepa panicea* Linn.  
*Ctenobium*  
     undescribed species  
*Ptinodes setifer* Lec.  
*Hadrobregmus gibbicollis* Lec.  
*Trypophytus punctatus* Lec.  
     *tenuilineata* Horn.  
*Petalium bistriatum* Say.  
*Vrilletta convexa* Lec.  
*Xyletinus lugubris* Lec.  
*Catorama frontalis* Lec.  
*Hemiptychus pusillus* Lec.  
     *latus* Horn.  
     *luteotectus* sp. nov.  
     *palliatu* sp. nov.

*Hemiptychus*  
     two undescribed species  
*Cænocara californica* Lec.  
*Ptilinus basalis* Lec.  
     *ramicornis* Csy.  
     *flavipennis* Csy.  
*Sinoxylon sericans* Lec.  
     *sextuberculatum* Lec.  
     *declive* Lec.  
     *suturale* Horn.  
*Bostrychus californicus* Horn.  
*Amphicerus fortis* Lec.  
     *punctipennis* Lec.  
     *teres* Horn.  
*Dinapate wrightii* Horn.  
*Dinoderus pacificus* Csy.  
     *sobrinus* Csy.  
*Polycæon stoutii* Lec.  
     *confertus* Lec.  
     *megalops* sp. nov.  
*Psoa maculata* Lec.  
     *quadrisignata* Horn.  
*Lyctus planicollis* Lec.  
     *parvulus* Csy.  
     *californicus* Csy.

## CUPESIDÆ.

*Cupes lobiceps* Lec.

## CIOIDÆ.

*Cis versicolor* Csy.  
     *vitula* Csy.  
     *duplex* Csy.

*Plesiocis cribrum* Csy.  
*Ennearthron grossulum* Csy.  
     *convergens* Csy.

## SPHINDIDÆ.

*Odontosphindus clavicornis Csy.*

## LUCANIDÆ.

*Sinodendron rugosum Mann.*

## SCARABÆIDÆ.

- |                                       |                                       |
|---------------------------------------|---------------------------------------|
| <i>Canthon simplex Lec.</i>           | <i>Trox punctatus Germ.</i>           |
| var. <i>humeralis Horn.</i>           | <i>gemmulatus Horn.</i>               |
| <i>lævis Drury.</i>                   | <i>atrox Lec.</i>                     |
| <i>perplexus Lec.</i>                 | <i>Amphicoma ursina Lec.</i>          |
| <i>Copris mœchus Lec.</i>             | <i>edwardsii Horn.</i>                |
| <i>Ægialia conferta Horn.</i>         | <i>canina Horn.</i>                   |
| <i>latispina Lec.</i>                 | <i>Oncerus floralis Lec.</i>          |
| <i>crassa Lec.</i>                    | <i>Hoplia sackenii Lec.</i>           |
| <i>Psammodius nanus De Geer.</i>      | <i>callipyge Lec.</i>                 |
| <i>cælatus Lec.</i>                   | <i>pubicollis Lec.</i>                |
| <i>Rhyssemus californicus Horn.</i>   | <i>Gymnopyge hopliæformis Linell.</i> |
| <i>Atænius desertus Horn.</i>         | <i>Dichelonycha crotchii Horn.</i>    |
| <i>abditus Hald.</i>                  | <i>fuscata Lec.</i>                   |
| <i>gracilis Melsh.</i>                | <i>truncata Lec.</i>                  |
| <i>californicus Horn.</i>             | <i>pusilla Lec.</i>                   |
| <i>lobatus Horn.</i>                  | undescribed species                   |
| <i>Aphodius granarius Linn.</i>       | <i>Cœnonycha rotundata Lec.</i>       |
| <i>vittatus Say.</i>                  | <i>socialis Horn.</i>                 |
| <i>lividus Oliv.</i>                  | <i>Serica fimbriata Lec.</i>          |
| <i>rugifrons Horn.</i>                | <i>elongatula Horn.</i>               |
| <i>consociatus Horn.</i>              | <i>mixta Lec.</i>                     |
| <i>subænius Lec.</i>                  | <i>alternata Lec.</i>                 |
| <i>alternatus Horn.</i>               | <i>Plectrodes riversi Csy.</i>        |
| <i>cribratus</i>                      | <i>carpenteri Lec.</i>                |
| <i>ochreipennis Horn.</i>             | <i>palpalis Horn.</i>                 |
| <i>rubidus Lec.</i>                   | <i>squamosa Csy.</i>                  |
| <i>militaris Lec.</i>                 | <i>blaisdelli Csy.</i>                |
| <i>coquilletti Linell.</i>            | <i>fossiger Csy.</i>                  |
| <i>luxatus Horn.</i>                  | <i>pistoria Csy.</i>                  |
| <i>ungulatus sp. nov.</i>             | <i>Orsonyx anxius Lec.</i>            |
| <i>pardalis Lec.</i>                  | <i>Diplotaxis mœrens Lec.</i>         |
| <i>Ochodæus californicus Horn.</i>    | <i>subangulata Lec.</i>               |
| <i>Pachyplectrus lævis Lec.</i>       | <i>tenuis Lec.</i>                    |
| <i>Bradycinetus serratus Lec.</i>     | <i>corvina Lec.</i>                   |
| <i>Pleocomma puncticollis Rivers.</i> | undescribed species                   |
| <i>Trox suberosus Fab.</i>            | <i>Lachnosterna lenis Horn.</i>       |

<i>Listrochelus mucoreus Lec.</i>	<i>Cyclocephala hirta Lec.</i>
<i>Polyphylla decemlineata Say.</i>	<i>dimidiata Burm.</i>
<i>crinita Lec.</i>	<i>Ligyris gibbosus De G.</i>
<i>Thyce marginata Csy.</i>	<i>Euphoria verticalis Horn.</i>
<i>Phobetus comatus Lec.</i>	<i>Cremastochilus wheeleri Lec.</i>
<i>Anomala centralis Lec.</i>	<i>ineptus Horn.</i>
<i>Cotalpa ursina Horn.</i>	<i>schaumii Lec.</i>
<i>granicollis Hald.</i>	<i>westwoodi Horn.</i>
<i>Cyclocephala immaculata Oliv.</i>	<i>pilosicollis Horn.</i>
<i>longula Lec.</i>	<i>crinitus Lec.</i>
<i>villosa Burm.</i>	

## SPONDYLIDÆ.

<i>Parandra brunnea Fab.</i>	<i>Spondylis uniformis Mann.</i>
------------------------------	----------------------------------

## CERAMBYCIDÆ.

<i>Ergates spiculatus Lec.</i>	<i>Megobrium edwardsii Lec.</i>
<i>Mallodon melanopus Linn.</i>	<i>Molorchus longicollis Lec.</i>
<i>Derobrachus geminatus Lec.</i>	<i>Callimoxys fuscipennis Lec.</i>
<i>Prionus californicus Mots.</i>	<i>Rosalia funebris Mots.</i>
<i>Asemum nitidum Lec.</i>	<i>Dendrobias mandibularis Serv.</i>
<i>Criocephalus productus Lec.</i>	<i>Lissonotus multifasciatus Dup.</i>
<i>asperatus Lec.</i>	<i>Tragidion annulatum Lec.</i>
<i>Hylotrupes amethystinus Lec.</i>	<i>armatum Lec.</i>
<i>ligneus Fab.</i>	<i>Metaleptus angulatus Chev.</i>
<i>Phymatodes blandus Lec.</i>	<i>Amanuus pectoralis Lec.</i>
<i>obscurus Lec.</i>	<i>Batyle saturalis Say.</i>
<i>decussatus Lec.</i>	<i>Oxoplus jocosus Horn.</i>
<i>juglandis Leng.</i>	<i>Crossidius testaceus Lec.</i>
<i>Callidium antennatum Newm.</i>	<i>Ischnocnemis bivittatus Dup.</i>
<i>hirtellum Lec.</i>	<i>Stenosphenus debilis Horn.</i>
<i>Malacopterus lineatus Guer.</i>	<i>Cyllene antennatus White.</i>
<i>Æme gracilis Lec.</i>	<i>crinicornis Chev.</i>
<i>Eucrossus villicornis Lec.</i>	<i>Calloides lorquini Buq.</i>
<i>Brothylus gemmulatus Lec.</i>	<i>Clytus lanifer Lec.</i>
<i>Romaleum simplicicolle Hald.</i>	<i>Xylotrechus nauticus Mann.</i>
<i>seminitidum Horn.</i>	<i>obliteratus Lec.</i>
<i>Elaphidion albofasciatum Linell.</i>	<i>Neoclytus irroratus Lec.</i>
<i>imbelle Lec.</i>	<i>Atimia dorsalis Lec.</i>
<i>Anellus linearis Lec.</i>	<i>Desmocerus cribripennis Horn.</i>
<i>Pœcilobrium chalybæum</i>	<i>californicus Horn.</i>
<i>Hybodera debilis Lec.</i>	<i>Necydalis barbaræ Rivers.</i>
<i>Callinus cyanipennis Lec.</i>	<i>Ulochætes leoninus Lec.</i>
<i>ruficollis Lec.</i>	<i>Centrodera nevadica Lec.</i>



- Toxotus vestitus* *Hald.*  
*Pachyta spurca* *Lec.*  
*Anthophilax tenebrosus* *Lec.*  
*Acmaeops tumida* *Lec.*  
     *falsa* *Lec.*  
*Strangalia delicata* *Lec.*  
*Leptura molybdica* *Lec.*  
     *læta* *Lec.*  
     *tribalteata* *Lec.*  
     *coquilletti* *Linell.*  
     *instabilis* *Hald.*  
         var. *convexa* *Lec.*  
     *sexspilota* *Lec.*  
     *pernigra* *Linell.*  
     *crassipes* *Lec.*
- Leptura valida* *Lec.*  
*Ophistomis ventralis* *Horn.*  
*Ipochus fasciatus* *Lec.*  
*Monilema spoliatum* *Horn.*  
*Synaphæta guexi* *Lec.*  
*Cœnopæus palmeri* *Lec.*  
*Acanthocinus obliquus* *Lec.*  
     *spectabilis* *Lec.*  
*Pogonocherus crinitus* *Lec.*  
*Lypsimena californica* *Horn.*  
*Saperda mœsta* *Lec.*  
*Oberea schaumii* *Lec.*  
*Tetraopes femoratus* *Lec.*  
*Idœmea californica* sp. nov.  
*Methia* sp.

## CHRYSOMELIDÆ.

- Anulacoscelis purpurea* *Horn.*  
*Zengophora californica* *Cr.*  
*Lema nigrovittata* *Guér.*  
*Euryscopa subtilis* *Horn.*  
     *vittata* *Lec.*  
     *lecontei* *Cr.*  
*Coscinoptera canella* *Lec.*  
     *mucorea* *Lec.*  
     *æneipennis* *Lec.*  
*Babia 4-guttata* *Oliv.* var.  
*Saxinis saucia* *Lec.*  
     *politula* *Horn.*  
     *speculifera* *Horn.*  
     *hornii* sp. nov.  
*Chlamys* sp.  
*Exema conspersa* *Mann.*  
*Cryptocephalus sanguinicollis* *Suffr.*  
     *castaneus* *Lec.*  
     *spureus* *Lec.*  
*Pachybrachys analis* *Lec.*  
     *pubescens* *Oliv.*  
     *hybridus* *Suffr.*  
     *cœlatus* *Lec.*  
     *livens* *Lec.*  
     *lustrans* *Lec.*  
     ten or twelve undescribed species
- Diachus auratus* *Fab.*  
*Myochrous longulus* *Lec.*  
*Glyptoscelis illustris* *Cr.*  
     *sqamulatus* *Cr.*  
     *alternatus* *Cr.*  
*Colaspidea cuprascens* *Lec.*  
     *smaragdulus* *Lec.*  
     *varicolor* *Cr.*  
     *subvittata* *Fall.*  
*Typophorus viridicyaneus* *Cr.*  
     *canellus* *Fab.*  
*Metachroma californica* *Cr.*  
*Chrysochus cobaltinus* *Lec.*  
*Leptinotarsa behrensi* *Harold.*  
     *11-lineata* *Stål.*  
*Calligrapha*  
     *elegans* var. *californica* *Linell.*  
     *sigmoidea* *Lec.*  
*Plagioderia prasinella* *Lec.*  
*Gastroidea cyanea* *Melsh.*  
*Trirhabda geminata* *Horn.*  
     *caduca* *Horn.*  
     *luteocincta* *Lec.*  
     *flavolimbata* *Mann.*  
*Galerucella nymphaeæ* *Linn.*  
*Monoxia puncticollis* *Say.*



- Monoxia consputa* Lec.  
*debilis* Lec.  
*sordida* Lec.  
*Diabrotica* 12-punctata *Fab.*  
     var. *tenella* Lec.  
*soror* Lec.  
*trivittata* Mann.  
*Phyllobrotica viridipennis* Lec.  
*luperina* Lec.  
*ngripes* Horn.  
*Scelolyperus flavicollis* Lec.  
*maculicollis* Lec.  
*graptoderoides* Cr.  
*Trachyscelida bicolor* Lec.  
*Luperodes bivittatus* Lec.  
     *transitus* Horn.  
     *laticeps* Horn.  
     *torquatus* Lec.  
     *smaragdinus* Lec.  
     *morrisoni* Jac.  
     *varipes* Lec.  
*Malacorhinus maculatus* Lec.  
*Ædionychis violascens* Lec.  
*Disonycha pennsylvanica* Illig.  
     5-vittata Say.  
     *maritima* Mann.  
*Haltica bimarginata* Say.  
     *carinata* Germ.  
     *californica* Mann.  
     *æruginea* Lec.  
     *obolina* Lec.  
     *punctipennis* Lec.  
     undescribed species  
*Hemiglyptus basalis* Cr.  
*Crepidodera helxines* Linn.  
*Epitrix cucumeris* Harris.  
     *subcrinita* Lec.  
     *parvula* Fab.  
*Leptotrix recticollis* Lec.  
*Chætocnema opacula* Lec.  
     *subviridis* Lec.  
     *opulenta* Horn.  
     *ectypa* Horn.  
*Systema tæniata* Say.  
*Longitarsus repandus* Lec.  
     *livens* Lec.  
     *montivagus* Horn.  
     *mancus* Lec.  
     undescribed species  
*Glyptina cerina* Lec.  
     *atriventris* Horn.  
*Phyllotreta lepidula* Lec.  
     *vittata* Fab. (?)  
     *albionica* Lec.  
     *ramosa* Cr.  
     *pusilla* Horn.  
     undescribed species  
*Dibolia ovata* Lec. (?)  
*Psylliodes punctulata* Melsh.  
     *convexior* Lec.  
*Microrhopala rubrolineata* Mann.  
     *melsheimeri* Cr.  
*Odontota californica* Horn.  
*Stenopodius flavidus* Horn.  
*Cassida texana* Cr.  
*Coptocycla aurichalcea* Fab.

## BRUCHIDÆ.

- Bruchus pisorum* Linn.  
     *sordidus* Horn.  
     *ramicornis* Boh.  
     *limbatus* Horn.  
     *pruininus* Horn.  
     *aureolus* Horn.  
     *pauperculus* Lec.  
     *prosopis* Lec.  
*Bruchus protractus* Horn.  
     *uniformis* Horn.  
     *obtectus* Say.  
     *exiguus* Horn.  
     *seminulum* Horn.  
     three or four undescribed species  
*Zabrotes spectabilis*.  
     undescribed species

## TENEBRIONIDÆ.

- Craniotus pubescens Lec.*  
*Edrotes ventricosus Lec.*  
     *nitidus Csy.*  
*Triorophus punctatus Lec.*  
     *lævis Lec.*  
     *subpubescens Horn.*  
*Stibia ovipennis Horn.*  
     *maritima Csy.*  
*Auchmobius sublævis Lec.*  
*Eurymetopon rufipes Esch.*  
     *fusculum Csy.*  
     *convexicollis Lec.*  
     *sodalis Horn.*  
     *serratum Lec.*  
     *inflatum Lec.*  
*Emmenastus longulus Lec.*  
     *piceus Csy.*  
     *obesus Lec.*  
     *thoracicus Csy.*  
*Epitragus pruinosis Horn.*  
*Chilometopon abnorme Horn.*  
     *helopioides Horn.*  
*Cnemodus testaceus Horn.*  
*Zopherus tristis Lec.*  
     *granicollis Horn.*  
     *induratus Csy.*  
*Phlæodes diabolicus Lec.*  
*Noserus plicatus Lec.*  
*Dacoderus striaticeps Lec.*  
*Aræoschizus sulcicollis Horn.*  
     *costipennis Lec.*  
     *armatus Horn.*  
*Anepsius delicatulus Lec.*  
*Typhlusechus singularis Linell.*  
*Nyctoporis carinata Lec.*  
*Cryptoglossa verrucosa Lec.*  
     *lævis Lec.*  
*Centrioptera muricata Lec.*  
     *asperata Lec.*  
     *seriata Lec.*  
*Schizillus laticeps Horn.*  
*Microschatia inæqualis Lec.*  
     undescribed species
- Asida ægrota Lec.*  
     *actuosa Horn.*  
     *carinata Lec.*  
     *confluens Lec.*  
     *parallela Lec.*  
     *impetrata Horn.*  
     *obsoleta Lec.*  
     *muricatula Lec.*  
     *hirsuta Lec.*  
     *hispidula Lec.*  
     *luctata Horn.*  
     *angulata Lec.*  
     *marginata Lec.*  
     *gabbii Horn.*  
*Conioutis abdominalis Lec.*  
     *robusta Horn.*  
     *elliptica Csy.*  
     *lata Lec.*  
         var. *insularis Csy.*  
     *opaca Horn.*  
     *punctipes Csy.*  
     *subpubescens Lec.*  
     *globulina Csy.*  
     *pallidicornis Csy.*  
     *parviceps Csy.*  
*Cœlomorpha maritima Csy.*  
*Cœlus globosus Lec.*  
     *grossus Csy.*  
     *arenarius Csy.*  
     *latus Csy.*  
     *pacificus Fall.*  
     *remotus Fall.*  
*Eusattus robustus Lec.*  
     *politus Horn.*  
     *coquilletti Linell.*  
     *lævis Lec.*  
     *dubius Lec.*  
     *productus Lec.*  
     *convexus Lec.*  
     *difficilis Lec.*  
     *muricatus Lec.*  
*Eleodes quadricollis Esch.*  
     *armata Lec.*

- Eleodes femorata* *Lec.*  
*gentilis* *Lec.*  
*interrupta* *Blais.*  
*gracilis* *Lec.*  
*grandicollis* *Mann.*  
*gigantea* *Mann.*  
*acuticauda* *Lec.*  
*pilosa* *Horn.*  
*hirsuta* *Lec.*  
*scabripennis* *Lec.*  
*consobrina* *Lec.*  
*tenebrosa* *Horn.*  
*parvicollis* *Lec.*  
*subnitens* *Lec.*  
*marginata* *Esch.*  
*Embaphion depressum* *Lec.*  
*Trogloclerus costatus* *Lec.*  
*Eulabis grossa* *Lec.*  
*rufipes* *Esch.*  
*pubescens* *Lec.*  
*laticornis* *Csy.*  
*crassicornis* *Csy.*  
*obscura* *Lec.*  
*Cerenopus concolor* *Lec.*  
*costulatus* *Lec.*  
*Argoporis costipennis* *Lec.*  
*bicolor* *Lec.*  
*inconstans* *Horn.*  
*Amphidora littoralis* *Esch.*  
*nigropilosa* *Lec.*  
*Cratidus osculans* *Lec.*  
*fuscipilosus* *Csy.*  
*Stenotrichus rufipes* *Lec.*  
*Iphthimus*  
*serratus* var. *sublævis* *Bland.*  
*Cœloclenemis dilaticollis* *Mann.*  
*rugosa* *Linell.*  
*obesa* *Lec.*  
*Cibdelis blaschkii* *Mann.*  
*bachei* *Lec.*  
*lævigata* *Csy.*  
*Tenebrio obscurus* *Fab.*  
*tenebrioides* *Beauv.*  
*Bius estriatus* *Lec.*
- Doliema plana* *Fab.*  
*Alæphus pallidus* *Horn.*  
*Eupsophus cantaneus* *Horn.*  
*Mecysmus angustus* *Lec.*  
*tenuis* *Csy.*  
*Trichoton sordidum* *Lec.*  
*Ulus crassus* *Lec.*  
*latus* *Blais.*  
*Blapstinus longulus* *Lec.*  
*validus* *Csy.*  
*dilatatus* *Lec.*  
*histricus* *Csy.*  
*coronadensis* *Blais.*  
*rufipes* *Csy.*  
*æqualis* *Csy.*  
*funebri* *Csy.*  
*brevicollis* *Lec.*  
*pubescens* *Lec.*  
*sulcatus* *Lec.*  
*Conibius parallelus* *Lec.*  
*seriatus* *Lec.*  
*crassipes* *Csy.*  
*elongatus* *Horn.*  
*Notibius puberulus* *Lec.*  
*puncticollis* *Lec.*  
*granulatus* *Lec.*  
*gracilis* *Csy.*  
*sulcatus* *Lec.*  
*Cnemeplatia sericea* *Horn.*  
*Alaudes singularis* *Horn.*  
*Tribolium ferrugineum* *Fab.*  
*confusum* *Duv.*  
*Gnathocerus cornutus* *Fab.*  
*Echocerus maxillosus* *Fab.*  
*Ulosonia marginata* *Lec.*  
*Merotemnus elongatus* *Horn.*  
*Aphanotus brevicornis* *Lec.*  
*Alphitobius diaperinus* *Panz.*  
*piceus* *Oliv.*  
*Cynæus angustus* *Lec.*  
*depressus* *Horn.*  
*Metaclisa marginalis* *Horn.*  
*Uloma longula* *Lec.*  
*Phaleria rotundata* *Lec.*

<i>Phaleria limbata</i> Horn.	<i>Apocrypha clivinoides</i> Horn.
<i>debilis</i> Lec.	<i>Helops edwardsii</i> Horn.
<i>Anemia californica</i> Horn.	<i>ovipennis</i> Csy.
<i>Platydema subquadratum</i> Mots.	<i>rugicollis</i> Lec.
<i>oregonense</i> Lec.	<i>strigicollis</i> Horn.
<i>Phylethus bifasciatus</i> Say.	<i>attenuatus</i> Lec.
<i>Hypophlœus substriatus</i> Lec.	<i>bachei</i> Lec.
<i>opaculus</i> Lec.	<i>blaisdelli</i> Csy.
<i>Apocrypha anthicoides</i> Esch.	<i>discipula</i> Csy.
<i>dyschirioides</i> Lec.	

## CISTELIDÆ.

<i>Stenochidus gracilis</i> Lec.	<i>Hymenorus punctatissimus</i> Lec.
<i>cyanescens</i> Lec.	<i>Mycetochares longipennis</i> Csy.
<i>Hymenorus infuscatus</i> Csy.	<i>procera</i> Csy.
<i>grandicollis</i> Champ.	<i>pubipennis</i> Lec.
<i>fusicornis</i> Csy.	<i>Isomira variabilis</i> Horn.
<i>fuscus</i> Csy.	<i>luscitiosa</i> Csy.
<i>macer</i> Csy.	<i>Cistela opaca</i> Lec.

## OTHNIIDÆ.

*Othnius longicornis* Horn.

## MONOMMIDÆ.

*Hyporhagus gilensis* Horn.

## MELANDRYIDÆ.

<i>Carebara longula</i> Lec.	<i>Mycterus concolor</i> Lec.
<i>Nothus luteus</i> Horn.	<i>quadricollis</i> Horn.
<i>Laeconotus pinicolus</i> Horn.	

## PYTHIDÆ.

<i>Cononotus macer</i> Horn.	<i>Cononotus sericans</i> Lec.
------------------------------	--------------------------------

## CEDEMERIDÆ.

<i>Copidita quadrimaculata</i> Mots.	<i>Oxacis bicolor</i> Lec.
<i>Asclera nigra</i> Lec.	<i>sericea</i> Horn.
<i>excavata</i> Lec.	<i>lucana</i> Lec.
<i>Chrysanthia repanda</i> Horn.	<i>debilis</i> Horn.
<i>Oxacis fragilis</i> Horn.	<i>Rhinoplatia ruficollis</i> Horn.

## MORDELLIDÆ.

<i>Pentaria nubila</i> Lec.	<i>Anaspis atra</i> Lec.
<i>hirsuta</i> Smith.	<i>collaris</i> Lec.
<i>pusio</i> Lec.	undescribed species

*Mordella scutellaris* *Fab.*  
undescribed species

*Mordellistena vilis* *Lec.*  
*comata* *Lec.*  
*aspersa* *Smith.*

*Mordellistena tosta* *Lec.*  
*nubila* *Lec.*

*unicolor* *Lec.*  
many new or unidentified  
species

## ANTHICIDÆ.

*Eurygenius constrictus* *Lec.*  
*Stereopalpus incannus* *Csy.*  
*nimius* *Csy.*  
*pruinosis* *Lec.*  
*Corphyra distinguenda* *Horn.*  
*bardii* *Horn.*  
*inconspicua* *Horn.*

*Formicomus mundus* *Lec.*

*Anthicus tenuis* *Lec.*  
*pinalicus* *Csy.* (?)  
*canonicus* *Csy.*  
*bipartitus* *Csy.*  
*turgidicollis* *Csy.*  
*californicus* *Laf.*  
*confinis* *Lec.*  
*formicarius* *Goeze.*  
*rufulus* *Lec.*  
*cribratus* *Lec.*  
*hecate* *Csy.*  
*luteolus* *Lec.*  
*pinguescens* *Csy.*  
*ovicollis* *Csy.*  
*biguttulus* *Lec.*  
*punctulatus* *Lec.*  
*seminotatus* *Csy.*  
*nanus* *Lec.*

*Anthicus obliquus* *Csy.*  
*militaris* *Csy.*  
*bellulus* *Lec.*  
*herifuga* *Csy.*  
*maritimus* *Lec.*  
*corticalis* *Lec.*  
*obesus* *Csy.*  
*albicans* *Csy.*  
*parviceps* *Csy.*

*Tanarthrus salinus* *Lec.*  
*alutaceus* *Lec.*

*Notoxus spatulifer* *Csy.*  
*denudatus* *Horn.*  
*debilitans* *Csy.*  
*conformis* *Lec.*  
*sparsus* *Lec.*  
*constrictus* *Csy.*  
*robustus* *Csy.*  
*alamedæ* *Csy.*  
*cavicornis* *Lec.*  
*calcaratus* *Horn.*

*Mecynotarsus delicatulus* *Horn.*

*Xylophilus brunnipennis* *Lec.*  
*brunnescens* sp. nov.  
*nucleus* sp. nov.  
*constrictus* sp. nov.

## PYROCHROIDÆ.

*Pyrochroa californica* *Horn.*

## MELOIDÆ.

*Cysteodemus armatus* *Lec.*  
*Melœ barbarus* *Lec.*  
*strigulosus* *Mann.* (?)  
*Nemognatha apicalis* *Lec.*  
*lutea* *Lec.* (?)

*Nemognatha dubia* *Lec.*  
*nigripennis* *Lec.*  
*scutellaris* *Lec.*  
two undescribed species  
*Guathium nitidum* *Horn.*

- |   |   |
|---|---|
| <i>Zonitis flavida</i> <i>Lec.</i>          | <i>Cantharis incommoda</i> <i>Horn.</i> |
| undescribed species                         | <i>stygica</i> <i>Lec.</i>              |
| <i>Epicauta alphonsii</i> <i>Horn.</i>      | <i>auriculata</i> <i>Horn.</i>          |
| <i>puncticollis</i> <i>Mann.</i>            | <i>crotchii</i> <i>Horn.</i>            |
| <i>oblita</i> <i>Lec.</i>                   | <i>æneipennis</i> <i>Lec.</i>           |
| <i>straba</i> <i>Horn.</i>                  | <i>nitidicollis</i> <i>Lec.</i>         |
| <i>maculata</i> <i>Say.</i>                 | <i>lugens</i> <i>Lec.</i>               |
| sp. nov. near <i>lemniscata</i> <i>Fab.</i> | <i>compressicornis</i> <i>Horn.</i>     |
| <i>fallax</i> <i>Horn.</i>                  | <i>Calospasta elegans</i> <i>Lec.</i>   |
| <i>maura</i> <i>Lec.</i>                    | <i>perpulchra</i> <i>Horn.</i>          |
| <i>Cantharis melæna</i> <i>Lec.</i>         | <i>mirabilis</i> <i>Horn.</i>           |
| <i>magister</i> <i>Horn.</i>                | <i>mæsta</i> <i>Horn.</i>               |
| <i>vulnerata</i> <i>Lec.</i>                | <i>nemognathoides</i> <i>Horn.</i>      |
| <i>childii</i> <i>Lec.</i>                  | <i>opaca</i> <i>Horn.</i>               |
| <i>tenebrosa</i> <i>Lec.</i>                | undescribed species                     |
| <i>mærens</i> <i>Lec.</i>                   | <i>Tegrodera erosa</i> <i>Lec.</i>      |
| <i>insperata</i> <i>Horn.</i>               | <i>race latecincta</i> <i>Horn.</i>     |
| <i>occipitalis</i> <i>Horn.</i>             | <i>Phodaga alticeps</i> <i>Lec.</i>     |

## RHIPIPHORIDÆ.

- |  |   |
|--|---|
| <i>Rhipiphorus flavipennis</i> <i>Lec.</i> | <i>Rhipiphorus cruentus</i> var. <i>rufus</i> <i>Lec.</i> |
| <i>cruentus</i> <i>Germ.</i>               | <i>Myodites californicus</i> <i>Lec.</i>                  |

## RHYNCHITIDÆ.

- |                                       |                                       |
|---------------------------------------|---------------------------------------|
| <i>Auletes nasalis</i> <i>Lec.</i>    | <i>Rhynchites aureus</i> <i>Lec.</i>  |
| <i>latifrons</i> <i>Csy.</i>          | <i>æratoides</i> <i>Fall.</i>         |
| <i>Rhynchites bicolor</i> <i>Fab.</i> | <i>Deporaus glastinus</i> <i>Lec.</i> |

## OTIORHYNCHIDÆ.

- |  |   |
|--|---|
| <i>Trigonoscuta pilosa</i> <i>Mots.</i>    | <i>Paraptochus sellatus</i> <i>Boh.</i>     |
| <i>Eupagoderes argentatus</i> <i>Lec.</i>  | <i>Stenoptochus inconstans</i> <i>Csy.</i>  |
| <i>desertus</i> <i>Horn.</i>               | <i>Thricolepis inornata</i> <i>Horn.</i>    |
| <i>varius</i> <i>Lec.</i>                  | <i>simulator</i> <i>Horn</i> (?)            |
| <i>geminatus</i> <i>Horn.</i>              | <i>Geoderces puncticollis</i> <i>Csy.</i>   |
| <i>plumbeus</i> <i>Horn.</i>               | two undescribed species                     |
| <i>Rhigopsis effracta</i> <i>Lec.</i>      | <i>Aragnomus hispidulus</i> <i>Csy.</i>     |
| <i>Amotus longisternus</i> <i>Csy.</i>     | <i>Dysticheus insignis</i> <i>Horn.</i>     |
| <i>gracilior</i> <i>Csy.</i>               | <i>Eucyllus vagans</i> <i>Horn.</i>         |
| <i>Adaleres ovipennis</i> <i>Csy.</i>      | <i>Rhyodes dilatatus</i> <i>Horn.</i>       |
| <i>humeralis</i> <i>Csy.</i>               | <i>Dirotoognathus sordidus</i> <i>Horn.</i> |
| <i>Orimodema protracta</i> <i>Horn.</i>    | <i>Elissa constricta</i> <i>Csy.</i>        |
| <i>Mimetes setulosus</i> <i>Lec.</i>       | <i>Aramigus fulleri</i> <i>Horn.</i>        |
| <i>Sciopithes setosus</i> <i>Csy.</i> var. | <i>Scythopus californicus</i> <i>Horn.</i>  |

## CURCULIONIDÆ.

- Sitones sordidus* *Lec.*  
*crinitus* *Gyll.*  
*nebulosus* *Csy.*  
*hispidiceps* *Csy.*  
*explicitus* *Csy.*  
 undescribed species  
*Apion pennsylvanicum* *Boh.*  
*hesperum* *Fall.*  
*sordidum* *Smith.*  
*antennatum* *Smith.*  
*œdorhynchum* *Lec.*  
*opacicolle* *Smith.*  
*trogodytes* *Mann.*  
*proclive* *Lec.*  
*cribricolle* *Lec.*  
*cordatum* *Smith.*  
*varicorne* *Smith.*  
*ventricosum* *Lec.*  
*attenuatum* *Smith.*  
*Listronotus*  
 undescribed species  
*Emphyastes fucicola* *Mann.*  
*Lixus asper* *Lec.*  
*perforatus* *Lec.*  
*semivittatus* *Csy.*  
*Dinocleus pilosus* *Lec.*  
*jacobinus* *Csy.*  
*fartus* *Csy.*  
*albovestitus* *Csy.*  
*molitor* *Lec.*  
*wickhami* *Csy.*  
*Cleonus inornatus* *Lec.*  
*pacificus* sp. nov.  
*erysimi* sp. nov.  
*basalis* *Fall.*  
*lobigerinus* *Csy.*  
*vittatus* *Kirby.*  
*modestus* *Mann.*  
*pleuralis* *Lec.*  
*Dorytomus inæqualis* *Csy.*  
*hystriacula* *Csy.*  
*hirtus* *Lec.*  
*luridus* *Mann.*  
*Dorytomus rufus* *Say.*  
*Desmoris constrictus* *Say.*  
*Smicronyx pusillus* *Dietz.*  
*cinereus* *Mots.*  
*instabilis* *Csy.*  
*defricans* *Csy.*  
*nubilus* *Dietz.*  
*californicus* *Dietz.*  
*obtectus* *Lec.*  
 two or three undescribed species  
*Synertha imbricata* *Csy.*  
*Promecotarsus maritimus* *Csy.*  
*Phyllotrox nubifer* *Lec.*  
*Endalus limatulus* *Gyll.*  
*ovalis* *Lec.*  
*Stenopelmus rufinasus* *Gyll.*  
*Bagous californicus* *Lec.*  
*Phycocætes testaceus* *Lec.*  
*Otidocephalus vittatus* *Horn.*  
*Magdalis cuneiformis* *Horn.*  
*lecontei* *Horn.*  
 undescribed species  
*Macrorhoptus hispidus* *Dietz.*  
*estriatus* *Lec.*  
*Cionistes insolens* *Dietz.*  
*Anthonomus peninsularis* *Dietz.*  
*confusus* *Dietz.*  
*sycophanta* *Walsh.*  
*apertus* sp. nov.  
*albopilosus* *Dietz.*  
*æneolus* *Dietz.*  
*ater* *Lec.*  
*ochreopilosus* *Dietz.*  
*pauperculus* *Lec.*  
*helianthi* sp. nov.  
*ornatulus* *Dietz.*  
*figuratus* *Dietz.*  
*inermis* *Boh.*  
*stolatus* sp. nov.  
*Epimechus mimicus* *Dietz.*  
*soriculus* *Dietz.*  
*nevadicus* *Dietz.*  
*æmulus* sp. nov.



- Elleschus ephippiatus Say.*  
*Orchestes parvicollis Lec.*  
     *puberulus Boh.*  
*Tychius lineellus Lec.*  
     *prolixus Csy.*  
     *setosus Lec.*  
     species unidentified  
*Sibynes fulvus Lec.*  
*Paragoges maculatus Lec.*  
*Rhyssematus pubescens Horn.*  
*Piazurus californicus Lec.*  
*Copturus mammillatus Lec.*  
     *adpersus Lec.*  
     *longulus Lec.*  
     two undescribed species  
*Gyrotus munitus Csy.*  
*Ceutorhynchus subpubescens Lec.*  
     *angulatus Lec.*  
     *disturbatus Dietz.*  
     *obliquus Lec.*  
     *hornii Dietz.*  
     *nodipennis Dietz.*  
     *mutabilis Dietz.*  
*Ceutorhynchus convexicollis Lec.*  
     *albopilosus Dietz.*  
*Cœlogaster zimmermanni Gyll.*  
*Pelenomus cavifrons Lec.*  
*Baris heterodoxa sp. nov.*  
     *dilatata Csy.*  
     *montana sp. nov.*  
     *rubripes Csy.*  
     *futilis Csy.*  
*Onychobaris densa Lec.*  
     *depressa Csy.*  
     *austera Csy.*  
     *insidiosa Csy.*  
     *arguta Csy.*  
     *audax Csy.*  
*Trichobaris mucorea Lec.*  
     *compacta Csy.*  
*Limnobaris nasuta Lec.*  
     *seclusa Csy.*  
*Barinus difficilis Csy.*  
*Barilepton falciger Csy.*  
*Balaninus uniformis Lec.*

## CALANDRIDÆ.

- Scyphophorus acupunctatus Gyll.*  
     *robustior Horn.*  
     *yuccæ Horn.*  
*Cactophagus validus Lec.*  
*Rhodobænnus tredecimpunctatus Ill.*  
*Sphenophorus simplex Lec.*  
     *vomerinus Lec.*  
     *pictus Lec.*  
     *pertinax Oliv.*  
     *sayi Gyll.*  
     *placidus Say.*  
     *gentilis Lec.*  
*Sphenophorus tardus sp. nov.*  
*Calandra granaria Linn.*  
     *oryzæ Linn.*  
*Yuccaborus frontalis Lec.*  
*Cossonus crenatus Horn.*  
*Macrorhyncholus protractus Horn.*  
*Elassoptes marinus Horn.*  
*Rhyncholus dorsalis Lec.*  
     *angularis Lec.*  
     *oregonensis Horn.*  
     *spretus Csy.*

## SCOLYTIDÆ.

- Monarthrum scutellare Lec.*  
     *dentigerum Lec.*  
*Gnathotrichus retusus Lec.*  
*Pityophthorus pubipennis Lec.*  
     *carinulatus Lec.*  
*Pityophthorus confinis Lec.*  
     *digestus Lec.*  
*Hypothenemus striatus Lec.*  
     undescribed species  
*Xyleborus xylographus Say.*



<i>Xyleborus cælatus</i> Eich.	<i>Chætophlœus hystrix</i> Lec.
<i>Xylocleptes cucurbitæ</i> Lec.	<i>Carphoborus simplex</i> Lec.
<i>Tomicus confusus</i> Lec.	<i>Dendroctonus terebrans</i> Oliv.
<i>Micracis hirtella</i> Lec.	<i>frontalis</i> Zimm.
undescribed species	<i>Hylastes macer</i> Lec.
<i>Scolytus ventralis</i> Lec. (?)	<i>longus</i> Lec.
<i>Hylesinus aspericollis</i> Lec.	<i>nigrinus</i> Mann.
<i>Phlœosinus punctatus</i> Lec.	<i>Hylurgops subcostulatus</i> Mann.

## ANTHRIBIDÆ.

<i>Toxotropis approximatus</i> Lec.	<i>Brachytarsus</i>
	three species

## NOTES ON DISTRIBUTION OF SPECIES.

## CICINDELIDÆ.

**Omus.**—All our species of this genus occur in California, the headwaters of Kings and Tule Rivers in the Sierras (*levis*), Fort Tejon (*californicus*), and Port Harford along the coast (*lecontei*) being the southernmost recorded limits. Two species—*dejeani* and *audouni*—extend their range northward into Oregon and Washington. The species occur in hilly and mountainous regions only, so far as known.

**Tetracha.**—*T. carolina* occurs at Yuma, especially along the margins of pools, where they conceal themselves in the cracks or fissures of the drying mud banks.

**Cicindela.**—The species, with the exception of *12-guttata* and its varieties, are more or less local and only found near water, none occurring along paths or dry, sandy roads, as is true of several eastern species. *C. late-signata* and *C. sigmoidea* are confined to the ocean beaches, both being very abundant at San Diego in midsummer: *hirticollis* occurs on various beaches and also at Yuma:

*gabbii* is infrequent on the mud of salt marshes at San Diego and Wilmington in August and September, and is rarely taken on the ocean beach; Long Beach, August (Daggett): *senilis*, near San Diego, in damp places away from the ocean (Dunn): *vulgaris* and its variety *obliquata*, at San Luis Rey (Dunn); the latter also on the borders of the Big Laguna, Temecula Valley (Blaisdell): *vibex*, on the ocean beach at Oceanside (Blaisdell) and near San Bernardino (Wright): *12-guttata* and varieties, widely diffused along fresh water, Feb. to Oct. The most common form is that which Le Conte called *oregona*; these graduate into *guttifera* as Arizona is approached; typical *12-guttata* is seldom taken. The various forms are so perfectly connected that it seems scarcely worth while to continue the three names. The impression is quite general among coleopterists that *12-guttata* is not a variety of *repanda*, and it seems quite certain to me that it must be given full specific standing.

*C. pusilla* occurs in Owen's Valley: *tenuisignata*, Palm Springs and Colorado River: *sperata*, plentiful along the Colorado at Yuma in July: *lemniscata*, between San Diego and the Mexican boundary (fide Fuchs): *prætex-tata*, San Diego County (Fuchs). *C. hæmorrhagica* and *C. pacifica* are found along fresh-water streams at San Diego and San Bernardino; the former occurs on the sandy margins of salt flats back of the ocean beach at Santa Barbara in August and September, also at Yuma (Horn), while the latter has been taken by several collectors on the ocean front at Del Mar. *C. sommeri* is reported from the vicinity of San Diego.

The following species or varieties are recorded from the more northern portions of the State: *perviridis*, *graminea*, *10-notata*, *fulgida*, *cinctipennis*, *imperfecta*, *lunalonga*, *depressula*, *lauta*, and *plutonica*.

## CARABIDÆ.

**Omophron.**—*O. dentatum* is quite common in many localities along the sandy or gravelly margins of streams, Feb.-Aug. This species, together with *Bembidium*, *Tachys* and other littoral forms, is easily secured by "washing" the banks. *O. ovale* Horn and *concinnum*, *solidum* and *gemma* Csy. are northern.

**Trachypachys.**—*T. inermis* and *T. gibbsii* occur both in the northern Sierras and in the Coast Range. I have taken the latter at Lake Tahoe in July, and Mr. Daggett has found it at an elevation of 6,000 feet on the Kern River.

**Cychrus.**—*C. obliquus*, Los Angeles and San Diego; not common: *striatus*, Santa Barbara, Pomona, Pasadena; scarce along streams, where they conceal themselves under rocks and logs during the day: *minus*, somewhat more common than the last and occurs in same situations; best obtained by trapping with fresh meat: *punctatus*, San Diego County, common at Fort Tejon (Le Conte): *subtilis*, Los Angeles, very rare; not seen alive by the writer; the example in my collection is from Tulare County; it is also reported from Mariposa.

The following species also occur in the State: *angusticollis*, northern; *cordatus*, *cristatus*, *interruptus*, *ventricosus*, *striatopunctatus*, region about San Francisco; *oreophilus*, middle Sierras; *fuchsianus*, northern; *dissolutus*, "Cal." In a recent review of the subgenus *Brennus*, Casey has added the following names: *basalis*, *duplicatus*, *incipiens*, *decipiens*, *symmetricus*, *gentilis*, *strictus*, *convergens*, *sculptipennis*, *porcatus*, *sinuatus*, *politus*, *corpulentus*, *compositus*, *longipes*, *gravidus*, and *catenulatus*. The last two are said to be from Southern California.

Judging from the description, *gravidus* is not separable from *punctatus*, and *catenulatus* is without much doubt *minus*.

**Carabus.**—*C. tectatus* or variety occurs at Lake Tahoe and north.

**Calosoma.**—*C. angulatum*, "So. Cal."; not seen by the writer: *prominens*, Riverside, April; San Bernardino; Pasadena, Sept.; southern Mojave region (Horn, "The Coleoptera of Baja California")\*: *peregrinator*, "So. Cal." (Horn, l. c.): *parviceps*, San Diego County (Fuchs): *triste*, San Diego County: *semilave*, Redondo, February; Pomona, April-May; frequently common, roads and fields in spring and early summer: *simplex*, Los Angeles County, not common: *cancellatum*, Los Angeles; Redondo Beach, one example, March; Santa Monica, Santa Barbara: *latipenne*, Antelope Valley: *subaneum* occurs about Fresno: *discors* and *luxatum* var. in the north.

**Elaphrus.**—*E. viridis* is described from California without definite locality. Specimens of *riparius* from the Kern River, near Bakersfield, are in my collection, and it is possible that this may be found along some of the small streams on the desert side of the mountains. Casey describes *politus* from San Francisco.

**Loricera.**—Specimens of *L. californica* have been taken by Mr. F. S. Daggett (June) and by Mr. F. D. Twogood (August) along the margin of the Bear Valley reservoir in the San Bernardino Mountains, elevation 6,800 feet. *L. foveata* and *L. cærulescens* are northern.

**Notiophilus.**—*N. semiopacus*, one example, Pomona, February 18: *obscurus*, San Bernardino Mountains, 6,000 feet.

**Nebria.**—*N. eschscholtzii*, a small number found along a stream in a mountain cañon near Pomona in March.

---

\* Proc. Cal. Acad. Sci., 2nd Ser., Vol. IV, p. 307.

*N. rathvoni* occurs near Sacramento: *ingens* and *ovipennis*, in the "Sierra Nevada," probably middle or northern: *metallica*, in the extreme north.

**Metrius.**—*M. contractus* is said by Horn to occur in the Coast Range south of Tejon. *Promecognathus levissimus* is also said to occur in the same places as the preceding. I have never met either of these species in Southern California, and it is probable that they will be found only along the northern limits of the territory we are considering.

**Scarites.**—*S. subterraneus*, Riverside, April; Yuma, July.

**Dyschirius.**—*D. tridentatus*, *D. basalis* and *D. gibbipennis* are common near water almost everywhere west of the mountains: *patruelis*, San Diego (Le Conte): *ceneus* and *analis* are common along the Colorado River: *marinus*, common, and found only on the sea beaches: *truncatus* (?), rather rare; Pasadena, May; Lake Tahoe, July: *unipunctatus*, not rare at Pomona, Riverside and San Bernardino: *aratus* and *terminatus* are both recorded from California, the former was described from the Gila River and is common in southwestern Arizona; it may be expected to occur along the Colorado: *terminatus* occurs from New York to Kansas and its occurrence in California may be looked upon with suspicion: *consobrinus* is found at San Francisco and in the middle Sierras: *obesus*, on the northern beaches: *ceneolus* must be northern if its occurrence in the State is correctly reported.

**Clivina.**—*C. dentipes* was found along the Colorado River at Needles by Wickham, and by myself at Yuma: *punctulata* appears to be very scarce; a single example only at Pomona, June.

**Schizogenius.**—*S. crenulatus* and *S. pluripunctatus*, Colorado River: *seticollis*, common in many localities west of the mountains: *depressus*, generally distributed both east and west of the mountains: *litigiosus* occurs at San Francisco and north.

**Bembidium.**—Very numerous represented, more than one-fourth of all our described species occurring in Southern California. *B. transversale* (of this I believe *lugubre* to be only a variety), *striola*, *californicum*, *lucidum*, *variegatum*, *timidum*, *versicolor* and *iridescens* are widely distributed within our area and are generally common: *nitidum* has occurred at Riverside, March, and Pomona, May, but is rather rare: *erasum* occurs in the Sierras from 5,000 feet upward; the mentum tooth is not bifid in our specimens: *longulum*, *brevistriatum*, *consanguineum* and *acutifrons* are also found in the higher mountain cañons or valleys between the ranges: *carinatum*, banks of Colorado River: *bifossulatum*, Riverside, June, and near San Diego: *insulatum*, Riverside and Pomona, June; one example on ocean beach at Santa Barbara, August; both these species prefer the muddy banks of reservoirs or ponds in the low country, rarely or never occurring along the small streams of the mountain cañons or foot-hills: *recticolle*, Pasadena, Pomona; March to October; not rare along mountain streams at no great elevation: *platynoides*, in same situations as the last, but less frequent: *nubiculosum* and *intermedium* occur wherever there is water in the desert regions of the southeast, the former being exceedingly abundant: *indistinctum*, found only in the immediate vicinity of the coast, where it is everywhere common: *ephippiger*, Pomona, June; Redondo, April; Long Beach, August; more frequent near the coast on salt flats, but nowhere



common: *grandicolle*, San Diego, Riverside; rare: *vile*, San Diego and Los Angeles, August; rare: *horni*, two examples; Riverside, June; Sierras, July: *bifasciatum*, often abundant along mountain streams: *dubitans*, Pomona, Riverside, Pasadena; March to June: *trechiforme*, Santa Monica, February; Ojai Valley, March; rather scarce: *laticeps*, San Diego; very few specimens known: *cautum* (from which *connivens* is not separable) is widely diffused and very common: *henshawi* and *scudderi* are recorded from Tehachapi, the northern limit of our area.

The following species occur in the middle or northern portions of the State: *littorale*, *lorquini*, *sculpturatum*, *quadrulum*, *planatum*, *planiusculum*, *complanulum*, *juncireum*, *nigrocæruleum*, *incertum*, *nebraskense*, *nevadense*, *rickseckeri*, *approximatum*, *dentellum*, *variolosum*, *mormon*, *obtusangulum*, *wickhami*, *anguliferum*, *oblongulum*, *tigrinum*,\* *assimile*, *spectabile*. Hayward cites *musculicola* with an implied doubt: *sordidulum* was described by Chaudoir from California, but is unrecognized.

**Anillus.**—*A. explanatus* has been found in Alabaster Cave, El Dorado County: *debilis*, under stones at San Jose.

**Tachys.**—*T. vittiger* and *T. virgo* are found only near the seacoast; both have been found running on salt mud at Santa Barbara and Redondo, while *vittiger* has also been found at San Diego and near the beach on Catalina Island: *mordax*, Colorado River: *corax*, in same locality and more frequent, also at Pomona: *vorax*, rare at Pomona and Riverside: *edax*, Ojai Valley, March: *nanus*, common under bark of fallen trees in the mountains:

---

\*Specimens of *B. tigrinum* have been recently taken by me on the sandy margin of the salt flats back of the ocean beach at Santa Barbara (August). It has previously been known to me only from Alameda, probably occurring in similar situations.

*anthrax* and *audax*, common nearly everywhere along streams, the latter being our most common species: *rapax*, not rare at Yuma in July. Two examples of *rufotestaceus* have occurred at Pomona. I have seen a specimen of *flavicauda* from the northern part of the State. *T. falli* occurs in the middle and northern Sierras (Lake Tahoe and Siskiyou).

**Pericompsus.**—A few examples of *P. sellatus* taken along the banks of the Colorado River at Yuma in July.

Mr. Hayward writes me that *Pericompsus* will be merged into *Tachys* in his forthcoming revision.

**Patrobus.**—*P. californicus* is northern.

**Pogonus.**—*P. planatus* is reported from "Cal." (Horn), probably from the desert region of the southeast.

**Trechus.**—*T. barbaræ* occurs at Santa Barbara: *pomona*, rare, at Pomona and Pasadena in foot-hill cañons; October, January, March. *T. chalybeus* and *T. ovipennis* are both recorded from California; the former is northern, the latter middle and northern.

**Pterostichus.**—The first division of the genus, viz., those species without dorsal puncture on the elytra, is well represented, and several are yet undescribed. *P. ater* occurs in the mountains north of San Bernardino, and probably throughout the higher Sierras: *vicinus* is common in the Sierras and throughout the region to the west: *gracilior* appears to be confined to the mountains: *inermis*, not common, in the foot-hill cañons of the San Gabriel Range: *isabellæ* and *congestus*, common and widely diffused: *menetriesii*, Santa Rosa Island: *letulus*, one example from Santa Barbara: *scitulus*, not common: *subcordatus*, plentiful along the Colorado



River at Yuma: *hornii* and *splendidulus* are also reported from Yuma, but I have not seen specimens: *texanus*\* is reported from Needles by Wickham: *lustrans* and *occidentalis* are rather widely diffused, the former being common nearly everywhere, the latter more local and scarcer. I have seen *californicus* from "So. Cal.", but it is doubtless from the northern limits of this region. Aside from *menetriesii*, *isabellæ*, *latulus*, and an undetermined species are all that have thus far been taken on the islands off the coast. Of the other Californian species, *serripes* is from the Yosemite region; *tarsalis*, *planctus* and *protractus* are all fairly common at Lake Tahoe: *angustus*, middle coast region; *inanus*, middle and northern Sierras: *validus*, *brunneus*, *morionides*, *caligans* and *vitreus* are all northern. I have been unable to determine exact locality for *castanipes*; it is probably central or northern.

**Amara.**—*A. jacobinæ* has occurred at Riverside, June, but is more common in the immediate neighborhood of the coast—San Diego, Santa Monica, Redondo, etc.: *scitula*, rather uncommon, Pomona, September-December; San Bernardino Mountains, July: *longula*, one example only, Santa Monica, February: *insignis*, widely distributed from islands and beaches to mountains, but not very common: *insularis*, abundant in spring and early summer on all the islands: *impuncti-collis*, Pomona and Pasadena; a few specimens seen: *interstitialis*, Sierras, not frequent: *californica*, abundant everywhere: *remotestriata*, mountains only, not rare: *gibba*, "So. Cal." (Horn); not seen by writer: *aurata*, not rare in early spring, less frequent later in the season; widely diffused. *A. farcta* and *A. robustula* are recorded

---

\* Probably an incorrect identification of *T. subcordatus*.

from "Cal." without definite locality: *rectangula* and *imitatrix* are northern, the latter occurring also on the Farallon Islands (Fuchs): *stipida* is described from Sacramento; it is very near *jacobinae*: *conflata* was described from San Francisco. There is a specimen of *laticornis* in the Le Conte Collection bearing a gold disk, Le Conte's customary California label.

**Badister.**—*B. anthracinus*, a single specimen from Los Angeles County (Van Dyke): *ferrugineus*, "Alaska-Cal."; doubtless confined to the northern part of the State.

**Calathus.**—The name *C. ruficollis* is generally used for our very abundant species, and it is not quite plain how it is to be separated from *obscurus*, which, according to description, is a "So. Cal." species.\* Le Conte described *quadricollis* from San Francisco, and *behrensii* is from "Cal." It is impossible to separate these four species by Le Conte's table, and it is more than likely that all the names apply to a single moderately variable species.

**Pristonychus.**—*P. complanatus*, San Bernardino, about stables in December; San Diego, uncommon.

**Platynus.**—*P. agilis*, found at St. Isabel, San Diego County (Le Conte); I have never taken it: *simplex*, specimens so referred occur on the banks of the Colorado River: *frater* is recorded as from San Diego and San Francisco; it is unknown to me: *brunneomarginatus*, *californicus*, *funerarius*, *maculicollis* and *variolatus* occur nearly everywhere and are generally very common: *fossiger* and *deplanatus* are less frequent, the latter

---

\* In his list of species from Guadalupe Island (Trans. Am. Ent. Soc., Vol. V, p. 198), Dr. Horn refers a large species of *Calathus* to *C. obscurus*. The Guadalupe species appears to me to be distinct, and I believe Major Casey is justified in giving it the name *C. guadalupensis* in a recent description.

I have taken at Santa Barbara. In the vicinity of Los Angeles, *brunneomarginatus* and *funnebris* are to be found along the margin of natural water courses, *californicus* and *variolatus* on the shores of permanent ponds or reservoirs; the latter species is also found along irrigating ditches and in damp situations on cultivated land: *maculicollis* alone occurs about houses, where it is generally common under any object that will afford shelter during the winter and spring. I have occasionally seen great numbers on the wing in early spring evenings.

In addition to the above named species, the following are reported from California, and with the possible exception of *larvalis* are probably all from the central and northern portions of the State: *larvalis*, *jejunus*, *ovipennis*, *bicolor*, *ferruginosus*, *quadratus*, *cupripennis*, *sulcatus*, *striatus*. *P. punctiformis* is found along the Gila not far from the Colorado and will probably occur at Yuma.

**Perigona.**—*P. nigriceps*, Pasadena, at electric light, September; one specimen (Fényes).

**Lachnophorus.**—*L. elegantulus* is abundant along the sandy margins of streams.

**Euphorticus.**—*E. occidentalis* is of similar habits to the preceding, but quite rare; Riverside, late in May.

**Galerita.**—*G. lecontei*, not frequent; Riverside, May and June, at electric lights.

**Zuphium.**—*Z. longicolle*, north of the central portion of the State (Yuba County).

**Diaphorus.**—*D. tenuicollis*, Sacramento and San Jose.

**Thalpius.**—*T. hornii*, not rare along the banks of the Colorado: *rufulus* is reported from Southern California (Horn); it was described from San Jose.

**Ega.**—*E. letula*, found at East Bridge, on the Arizona side of the Colorado, by Mr. Wickham, and without a doubt occurs also on the California side; found also at Yuma.

**Tetragonodera.**—*T. fasciata*, margin of Colorado River; Yuma, July: *pallidus*, one specimen, taken under dried remains of *Opuntia*, Pomona, May; others under boards at Palm Springs in April; a very active insect and everywhere rare; so far as observed, it does not seem to share the littoral habits of its congeners.

**Lebia.**—*L. cyanipennis*, not rare: *ruficollis*, much less frequent; Riverside, April; Redondo, March: *viridis*, quite common in numerous localities; generally smaller than Eastern specimens: *furcata*, one example, Pomona, July: *guttula*, usually rare; occasionally found in small colonies under the loose bark of *Eucalyptus* in December and January: *bilineata*, San Bernardino Mountains, July; rare.

**Dromius.**—*D. piceus*, not rare under bark of live-oak in foot-hills near Pomona in February and April.

**Apristus.**—*A. laticollis*, rather common on banks of streams or ditches, and occasionally running on garden walks, usually in April and May.

**Blechrus.**—*B. glabratus* and *B. lucidus* occur in situations similar to the preceding, but are less common.

**Axinopalpus.**—*A. biplagiatus*, not rare under *Eucalyptus* bark in winter: *fusciceps*, only one specimen seen, Los Angeles County (Van Dyke).

**Tecnophilus.**—*T. croceicollis* is said by Horn to occur "from San Diego to Oregon," but I have not yet met it.

**Callida.**—*C. platynoides*, "Mountains east of Visalia" (Horn).

**Philophuga.**—*P. castanea*, one example, Pomona, May, from between the leaves of decaying yucca: *P. amæna*, middle California.

**Plochionus.**—*P. timidus* and *P. pallens* are in the Horn Collection, from "Cal.," without more definite locality.

**Pinacodera.**—*P. punctigera*, occurs at Yuma; other species are found in the Peninsula.

**Cymindis.**—*C. cribricollis*, San Diego County (Fuchs): *californica* and *Apenes limbata* were described by Horn ("Coleoptera of Baja California," Supp. I)\* as from San Luis Obispo. The occurrence of these genera in maritime California seems so surprising that one might almost suspect some confusion of localities. I have not yet seen specimens.

**Brachynus.**—The species are as elsewhere very indefinite. *B. lateralis* is readily known by the pale elytral margin; it is rare west of the mountains, but common along the Colorado River. One or two of the other species are abundant near water. *B. fidelis* is included on the authority of Mr. Fuchs.

**Chlænius.**—*C. ruficauda*, common along the Colorado from Needles to Yuma: *viridifrons*, widely distributed throughout the southern part of the State, but generally uncommon: *cursor*, Southern California (Horn); doubtless from the desert regions of the south-east: *cumatilis*, common along streams in foot-hill cañons: *leucoscelis*, occurs only along the Colorado, where it is not rare: *obsoletus*, not rare both east and west of the mountains: *variabilipes*, not common, though generally diffused: *glaucus*, Yuma, March (Daggett): *tricolor*, rather common everywhere: *harpalinus*, very

---

\* Proc. Cal. Acad. Sci., 2nd Ser., Vol. V, 1895, pp. 231, 232.

scarce, Los Angeles County (Van Dyke): *simillimus* occurs at San Francisco (Horn): *pennsylvanicus* in the north: *sericeus* is reported from the State without definite locality.

**Brachylobus.**—*B. caurinus* is northern (Yreka, Horn).

**Oodes.**—*O. elegans*, Colorado River, March (Daggett).

**Agonoderus.**—*A. lineola* moderately, and *A. pallipes* exceedingly common in many localities; both are frequently attracted in numbers to lights in early summer.

**Discoderus.**—A single example of *D. amœnus* was found under a stone near Palm Springs on the desert side of the mountains.

**Harpalus.**—This genus is not very well represented in California and only one species is known to me from the southern part of the State; this is superficially very much like *ventralis* and may be that species; specimens were taken in Bear Valley (elevation 6,000 to 7,000 feet) by Mr. Daggett. I have taken *caliginosus* at Sacramento, *cantus* at San Francisco, *somnulentus* at Lake Tahoe, and have specimens referred to *carbonatus* from the middle Sierras. There are two other apparently undescribed species in my collection from still further north.

**Stenolophus.**—*S. limbalis*, *S. flavipes* and *S. anceps* are common and widely distributed: *cineticollis* is described from the Colorado River: *alternans* from San Jose: *unicolor* is in the Horn Collection, from San Francisco, and others have been seen labeled simply "Cal".

**Bradycellus.**—*B. cognatus*, not common, Los Angeles County, June: *rupestris* and *californicus*, exceedingly abundant everywhere in moist places: *rivalis*, Colorado River and adjacent region.



**Tachycellus.**—Two forms are at present included under the name *T. nitidus*. The larger, pale form is more common near the coast, though I have seen specimens from as far inland as San Bernardino. The smaller, dark form is in my opinion specifically distinct; it is widely dispersed. There is a still smaller piceous form, the thorax with obtuse hind angles, specimens of which have been seen from Pasadena and from the southern Sierras. The proper elucidation of the species requires study.

**Anisodactylus.**—*A. dilatatus* is rather uncommon or at least local; I have taken it in early spring under stones in a dry hillside ravine near Pomona. *A. piceus*, *semipunctatus*, *consobrinus* and *californicus* are all widely distributed and more or less common. *A. amaroides* is rare in our region; I have specimens taken at Riverside and in the San Bernardino Mountains. Of the other species recorded from California, *strenuus* occurs at Fort Tejon; *obtusus* at San Jose; *pilosus* and *immanis* in the San Joaquin Valley; *nivalis* at Santa Rosa and north; *pitychrous* in the middle Sierras; *porosus* in the northeastern desert regions.

**Anisotarsus.**—*A. flebilis* or a closely allied species has been taken sparingly at Pomona.

**Pseudomorphus.**—*P. cronkhitei* occurs in Owens Valley; *behrensi* in the San Joaquin Valley.

#### AMPHIZOIDÆ.

**Amphizoa.**—Four examples of *A. insolens* were taken in a mountain stream near Pomona in September.

#### HALIPLIDÆ.

**Haliplus.**—*H. concolor* is rare at Pomona and Riverside, April and May; described from the Colorado River.

**Cnemidotus.**—*C. callosus* and *C. simplex* are both common and are to be found at all seasons. I have taken both most frequently in algæ occurring on stagnant pools in August and September.

#### DYTISCIDÆ.

**Laccophilus.**—*L. decipiens*, everywhere plentiful: *terminalis* is moderately common, and *4-lineatus* less so, at Yuma: *mexicanus* has occurred twice at Pomona: *fasciatus* is said by Crotch to occur in California, but I have not seen specimens.

**Hydrovatus.**—To *H. brevipes* is referred a single example taken at Long Beach.

**Desmopachria.**—*D. latissima*, San Diego, Santa Barbara; not common.

**Bidessus.**—*B. cinctellus*, "So. Cal." (*teste* Horn), probably from the desert region, as it is known to be a common species in adjacent parts of Arizona: *pictodes*, generally scarce; I have seen specimens from Ventura only; an allied form is not rare in the San Bernardino Mountains: *affinis*, occurs commonly in most localities; the form taken west of the mountains is the dark variety to which Le Conte gave the name *obscorellus*; specimens from east of the mountains are more or less distinctly vittate: *subtilis*, to this should probably be referred the commonest species in maritime Southern California, though few specimens are as distinctly marked as the Le Conte type: *amandus*, reported from "So. Cal." by Horn ("The Coleoptera of Baja California")\* without definite locality; it was described from the Gila River, and should occur along the Colorado; in the type the sutural stria is very feeble, but I suspect that this may be merely an individual variation.

---

\* Proc. Cal. Acad. Sci., 2nd Ser., Vol. IV, 1894, p. 313.



**Hygrotus.**—*H. hydropicus*, rather rare, Pomona and San Diego.

**Cœlambus.**—*C. medialis*, rather common everywhere: *fraternus*, Colorado River: *pedalis*, not common, Riverside and Pomona, May and June: *lutescens* is quite common in middle California,—San Francisco, San Jose, Sacramento: *impressopunctatus*, one specimen, Lake Tahoe (Fényes).

**Deronectes.**—*D. griseostriatus*, not common, Redondo, December and February: *striatellus*, exceedingly abundant everywhere.

**Hydroporus.**—*H. addendus*, rare at Pomona and Ojai Valley, more common a little further north: *fortis*, rare, San Diego; *subpubescens*, not common, though widely diffused: *axillaris*, southeastern region, probably Yuma: *vilis* and *latebrosus*, common in most localities. The following species are also reported from the State and some may possibly occur within our limits, though the species of the genus are as a rule more northern in habitat: *eximius*, *rivalis* (= *sanmarki* Sahlb, according to Fauvel), *terminalis* and *hardyi*. Mr. Fuchs has a record of *H. funereus* from San Diego. This is quite likely correct, but needs verification.

**Coptotomus.**—*C. interrogatus* is common at Needles in August (Wickham).

**Hydrotrupes.**—*H. palpalis* is one of our rarest species, occurring in the mountain streams of the southern Sierras.

**Ilybiosoma.**—*I. regularis* is plentiful everywhere.

**Agabinus.**—*A. glabrellus*, rather scarce as a rule, but once found in some numbers in a small spring on Catalina Island.

**Agabus.**—*A. lugens*, abundant everywhere: *lineellus*, very rare, Pomona, April and May: *morosus*, Pomona,

San Diego; not common: *obsoletus*, San Diego (Le Conte); *lecontei*, rather common, various localities. The following species are found farther north: *intersectus*, *brevicollis*, *confertus*, *strigulosus*, *tristis* (Lake Tahoe), *suturalis*, *erichsoni*.

**Rhantus.**—*R. binotatus*, common generally: *anisonychus*, much rarer, Pomona: *flavogriseus* and *sinuatus* are reported from the State, but I have not seen specimens from the south.

**Colymbetes.**—*C. strigatus*, rare; a single specimen taken at San Bernardino in April: *sculptilis* is recorded from the State by Crotch: *seminiger* occurs in the northeast.

**Eretes.**—*E. sticticus* is a cosmopolitan species which has occurred once at Soldiers' Home (Van Dyke).

**Hydaticus.**—*H. stagnalis* is recorded from "Cal." (Horn, "The Coleoptera of Baja California"\*).

**Dytiscus.**—*D. marginicollis*, scarce; several specimens taken by Mr. Daggett in Bear Valley, elevation 7,000 feet, June; one example taken by myself at Santa Monica.

**Thermonectes.**—*T. basilaris* var. *intermedius*, one example, Pomona, September: *marmoratus*, near San Diego (Fuchs).

**Cybister.**—*C. explanatus*, not rare, many localities: *ellipticus*, Colorado River. Both species are frequently attracted by electric lights.

## GYRINIDÆ.

This family is very poorly represented in Southern California.

---

\* Proc. Cal. Acad. Sci., 2nd Ser., Vol. IV, 1894, p. 314.

**Gyrinus.**—*G. plicifer* occurs in the small streams of the mountains and foot-hills, while *consobrinus* generally prefers larger and more permanent waters; the former is not rare about San Francisco, more especially in the small streams of the redwood region, a few miles north of the Bay; the latter extends its range into Oregon. I have seen hundreds of *consobrinus* about the electric lights at Riverside, in May. Le Conte mentions having two examples of *affinis* from middle California, and I have a species of doubtful identity from the middle Sierras.

**Gyretes.**—*G. sinuatus*, occurs at Yuma (Le Conte).

#### HYDROPHILIDÆ.

**Helophorus.**—*H. obscurus*, common and widely diffused: species doubtful, one example, Ventura, March: *fortis*, San Francisco (Le Conte): *angustulus*, northern.

**Hydrochus.**—Undescribed species, Ojai Valley, March; San Bernardino Mountains, July; rare: *variolatus*, San Diego (Le Conte): *vagus*, Colorado River.

**Ochthebius.**—The species occur sometimes in great numbers in the mud or silt or under stones in the shallow water of the margins of streams or pools, where any agitation of the bottom brings them to the surface, upon which they float and are easily taken. *O. rectus* is our commonest species and occurs nearly everywhere; I have taken it plentifully in muddy irrigating ditches at Riverside, in clear streams, and at San Diego on a salt marsh which is covered by tide-water: *interruptus*, Riverside, San Diego, Redondo; common at the latter place, living in algæ in a small lake much saltier than the ocean: *discretus*, widely diffused and moderately common: *puncticollis*, rare, San Bernardino Mountains, Ojai Valley: *sculptus*, rare, Ventura: *costipennis*, one

example, Ventura, March: *holmbergi*, Santa Monica and San Diego, not very common: *lineatus*, abundant at Yuma, July: *nitidus*, Yuma (Le Conte). Dr. Horn records *laevipennis* from Fort Tejon, and I have taken *cribricollis* at San Francisco and Lake Tahoe.

**Hydræna.**—*H. pennsylvanica*, common everywhere in same situations as *Ochthebius*.

**Hydrophilus.**—*H. triangularis*, not rare, comes quite often to electric lights: *insularis*, said by Horn ("The Coleoptera of Baja California")\* to occur in Southern California.

**Tropisternus.**—All the species common and generally distributed except *salsamentus*, which has occurred only in the Redondo salt lake, April to June, and in salt pools at Santa Barbara, August.

**Hydrocharis.**—*H. glaucus*, three examples taken by Mr. F. D. Twogood in a watering trough at a lumber camp in the San Bernardino Mountains, elevation 5,000 feet, September; found also on Santa Cruz and Santa Rosa islands, but everywhere rare; and by Dr. Blaisdell in Calaveras Mountains, Mokelumne Hill and on Mt. Diablo (Fuchs): *obtusatus*, one example in mountains near Pomona, elevation 4,000 feet, August: *rickseckeri* occurs near Santa Rosa (Ricksecker).

**Berosus.**—*B. punctatissimus*, not rare, Pomona to San Diego: *miles*, plentiful at Yuma: *subsignatus*, Colorado River: *salinus*, Redondo salt lake, not rare in early spring; found also in salt pools at Santa Barbara in August: *infuscatus*, Yuma, San Diego, Redondo, Pomona, etc., rather common: *rugulosus*, very rare, only two examples found, Pomona, June and September.

**Chætarthria.**—All three species are found about equally common by washing the sandy margins of streams.

---

\* Proc. Cal. Acad. Sci., 2nd Ser., Vol. IV, 1894, p. 315.

**Limnebius.**—*L. piceus*, common enough when found, but inclined to be local; I have found it in numbers in the streams of the foot-hills near Pomona and Pasadena; *politus*, *alutaceus* and *congener* are names given by Casey to forms occurring from San Francisco north; they are, I think, based merely on local or individual and sexual variations of *piceus*.

**Laccobius.**—*L. ellipticus* is extremely abundant everywhere.

**Philhydrus.**—*P. carinatus* and *P. nebulosus* are common and widely diffused: *diffusus*, not rare in the Redondo salt lake, also in salt marsh at Santa Barbara: *cuspidatus* and *hamiltoni* are recorded from the northern part of the State. Specimens taken by myself at Sacramento are by their color to be referred to *hamiltoni*; this difference, however, is of little consequence, and they are structurally not materially different from *diffusus* of Southern California. *P. californicus* is from an unrecorded part of the State, "probably northern": *conjunctus* occurs at Lake Tahoe.

**Helochares.**—*H. normatus* is found in most localities, but nowhere very common.

**Cymbiodyta.**—*C. punctatostriata*, found only in mountain streams: *dorsalis*, common and generally distributed: *imbellis* occurs from Fort Tejon north, and is not rare in the neighborhood of San Francisco.

**Hydrobius.**—*H. fuscipes*, not common, Pomona, Riverside, San Bernardino: *latus*, extreme north: *scabrosus*, San Francisco and north.

**Creniphilus.**—*C. subcupreus*, not very frequent; Pomona, San Bernardino Mountains, Yuma: *elegans*,\* not common, in the salt lake at Redondo: *rufiventris*, rare at

---

\* Specimens of this species recently received from Mr. Schwarz were taken by the late Henry G. Hubbard in salt springs at Salton on the Colorado Desert.

low altitudes, but more plentiful in the higher parts of the Sierras: *infuscatus*, common: *dissimilis*, San Francisco and north.

**Dactylosternum.**—*D. cacti*, one example, Los Angeles County (Van Dyke); occurs abundantly in decomposing cactus (*Cereus*) in Arizona, but is thus far rare in Southern California.

**Cercyon.**—*C. fimbriatus*, everywhere abundant along the seashore in decomposing seaweed: *luniger*, much less frequent in same situations, and on Catalina Island, where *fimbriatus* has not yet been found: *fulvipennis*, common in excrementitious matter: *lugubris*, not rare, about the roots of grass in marshy places, and in decaying vegetable matter: *nigriceps*, rare, San Diego, Pomona.

*C. depressus*, *quisquilius*, *lateralis*, *tristis* and *navicularis* are quoted from California in Horn's Revision.

**Megasternum** —*M. posticatum*, rather common in decomposing vegetable matter, especially along the mountain streams.

#### SILPHIDÆ.

**Necrophorus.**—*N. marginatus*, rare; one specimen taken at Riverside, June: *guttula*, numerous on San Clemente Island, May and June, rather scarce on the mainland: *pustulatus* var. *nigritus*, not rare on both islands and mainland; numerous specimens taken in electric light trap at Pasadena, April and May; var. *melsheimeri* is northern.

**Silpha.**—*S. lapponica* and *S. ramosa* are generally common: *opaca*, one specimen mentioned by Dr. Horn as taken near Mono Lake.

**Necrophilus.**—*N. hydrophiloides*, from middle and northern part of State.



**Agyrtes.**—One example of *longulus* taken at Pomona under live-oak chips, February 15, 1896; one example also taken by Dr. Fényes at Pasadena, January, 1898.

**Sphærites.**—*S. glabratus* is northern.

**Pinodytes.**—One dead and mutilated specimen of *cryptophagoides* (?) found with ants at Pasadena (Fényes).

**Platycholeus.**—Several specimens of *leptinoides* were taken by me in very rotten wood at Lake Tahoe, July; the species is termitophilous (Koebele, Hubbard, Fényes).

**Choleva.**—*C. basillaris* from middle and northern part of State.

**Ptomaphagus.**—*P. consobrinus*, Pomona, San Bernardino Mountains, uncommon; one specimen found at Pomona with ants in March: *californicus*, "San Diego northward" (Horn): *pusio*, northern: *flsus*, Owens Valley.

**Colon.**—*C. clavatum* and *C. inerme* are both northern.

**Triarthron.**—*T. lecontei*, "high Sierras east of Visalia."

**Hydnobius.**—*H. latidens*, one example, so referred by Dr. Horn, Ojai Valley, March: *longulus*, northern.

**Anisotoma.**—Specimens of this and the preceding genus are rare, and are most often taken on the wing an hour or two before sunset on warm days in early spring: *paludicola* has thus occurred at Pomona and Pasadena, January, February: *obsoleta* (?), Ojai Valley, March: *humeralis*, one example taken by Dr. Fényes in leaves beneath chopped meat which had been placed as a bait for *Cychnus*, Pasadena, January: *curvata* and *collaris* are northern: *difficilis* is found in Owens Valley.

**Liodes.**—One specimen of *confusa* was found at Lake Tahoe (Fényes).

**Cyrtusa.**—A single specimen of *picipennis* was taken by myself in the San Bernardino Mountains, August.

**Agathidium.**—*A. revolvens*, rare, San Bernardino Mountains, July: *concinnum*, Pomona, May, not common, more frequent farther north: *virile*, found rarely at Pomona and Riverside in early spring: *pulchrum*, taken on decaying mushrooms at Pomona, February: *californicum* and *sexstriatum* occur in the middle and northern parts of the State.

**Clambus.**—One example of a species near *seminulum*, but smaller; Pomona, April.

#### SCYDMÆNIDÆ.

This family is poorly represented in California, only seventeen species being known from the entire State, and of these but six have been found in Southern California; moreover all of these six are of greater or less rarity, a state of affairs surely not due entirely to the small size of the specimens or lack of proper investigation.

**Connophron.**—*C. occidens* and *C. digressum* are represented, the first by a single specimen taken at or near San Bernardino by Mr. W. G. Wright, and now in the collection of Captain Casey, the latter by three specimens taken by myself about the roots of grass in a marshy locality near Pomona.

**Scydmaenus ovipennis.**—A single specimen has been taken at Pasadena by Dr. Fényes in February. The three other species listed are doubtless all members of the Sonoran fauna. *V. colon* has occurred in San Diego County, and *C. deformata* is said to have been collected by Crotch at Los Angeles.

**Papusus macer.**—This was described from a unique example taken by me under a stone at Palm Springs, on



the western border of the Colorado desert, in April. Captain Casey writes me that the type which was sent to him with other specimens for study has been lost. The remaining Californian species with their localities are as follows: *Lophioderus gracilis*, San Jose; *L. myops*, San Francisco; *Drastophus levicollis*, San Francisco; *Scydmenus sparsus*, San Jose; *S. pacificus*, San Francisco and north; *S. fuchsi*, San Francisco; *S. californicus*, northern; *S. catalinus angustus*, San Jose; *Cephennium anophthalmicum*, San Francisco; *Veraphis impressa*, Lake Tahoe; *Scydmenus ovithorax* Bndl., Santa Clara County. According to Casey's recent revision, the last-named species would not be a true *Scydmenus*, and would perhaps require the erection of a new genus for its reception.

#### PSELAPHIDÆ.

**Adranes.**—An undescribed species found by Mr. Hubbard in a colony of *Lasius rubripes* at Lake Tahoe.

**Articerus fuchsi** var. **californicus.**—Occurs at Los Angeles in March (Fuchs); specimens were taken near Los Angeles by Mr. Koebele, in the nests of *Cremastogaster lineolata*.

**Biotus.**—*B. formicarius*, found living in the nests of a pale brown ant, Los Angeles (Casey).

**Sognorus.**—*S. pulvereus*, San Jose (Le Conte).

**Ctenisis.**—Several specimens of *dispar* were taken at an electric light, Pasadena, April (Fényes).

**Tyrus.**—*T. corticinus* is not uncommon under bark of fallen conifers at Lake Tahoe.

**Valda.**—*V. frontalis*, Siskiyou County (Casey).

**Tychus.**—A single example of the tiny species *T. tenellus* was taken on the wing at Pomona in January: *hexagonus*, three examples also taken flying, on a warm day in March, in the Ojai Valley; and six examples taken in like manner by Dr. Fényes at Pasadena in February: *puberulus* occurs at San Jose (Le Conte): *sonomæ*, Mendocino County.

**Pselaptrichus.**—*P. tuberculipalpus* was taken in the neighborhood of San Francisco in some numbers by Mr. Fuchs.

**Scalenarthrus.**—*S. hornii* was found at Needles, on the Colorado River, by Mr. Wickham, and by myself further down the river at Yuma, in July, under vegetable debris accumulating in damp situations along the bank of the stream.

**Decarthron.**—*D. brendeli*, Riverside and Pomona, April, in damp places near streams.

**Pselaptus.**—*P. belfragei* is described from Texas, but is said to occur as far west as Yuma (Casey).

**Reichenbachia.**—*R. deformata* and *R. tumidicornis* are both easily obtained in numbers by sifting along the margins of streams and in moist places nearly everywhere: *sagax* is far rarer, having been taken by myself once at Santa Monica in early spring, and by Dr. Fényes at Pasadena by sifting, March: the types of *falli* and *turgidicornis* were both taken by me under dead fish on the ocean beach at Santa Monica in February, 1892, and no other specimens were obtained until March, 1897, when both species were again found by Dr. Fényes and myself on the ocean beach at Redondo. It is probable that the occurrence of these species on the ocean front is due to winds or other fortuitous

circumstances. The following species are also Californian: *informis*, *fundata*, *tumorosa*, and *fusticornis* from the north; *compar* and *taphrocera* from the middle region; and *nevadensis* from western Nevada and the adjacent parts of California.

**Bryaxis.**—*B. foveata*, Yuma, San Diego, Riverside, Pasadena; not common: *loripes*, Southern California (Casey).

**Batrisus.**—*B. zephyrinus*, *B. monticola* and *B. cicatricosus* are from the middle Sierras: *occiduus* and *denticauda* from the north: *pygidiulis* is described from "Cal."

**Rhexidius.**—*R. granolus* and *R. asperulus* both occur in the vicinity of San Francisco.

**Oropus.**—Two examples of a species near, if not identical with, *abbreviatus*; Pasadena, March, by sifting (Fényes). The other species, *striatus*, *convexus*, *montanus*, *interruptus* and *cavicauda*, are all from the middle or northern parts of the State.

**Morius.**—*M. occidens*, Santa Cruz County (Casey).

**Sagola.**—*S. isabellæ*, not rare in moist situations, especially along the small streams of the foot-hill cañons: *grandiceps*, *rubida* and *longicollis* are from the middle, and *cavifrons*, *corticina* and *subsimilis* from the northern portions of the State.

**Euplectus.**—*E. orbiceps*, Los Angeles County (Casey): *californicus*, not common, Pomona, Pasadena, Ojai Valley; by sifting; March to June.

**Actium.**—*A. californicum* occurs in same situations as the preceding, and in about equal numbers: *candidum*, *marinicum* and *brevipenne* are described from middle California, and *politum*, *robustum*, *testaceum* and *pacificum* from farther north.

## STAPHYLINIDÆ.

The members of the Aleocharini are very numerous, but for the most part unstudied, and are consequently much neglected by most collectors.

**Falagria.**—*F. leviuscula* is very common along the margin of streams in the foot-hills: an undescribed species occurred in the San Bernardino Mountains (elevation 5,000 feet): *cavipennis* was described by Le Conte from San Pedro: *occidua* and *laticollis* are described from a little south of San Francisco (Santa Clara County).

**Echidnoglossa.**—Several species occur in the wet moss in mountain cañons; some of these are probably the same as some of the following species described by Casey from the middle and northern portions of the State, viz., *valida*, *eximia*, *brevicornis*, *gracilis*, *grandicollis*.

**Hoplandria.**—A species referred with some doubt to this genus was taken in the Ojai Valley in March.

**Pontomalota.**—*P. opaca* is usually plentiful on the sands of ocean beaches: *californica* and *nigriceps* are closely allied forms found at San Francisco and Santa Cruz.

**Colpodota.**—*C. parva* is said by Fauvel to occur at Mariposa.

**Atheta.**—This genus will contain many of the species hitherto lumped under *Homalota*. The species are numerous, and while no special effort has been made to collect them there are in my collection some twenty or thirty species from Southern California. *A. picipennis*, Mann, and *oraria* and *coriaria*, Kraatz, are mentioned by Hamilton and Fauvel as having been found in California.

**Lomechusa.**—*L. angusta* has occurred twice, the first specimen having been found with ants in the interior of an oak gall at Pasadena (Fényes), and the other under rubbish at Redondo beach (Daggett): *montana* was originally described from a specimen found under a stone at Truckee, near the summit of the middle Sierras; it has since been taken by Dr. Fényes under bark with ants, at Lake Tahoe, and farther north at Castle Crag in like manner.

**Tarphiota.**—*T. pallidipes* and *T. fucicola* are common on sea beaches.

**Tachyusa.**—Four species occur more or less commonly along streams, where they may be taken in numbers by "washing" the banks. It is not yet known whether they are identical with any of the following middle and northern species: *experta*, *linearis*, *laticeps*, *faceta*, *harfordi*, *crebrepunctata*.

**Tinotus.**—Two examples of *caviceps* were taken at Pomona in January.

**Myrmedonia.**—*M. sonomæ* is described from Sonoma County, and *M. fauveli* is said by Casey to occur near Los Angeles.

**Anepsiota.**—*A. insignis*, San Francisco.

**Phlæopora.**—Two species so referred were found at Pomona and San Diego.

**Calodera.**—*C. attenuata*, middle California (Monterey and Napa counties).

**Polystoma.**—*P. arenaria*, not very abundant at Redondo beach, San Diego, Santa Monica, and the islands off the coast. I have seen one specimen from

San Bernardino, but all others are from the immediate vicinity of the coast. *P. pacifica* is described from Santa Barbara.

**Microglossa.**—Fauvel mentions the occurrence of *suturalis* in "California."

**Aleochara.**—*A. sulcicollis* occurs only on the beaches, where it is quite abundant; several other species are generally common in excrementitious matter. One example of what seems surely identical with the European *puberula* has been taken at Pomona.

**Maseochara.**—*M. valida*, found at Pasadena, April to July, under decaying fruits and vegetables: *puberula*, one example at Arrowhead Hot Springs, May 31.

**Thiasophila.**—Several specimens of a species taken at Pomona early in September: *asperata* occurs at Lake Tahoe and Truckee.

**Isoglossa.**—A fine species, doubtfully referred to this genus, occurred in the San Bernardino Mountains in July: *arcuata* was described from Lake Tahoe.

**Oxyptoda.**—About five species of this genus are known to me from Southern California. Casey describes *fustiger* from Humboldt County, and *californica* from "Cal."

**Myrmecochara.**—An undescribed species from Lake Tahoe (Schwarz).

**Autalia.**—*A. elegans*, Lake County.

**Thinusa.**—*T. maritima* is described from Oakland. I have found a closely allied if not identical species on the beach at Santa Barbara.

**Silusa.**—*S. vesperis*, Humboldt County.

**Bolitochara.**—*B. californica*, common throughout the State in or about fungi: *nigrina* is found near San Francisco and in Humboldt County.



**Leptusa.**—A single specimen of a species taken at Pomona.

**Oligota.**—A minute species is sometimes beaten abundantly from the branches of trees; it is quite likely not separable from *pusillima*.

**Sipalia.**—*S. frontalis*, "California."

**Bryobiota.**—*B. bicolor* is a very small linear species, occurring rather plentifully on the ocean beach at San Diego and Santa Barbara.

**Somatium.**—*S. oviforme* is occasionally found in some numbers at Pomona, usually by beating.

**Gyronecha.**—*G. obscura*, rather common at times along the banks of streams, Pomona; a smaller form found in the Ojai Valley seems different. *G. attenuata* occurs at Monterey and in Napa County.

**Bamona.**—*B. falliana* is not rare at Pomona along the margin of streams.

**Myllæna.**—One example of a species taken in Los Angeles County.

**Acylophorus.**—*A. pronus*, taken only in small numbers at Pomona, January, and Palm Springs in April.

**Heterothops.**—*H. fumatus* var. *californicus* and *H. carbonatus* occur not uncommonly near springs or streams: *pusio* is less frequent in like situations: *occidentis* was taken deep in the detritus of hollow oak stumps at Pomona, February.

**Quedius.**—The species occur for the most part under stones or drift along the margins of streams in the foothills and mountains, though *fulgidus* is likely to be found almost anywhere. *Q. capucinus* and *Q. limbifer* are



most often taken: *explanatus* and *erythrogaster* are rather scarce: *desertus* is found east of the mountains: *prostans* is rather common in wet moss, at considerable elevations, but is only rarely taken in the lower valleys: *debilis* and *seriatus* are northern; of the latter I secured several examples near San Francisco (Marin County).

**Thinopinus.**—*T. pictus* is not rare on various beaches from San Diego northward; specimens in my collection were taken at Redondo, February; Coronado, June; Santa Monica, August.

**Creophilus.**—*C. villosus*, wide-spread, but usually not abundant; very common on San Clemente Island in May and June, about the carcasses of birds and foxes shot by hunters.

**Hadrotes.**—*H. crassus* is taken on sea beaches; Redondo, March to May; Santa Barbara, February; San Diego; not numerous.

**Staphylinus.**—*S. saphyrinus*, not uncommon under stones near water, especially in the foot-hills: *luteipes*, less common; Pomona, January; Santa Barbara, February: *tarsalis*, quite rare, Pomona and Los Angeles: *rutilicauda*, found in the middle Sierras: *nigrellus*, found farther north: *cinnamopterus* is said by Horn to occur from Maine to California; I have not yet seen specimens.

**Belonuchus.**—*B. ephippiatus* was taken abundantly by Dr. Fényes and myself under rotting squash, etc., Pasadena, Pomona, etc., July to October. Specimens of an apparently undescribed species were taken in decaying yucca, in mountains near Pomona, May.

**Philonthus.**—I have found *nitescens* only along the wet margin of a small salt lake or pond at Redondo, April: *alutaceus*, taken sparingly in a low, marshy spot

near Pomona, November to January: *semiruber*, rare; two examples only taken along a stream in foot-hills near Pomona: *hepaticus*, very scarce, same locality as preceding: *flavolimbatus*, has occurred at Pasadena in November, but is very rare west of the mountains; not rare at Yuma: *grandicollis*, not common; my specimens were taken in the bed of the Los Angeles River, near the city, in August: *longicornis*, widely distributed, but not very common: *discoideus*, rare at Pomona about stables: *alumnus*, the form with red thorax and elytra is very common at Yuma, on the river shore: *pettiti* and *triangulum*, not rare, many localities: *ferreipennis*, rather infrequent, Pomona, June to September: *sordidus*, common about houses and stables: *versutus*, two specimens only, San Bernardino Mountains, elevation 5,000 feet: *virilis*, not rare, same locality as the last: *nigritulus*, everywhere abundant: *instabilis*, has occurred on one occasion at Mojave: *quadrulus*, not infrequent in the foot-hills near Pomona: *lecontei*, not common, but widely dispersed. In addition to the above the following occur within the State: *politus* and *albionicus*, from San Francisco, north; *clunalis*, at Alameda; *puberulus*, from Tulare, north; *theveneti*, *crotchi*, *punctatellus* and *picicornis*, in the middle Sierras and Lake Tahoe region; *decipiens*, *bidentatus*, *siegwaldi*, *bucephalus*, *varians* and *caurinus*, from the north; *furvus* and *ventralis*, from "Cal.," but probably not southern. There are in my collection several probably undescribed species, for the most part represented by females only.

**Actobius.**—*A. ocreatus*, not uncommon at Los Angeles, Pomona, etc.: *sobrinus*, one specimen, Fort Yuma (Horn): *puncticeps*, abundant everywhere in the foot-hills and mountains; rare in the valleys: *gratus*, widely diffused, but nowhere common: *paederoides*, very common

everywhere from the sea-level to moderate altitudes, both east and west of the mountains: *formosus*, not common, Pomona and Pasadena: *elegantulus*, usually not plentiful, but at times found in numbers, Pasadena and Pomona. All the species are found only in the vicinity of water. The following occur in middle California: *senilis*, *infinus*, and *semipunctatus*—the latter now identified with *Bisnius procerulus* Grav.

**Cafius.**—The species of this genus are confined to the seacoast, and are often found in great numbers in or beneath decaying seaweed, or other littoral rubbish. *C. canescens* is our most abundant species, but *seminitens*, *lithocharinus*, and *luteipennis* often occur nearly as plentifully: *decipiens* is said to occur rather commonly at San Diego: *sulcicollis* and *opacus* are rather rare. All the species, except *seminitens* and *decipiens*, have been reported from one or another of the islands.

**Xantholinus.**—*X. cephalus*, not rare in the Sierras, under bark: *picipennis*, common in fields and pastures beneath stones, cow-droppings, etc.: *dimidiatus*, "San Bernardino, Mojave Desert," Pomona, May; in decaying yucca; rare: *obscurus*, not rare in damp situations near Pomona, September to February: *nanus*, San Diego; not seen by the writer: *pusillus*, not frequent, Riverside and Pomona.

**Leptacinus.**—*L. brunnescens*, common in most localities under any object upon the ground, especially from November to May: *pallidulus*, one specimen, Fort Yuma (Le Conte). There are three undescribed species from Southern California in my collection, of which one, at least, is not rare. *L. grandiceps* was described from San Francisco and *parvus* from San Jose; I have taken the latter at Sacramento.

**Othius.**—*O. californicus*, Mann, "Alaska and California." I have seen specimens taken at Stewart's Point (Ricksecker), and at Monterey (Fényes).

**Baptolinus.**—*B. punctiventris* is not uncommon under bark at Lake Tahoe; found also by Mr. Daggett, on Mt. Whitney, at an elevation of 11,000 feet.

**Stenus.**—*S. renifer*, *S. incultus* and *S. gila* occur on the Colorado River, the two latter being moderately common in Yuma early in July: *sculptilis* and *californicus* are very abundant along mountain streams, the former also occurring occasionally in the valleys: *costalis*, less common in same situations: *terricola*, one example, San Bernardino Mountains: *zunicus* and *pacificus* are found everywhere in the lower country, the former common at times, the latter always abundant: *insignis*, not rare at Pomona, November to May: *vestalis*, one example, Riverside: *sayi*, found in some numbers about the roots of grass in marshy spot near Pomona, January; not seen elsewhere: *pinguis*, common in many localities, occurring in prodigious numbers at times; hundreds have been found on every square yard of ground about a small lake near Pasadena, great numbers flying about and swarming on the shrubbery about the lake at the same time: *letulus*, rather rare, Ventura, Riverside, Pomona; March to May: *alveolatus*, one example, so named, San Diego: *lucidus*, a single example taken at Pomona, May.

The following additional species are recorded from the State: *tristis*, *luctuosus*, *luculentus* and *vacuus* from the middle regions; *corvus* from Fort Tejon; *dives* and *subgriseus* from the north; *ellipticus*, *colonus*, *villosus*, *exilis*, and *pollens* from "California" without definite locality.

**Cryptobium.**—This genus is not well represented in California, *californicum* being the only species known from west of the mountains; it is not rare, especially in the foot-hills and mountains. *C. tumidum* is recorded from California, Arizona and Utah, and is doubtless confined to the region east of the mountains. I have taken females of an undetermined species at Yuma.

**Lathrobium.**—*L. puncticeps*, rather rare: *jacobinum*, very common along any of the streams from the Sierras: *californicum*, rare, Pomona, May; Redondo beach, March: *lituarium*, a few specimens taken at Indio and Yuma: *subseriatum* occurs in the northern part of the State. There are three undescribed species in my collection, one from Southern California (Pasadena), and the other two from the north.

**Caloderma.**—*C. rugosa*, *C. mobilis* and *C. reducta* are all abundant under vegetable debris near water, occurring sometimes in great numbers in damp places in the mountain cañons: *continens* is recorded from San Diego; it is probably not different from *rugosa*. Three species not yet determined have been found in the cañons near Pomona.

The other species described by Casey—*angulata*, *contracta*, *luculenta* and *tantilla*—occur in the middle and northern parts of the State.

**Oligopterus.**—*O. cuneicollis*, San Francisco.

**Medon.**—*M. maluca* occurs in the same situations as the species of *Caloderma*, but specimens are not quite so numerous: *latuiscula*, "Southern California and Lake County" (Casey). Several unidentified species are from Southern California, while the following are from middle and northern California: *sinuaticollis*, *convergens*,

*lepida*, *puberula*, *sublesta*, *consanguinea*, *contigua*, *luctuosa*, *retrusa*, *gregalis*, *minula*, *languida*.

**Lithocharis.**—*L. ochracea*, not common, Pomona in June, flying at twilight: *alutacea* is described from Santa Clara County: *quadricollis*, from Lake County.

**Stilicus.**—*S. occiduus*, one example, Los Angeles, August: *quadriceps* and *opaculus* are described from "Cal."

**Pæderus.**—*P. femoralis*, "So. Cal." (teste Horn),\* doubtless from the Colorado River region: *compotens*, common in the vegetable detritus on the shore of Lake Wilson at Pasadena, May: *ustus*, abundant along the banks of the Colorado at Yuma.

**Sunius.**—*S. californicus*, not common: *longiusculus*, frequent. Both are quite widely distributed.

**Scopæus.**—*S. truncaticeps*, a single example, Pomona, June, so determined by Casey: *armiger*, a single specimen, Pomona, December.

**Scopæodera.**—A few examples of *nitida* were taken at Yuma in July.

**Leptorus.**—*L. texanus*, Colorado River, Yuma: *californicus*, Palm Springs, western border of Colorado Desert: *longipennis*, one specimen, Pomona.

**Orus.**—*O. punctatus*, common nearly everywhere west of the mountains: *fraternus*, not rare at Pomona and San Bernardino: *montanus*, two examples, male and female, San Bernardino Mountains: *parallelus*, northern (Napa and Sonoma counties).

**Pinophilus.**—*P. densus*, valley of Colorado River.

**Palaminus.**—*P. lividus*, taken on leaves of *Salix*, Yuma (Le Conte).

---

\* "The Coleoptera of Baja California," Proc. Cal. Acad. Sci., 2nd Ser., Vol. IV, 1894, p. 319.



**Tachinus.**—*T. agilis* and *T. debilis* are both rather widely distributed and moderately common. The genus is better represented farther north, *semirufus* and *angustatus* occurring in the middle regions, and *tachyporoides*, *pallipes* and *instabilis* in the upper portions of the State. A female specimen taken at Lake Tahoe is referred somewhat doubtfully to *memnonius*.

**Tachyporus.**—*T. californicus*, exceedingly abundant nearly everywhere; often seen flying at twilight in spring and early summer: *natidulus*, rare, Pomona, Lake Tahoe.

**Cilea.**—One example of *C. silphoides* from Pasadena, taken in February (Fényes).

**Erchomus.**—A single specimen of *E. punctipennis* was taken in the San Bernardino Mountains.

**Conosoma.**—*C. bipustulatum*, Wilmington and Santa Barbara (Horn); I have an example from the middle Sierras: *castaneum*, not rare in or under decaying wood, especially in the foot-hills of the Sierras; fungivorous according to Horn.

**Boletobius.**—*B. cincticollis* is common in mushrooms.

**Bryoporus.**—*B. rufescens* is said to occur from "Pennsylvania to California;" I have not seen California specimens. A single specimen of a possibly undescribed species has been taken at Pomona.

**Mycetoporus.**—*M. humidus*, "Michigan to Florida and west to California;" I have not yet seen it here: *splendidus* or a closely allied species has been found in the Santa Cruz Mountains.

**Habrocerus.**—*H. tarsalis* is described from San Mateo.

**Olisthærus.**—*O. megacephalus*, included on the authority of Fauvel; it is unquestionably northern.



**Pseudopsis.**—*P. obliterated*, *P. detrita* and *P. minuta* occur not uncommonly, and occasionally in large numbers, in the vegetable detritus along our mountain streams; the first named species was described from the valley of the Gila.

**Oxyporus.**—Specimens of an undescribed species have been taken at Lake Tahoe in Nevada by Mr. Schwarz, and on the California side of the lake by Dr. Fényes.

**Bledius.**—The species are numerous, and are found for the most part burrowing in wet sand or mud on the margins of fresh-water streams and ponds. *B. diagonalis* is recorded by Le Conte as occurring on salt marsh at San Diego: *cribricollis* appears to be confined to the sea-coast; I have seen it in abundance on the beach at Santa Barbara in February, but it is likely to be plentiful at all seasons: *ferratus*, *jacobinus*, *flavipennis* and *eximius* are all reported from San Diego, but I do not know them in nature: *opacifrons*, San Diego and Los Angeles: *luteipennis*, San Bernardino: *armatus*, Riverside, Pomona, Long Beach; not common: *nitidiceps*, Los Angeles, Wilmington (Le Conte), Pomona, one example in nearly dry bed of brook, November: *relictus*, two examples, Pomona: *laticollis*, common, many localities: *rusticus*, rare, Pomona: *ruficornis*, abundant everywhere: *clurus*, Riverside, Pomona, not common; *pleuralis*, occasionally common; once taken in numbers in San Dimas Cañon near Pomona, April: *ornatus*, not rare near the sea-coast, rare inland; taken in numbers at Redondo, February: *forcipatus*, Yuma (Le Conte): *punctatissimus* and *phytosinus* are recorded from "Southern California" without definite locality. A single specimen of a small black species not yet determined has been taken at Pomona. The following occur further

north: *deceptivus*, sp. nov, Kern County; of Casey's species, *strenuus* and *monstratus* occur at San Francisco, *lectus*, *foraminosus*, *gentilis*, *gracilis* and *bicolor* somewhat farther north, *parvicollis* in Mendocino County, *monticola* at Lake Tahoe, and *villosus* from "California".

**Platystethus.**—*P. americanus* occurs chiefly about cow-droppings, both east and west of the mountains.

**Oxytelus.**—*O. sculptus*, not rare about stables, etc.: *sobrinus*, one example, Pomona, December: *laqueatus*, *nitidulus*, *vegrandis*, and *niger* are found near San Francisco and farther north, the last-named species and *montanus* occurring at Lake Tahoe.

**Haploderus.**—*H. flavipennis* and *H. cephalotes* are both often taken in the vegetable detritus which collects in damp situations along streams or ponds: *linearis* is quite surely more northern in habitat: *annectens*, reported from San Francisco and Sonoma County.

**Trogophlæus.**—*T. gila* and *T. pacificus* are common under rubbish about gardens and along water courses at Pomona: *diffusus* occurs along streams in the higher parts of the Sierras: *prominens*, one example, Los Angeles (Casey): *confinis*, *debilis*, and *bledimus* are described from San Diego: *pauperculus* and *filum*, from Yuma: *armatus*, *occiduus*, and *lithocharinus* are from the middle coast region: *lapsus* and *obliquus*, from the middle Sierras (Truckee and Lake Tahoe): *sculptilis*, from Sonoma and Lake counties: *insolitus*, from Hoopa Valley, Humboldt County. Wickham reports *gila*, *dentiger*, and *tantillus* from Yuma. There are several unidentified species in my collection, some of which may be undescribed.

**Apocellus.**—*A. analis* and *A. gracilicornis* are both rather common near water, especially about reservoirs, irrigating ditches., etc., in cultivated regions: *sphaericollis*, one example, Pomona, April.

**Ancyrophorus.**—A small colony of *A. planus* was once taken in drift in a mountain cañon near Pomona.

**Thinobius.**—*T. hesperius*, Yuma (Casey): two unidentified species from Southern California, one of which is quite surely undescribed, and the other is probably *oxytelinus*. The following species are described from farther north: *macropterus*, "various localities" (Le Conte); *sonomæ*, Sonoma County; *gracilicornis*, Santa Clara to Sonoma counties: *validus*, Marin County and north.

**Zalobius.**—*Z. serricollis*, common under rubbish near water, more especially in the foot-hill and mountain cañons: *spinicollis*, middle and northern parts of the State; found in situations similar to the last.

**Asemobius.**—*A. calatus*, "California" (Horn, "Coleoptera of Baja California," Supp. I.).\*

**Deleaster.**—*D. concolor*, from San Francisco and north.

**Geodromicus.**—*G. temporalis*, quite rare along mountain streams, near Pomona, September: *humboldtianus*, northern.

**Tilea.**—*T. cavicollis*, *T. rufitarsis* and *T. filicornis* occur in the middle and northern Sierras.

**Vellica.**—*V. longipennis*, from Middle Sierras and north.

**Artochia.**—*A. productifrons*, from Santa Clara County.

**Unamis.**—*U. truncata*, from Middle Sierras.

**Lesteva.**—*L. fusconigra*, from Eldorado County.

---

\* Proc. Cal. Acad. Sci., 2nd Ser., Vol. V, 1895, p. 238.

**Brathinus.**—*B. californicus* was found in some numbers by Mr. Hubbard at Lake Tahoe, where it occurred in wet moss along a mountain stream; other specimens were found at Sisson.

**Amphichroum.**—*A. floribundum*, very abundant in many localities, especially on flowers of *Ceanothus*, in April: *puberulum*, described from San Diego; specimens taken in the San Bernardino Mountains are so referred with some doubt. The species are more numerous in the middle Sierras, and from there north to Vancouver. The following are recorded from the State: *sparsum*, *scutatum*, *opaculum*, *flavescens*, *pallidum* and *maculatum* from the middle Sierras (Mariposa and Tahoe); *bino-tatum*, *pilosellum*, *alutaceum*, *modestum*, from the region about and to the north of San Francisco; *testaceum* and *crassicornum* from still further north.

**Tanyrhinus.**—*T. singularis* occurs in the Santa Cruz Mountains, but is quite rare.

**Lathrimæum.**—One example of *L. subcostatum* obtained in the mountain cañon near Pomona, March. The following are recorded: *pictum*, northern; *nigropiceum*, Santa Cruz County; *spretum*, Siskiyou; *atrocephalum*, Tahoe (Fauvel).

**Homalium.**—This genus is numerously represented in California, about half our listed species being recorded from the State. *H. plagiatum* has occurred in mushrooms, at Pomona, in early spring, and *theveneti* at Santa Barbara and Redondo, on the ocean beach, in March; but, aside from these, I am able to find definite records of only four species from Southern California, viz.: *strigipenne*, San Diego; *repandum*, "Southern California and Texas" (Fauvel); *humile*, San Bernardino Mountains, identified by Casey; *alutaceum*, San Diego (Fauvel). When the

whole State is considered, we must add the following: *algarum* and *rugipenne*, from San Francisco; *ater*, from Santa Cruz; *pacificum*, from Siskiyou; *longulum*, *pusillum*, *megarthroides*, *rivulare*, *lapponicum*, *cæsum* and *fractum*, from "California," without indication of locality. It is probable that few, if any, of these latter will be found within our region. Several species in my collection from Southern California are as yet unidentified.

**Anthobium.**—*A. atriventre* and *A. nigerrimum* occur on flowers, the former very abundantly at Pomona, in April. Two other unidentified—possibly undescribed—species are known to me from Pomona, Ojai Valley and Palm Springs. The species of this genus are somewhat numerous, but not identifiable as the literature now stands. The following are to be added to the above: *aurifluum*, *diversicolle* and *subangulatum*, from Lake Tahoe; *gilvipenne* and *punctatum*, Santa Cruz County; *fraternum*, northern; *californicum*, "California."

**Orobanus.**—*O. rufipes*, San Diego and Pomona; one example from the bank of a mountain stream near Pasadena.

**Protinus.**—Two species of this genus have been found on decaying mushrooms at Pomona, in early spring; neither species has as yet been identified. In the mountains of the middle coast region occur *limbatus*, *basalis* and *salebrosus*: *mäklini* is middle or northern; *sulcatus* is described from the middle Sierras.

**Megarthrus.**—*M. pictus*, reported from middle and northern portion of the State.

**Lispinus.**—*L. californicus*, not rare in many localities, under bark: *linearis*, less frequent, Riverside,

Pomona and Palm Springs—at the last-named locality found under cottonwood bark.

**Trigonurus.**—All our species occur from middle California north, under bark of conifers. *T. edwardsii* has been taken in Los Angeles County (fide Fuchs).

**Micropeplus.**—*M. punctatus*, occurs at San Diego (Crotch): a single example of a probably undescribed species was taken at Pasadena, in March, by sifting (Fényes): *tesserula* and *brunneus* are more northern.

### TRICHOPTERYGIDÆ.

These minute things have not been systematically searched for, and the number recorded from Southern California, and indeed from the whole State, is very inconsiderable.

**Actidium.**—Four species are on record: *robustum*, *granulosum* and *attenuatum* from the middle coast region (Santa Cruz), and *politum* from “California.”

**Motschulskium.**—*M. sinuatocolle* occurs plentifully at times on various beaches from San Diego northward.

**Ptilium.**—An unidentified species has been taken sparingly at Pomona: *columbianum*, *humile* and *collani* are said to be from “California.”

**Ptenidium.**—*P. pullum*, quite common in vegetable detritus near water at Pomona, etc.

**Trichopteryx.**—*T. laticollis*, “Southern California,” in Horn Collection: *californicus* and *crotchii* bear “California” labels in the Le Conte Collection; and *sitkaensis* bears a gilt locality label which is supposed to signify a like origin.

**Smicrus.**—*S. filicornis*, from Colorado River; Le Conte Collection.



## HYDROSCAPHIDÆ.

**Hydroscapha.**—*H. natans* is probably not a rare insect, though very seldom taken; when one is discovered a considerable number may usually be obtained by patient search. I have taken it close to the bank in a mountain stream, and also in a muddy spot in a cow pasture near Pomona; Crotch took the first examples in the Los Angeles River.

## SPHÆRIIDÆ.

**Sphærius.**—*S. politus* is rather common on the sandy margins of streams, and is often started in numbers by "washing" the banks.

## SCAPHIDIIDÆ.

**Scaphisoma.**—*S. castaneum*, not very common, found in and about decaying and fungus-grown wood in damp or wet places: *rufulum*, occurs at Yuma (Le Conte).

## PHALACRIDÆ.

The species of this family are for the most part common on flowers; they are difficult to identify with certainty, and have been generally neglected by collectors.

**Phalacrus.**—*P. ovalis*, common at Pomona, Riverside, etc.: of *penicillatus* (so named by Casey) a number of specimens were taken in the San Bernardino Mountains: *conjunctus*, same locality as the last; also San Diego (Casey). The species here called *ovalis* is possibly not that species; it was pronounced "n. sp." by Casey, to whom specimens were sent for identification, but a comparison with Le Conte's types shows scarcely any tangible



difference; in fact, the three species here named are separated by characters of very doubtful significance.

**Olibrus.**—*O. wickhami*, sp. nov., Southern California (Casey); many specimens collected in San Diego County by Dunn.

**Acylomus** —*A. nebulosus*, taken at Yuma, in March (Daggett).

**Eustilbus.**—*E. apicalis*, widely distributed and rather common; I have beaten it in some numbers from a very tall dried grass at Pasadena, in February: *obtusus* and *nanulus* both occurred plentifully about the roots of grass in a marshy spot near Pomona, in January: *notabilis* occurred sparingly in the same situation: *aquatilis*, described from San Jose.

### CORYLOPHIDÆ.

**Sacium.**—*S. amabile*, found under bark of sycamores, sometimes in large numbers. I took on one occasion over seven hundred examples from beneath the bark of a fallen sycamore in the Ojai Valley, early in March: *decolor*, rare, Pomona: *scitulum*, described from Yuma (Le Conte).

**Sericoderus.**—*S. flavidus* is found in debris near water, Riverside, Pomona, etc.

**Orthoperus.**—Two apparently undescribed species have been taken; one of these was found plentifully about grass roots, in moist situations, in winter; of the other there is but a single example from the Ojai Valley.

**Ænigmaticum.**—*Æ. californicum*, not rare, Santa Barbara, Ventura, San Diego; found by beating.

## COCCINELLIDÆ.

Since the following notes were written there has come to hand a Revision of the American Coccinellidæ by Captain Casey. Among the large number of new species are many that are accredited to California, but as not a few appear to me to be merely slight color variations of old forms, I have thought it best to exclude them till the matter could be more carefully investigated. Except in the case of *Exochomus*, no reference will be made to this paper.

**Anisosticta.**—*A. seriata*, plentiful in September, along the shore of San Diego Bay (Blaisdell).

**Næmia.**—*N. episcopalis*, one specimen, swept in marshy spot at Tallac, Lake Tahoe (Fényes).

**Megilla.**—*M. maculata*, Colorado River and Palm Springs; I have not seen specimens from the maritime slope as yet: *vittigera*, not uncommon, especially on weeds or low herbage, in river bottoms.

**Eriopis.**—*E. connexa*, "California" (Mulsant).

**Hippodamia.**—*H. ambigua* and *H. convergens* are exceedingly abundant at nearly all seasons throughout Southern California, and intermediates are common: *parenthesis*, very rare, on squash vines, at Poway (Blaisdell): *5-signata*, an example in Mr. Fuch's collection from Los Angeles County, and bearing the label *lecontei*, is scarcely distinguishable from eastern specimens of *5-signata*; it is more than likely that *lecontei* is only a slight variety of Kirby's species: *spuria*, San Diego County (Fuch's collection).

**Coccinella.**—Blaisdell records *9-notata* from San Diego County, but I have not yet seen typical specimens from

our region; the var. *franciscana* is found in most localities, and is not rare. *C. transversoguttata* var. *transversalis* was abundant during the season of 1897, on the islands, and occurred rather frequently at Pasadena, but I do not recall seeing it either before or since: *californica* is rather common every season, and in nearly all localities: *monticola* has been taken by Dr. Fényes at Lake Tahoe (Mt. Tallac, elevation 9,000 feet), and *prolongata* is said by Crotch to occur in California; these two species are exceedingly closely related, and I am inclined to doubt their distinctness. According to Fauvel, the European *11-punctata* Linn. has been taken in California.

**Cycloneda.**—*C. sanguinea* is common and widely diffused: *oculata*, rare, Pasadena: *abdominalis*, moderately common, especially in cultivated grounds.

**Adalia.**—*A. frigida*, "California," must be northern: *bipunctata* var. *humeralis*, one example, Santa Rosa (Ricksecker).

**Harmonia.**—*H. picta*, common on pines in the Sierras.

**Anatis.**—*A. rathvoni*, "California, Oregon": *subvittata*, middle and northern Sierras: an undescribed species from Lake Tahoe, elevation 9,000 feet (Fényes).

**Mysia.**—*M. hornii*, San Bernardino Mountains, July; one example beaten from a pine (identified by Dr. Horn); it is not easy to distinguish this from specimens from the middle Sierras which pass as *Anatis subvittata*, in fact, I believe they are the same thing; it is possible that we do not know Mulsant's species.

**Psyllobora.**—*P. 20-maculata* var. *taedata*, very common everywhere, by beating.

**Chilocorus.**—*C. bivulnerus*, plentiful about orchards and cultivated grounds everywhere: *cacti*, generally much less frequent, probably more common in the southeast.

**Exochomus.**—*E. pilatii*, generally quite rare; on one occasion I found it in some numbers in an olive orchard at Pomona, where it was feeding on the black scale with which the trees were infested; it has also been found rather plentifully at Pasadena, by Mr. Daggett, on certain pepper trees (*Schinus molle*) infested by this same scale. *E. californicus* occurs in the Sierras from the San Bernardino Mountains to Siskiyou, and also in Marin County: *fasciatus* is common throughout our district: *childreni* has occurred at Pasadena and Redondo: *histrion*, Pomona, in the foot-hills.

**Cryptognatha.**—*C. pusilla*, abundant near Pomona, on live-oak, October: *catalinae*, a small number taken by beating on Catalina Island, August; probably not different from the preceding.

**Smilia.**—*S. reversa*, occurs on pines in the Sierras (San Bernardino and Lake Tahoe), and is apparently not uncommon: *ovalis*, specimens so referred are common on live-oak at Pomona, September to November.

**Hyperaspis.**—The species occur by beating and sweeping the shrubbery and herbage of uncultivated ground. *H. fimbriolata* and its variety *dissoluta* are represented in my collection by single specimens taken at Pomona in May and June: *lateralis* is our commonest species: *undulata* (?) is also rather common and very variable: *taeniata*, not rare at Pomona and Pasadena: *excelsa*, rare, Pomona, November; Pasadena, April; San Bernardino Mountains, July: *spiculinata*, not common at Pasadena and Pomona; usually if not always on a species of

tall grass or rush both living (June) and dead (November): *annexa*, not rare; Pomona, June to November: *tristis*, specimens so referred by Dr. Horn were bred in numbers from cactus by Professor A. J. Cook, at Claremont: *postica* was described from northern California, and *osculans* and *4-oculata* from "California" simply.

**Hyperaspidius.**—*H. trimaculatus*, widely dispersed but not very common: *arcuatus*, Colorado River (Le Conte).

**Scymnus.**—A few specimens of *lucidatus* were beaten from pines in the San Bernardino Mountains: *sordidus*, scarce; San Bernardino, Pomona: *guttulatus*, occurs at Pomona, Pasadena, Redondo, Catalina Island, etc., most frequently in March and April, but at no time common: *nebulosus*, tolerably common in the foot-hills near Pomona; taken also in the Ojai Valley and at Catalina Island: *pallens*, common on live-oak nearly everywhere, especially from October to February: *minus*, two examples only; Riverside and Pomona: *cinctus*, moderately common and wide-spread: *pacificus*, rare; Sierras, 3,000 to 5,000 feet elevation: *flexilis*, two examples taken at Yuma: *cervicalis*, rare; Pomona and Catalina Island: *marginicollis*, our commonest species and the only one occurring in orchards as well as on wild growth: *ardelio*, nearly as abundant as the preceding and very widely dispersed, but does not occur on cultivated lands: *punctum*, common, especially on live-oak, in many localities: *nunus*, less frequent than the preceding but equally wide-spread: *coniferarum*, very abundant on pines in the southern Sierras. The following species are more northern in habitat: *debilis*, San Jose and Alameda; *bisignatus*, Siskiyou; *phelpsii*, northern; *caurinus*, Owens Valley (Horn).

*/m* **Scymnillus.**—*S. aterripus* is moderately common in the San Bernardino Mountains, probably on pines; taken by beating.

**Cephaloscymnus.**—*C. occidentalis*, rare; Long Beach, Pomona, Catalina Island: *ornatus*, generally uncommon; Pomona, San Diego, Catalina Island, San Bernardino Mountains. These two species are not congeneric, and it is probable that a new genus will have to be erected for the latter.

**Rhizobius.**—*R. lopanthæ* is now found quite frequently in many localities west of the mountains; it is the only one of the introduced species that has so thoroughly established itself as to be independent of artificial support.

#### ENDOMYCHIDÆ.

Two specimens of an undescribed species allied to *Mycetæa* were found in rotten wood in a cañon near Pomona in June.

**Phymaphora.**—*P. californica*, "San Francisco and western Nevada" (Horn).

**Xenomycetes.**—*X. morrisoni* is found at Castle Crag in fungus (Fényes).

**Aphorista.**—*A. morosa* and *A. letu* occur rather uncommonly at various places in the foot-hills and mountains; both are fungivorus.

**Mycetina.**—*M. hornii* occurs sparingly under bark at Lake Tahoe: *limbata* is probably from the middle regions.

#### EROTYLIDÆ.

**Languria.**—A single specimen of *californica* was taken at Redondo, on *Astragalus crotolaræ*: *convexicollis*, Owens Valley.



**Dacne.**—*D. californica*, common at Pomona, Pasadena, Ojai Valley, etc., in fungi; March: *picea*, described from "California."

**Tritoma.**—*T. californica* is found abundantly in fungi, Ojai Valley, March.

### COLYDIIDÆ.

**Rhagodera.**—*R. tuberculata*, rare, Pomona, May.

**Anchomma.**—*A. costatum* is usually rather rare, occurring singly or in small numbers under stones; once taken in considerable numbers beneath a stone with *Pheidole hyatti* at Redondo in April.

**Megataphrus.**—*M. tenuicornis* is northern (Casey).

**Synchita.**—*S. variegata*, rare; taken occasionally on the wing on warm days in January and February at Pomona; once taken in some numbers under bark at Pasadena in February (Fényes).

**Ditoma.**—*D. ornata*, not rare under bark; Pomona, Yuma, San Bernardino, etc.: *sulcata*, usually not common; once found in numbers under the bark of an old dead stump nearly covered with a woody fungus; Pomona. *D. suffusa* is probably a color variety of *ornata*; it occurs at Yuma under bark with the typical form.

**Phlæonemus.**—*P. catenulatus*, Yuma (Horn).

**Coxelus.**—*C. serratus* and *C. pacificus* occur in the middle coast regions.

**Lasconotus.**—*L. linearis*, not common; on trunks of recently felled alders, San Bernardino Mountains, August: *servus*, a single specimen, flying at twilight; San Bernardino Mountains: *pusillus*, common under pine



bark in the same locality as the preceding: *complex* is northern, while *vegrandis*, *pertenuis*, *nucleatus* and *apicalis* inhabit the middle coast region.

**Aulonium.**—*A. longum*, rare under pine bark; San Bernardino Mountains: *parallelopipedum*, one specimen taken in "California" (Horn).

**Aglenus.**—*A. brunneus*, one or two examples were taken each morning from beneath a portion of the trunk of a recently cut fig tree; others were sifted from stable refuse near by; Pomona, October, February.

**Oxylæmus.**—A few specimens of *californicus* were taken flying at twilight about lumber piles; San Bernardino Mountains.

**Deretaphrus.**—*D. oregonensis*, middle Sierras and north.

**Cerylon.**—*C. castaneum*, not common; found once under oak bark in the San Bernardino Mountains: *californicum* and *sylvaticum* are described by Casey from Lake Tahoe.

**Lapethus.**—*L. discretus*, found under bark; Humboldt County (Casey).

#### RHYSSODIDÆ.

**Rhyssodes.**—*R. humatus*, not common, under bark of fallen conifers, San Bernardino Mountains; much more plentiful at Lake Tahoe and further north.

#### CUCUJIDÆ.

**Silvanus.**—*S. surinamensis*, occurs commonly in stored cereals, etc.: *bidentatus*, "not very common, occurs with the preceding;" San Diego County (Blaisdell):

*imbellis*, one example; San Diego (Blaisdell): *nitidulus*, now referred to *planatus*, was described from California.

**Cathartus.**—*C. advena*, common about gardens, stables, etc.: *opaculus*, not very common but generally diffused; Yuma, San Bernardino, Pomona; once taken in some numbers under bark of eucalyptus in December.

**Nausibius.**—*N. clavicornis*, "Commercial storehouses occasionally" (Blaisdell).

**Prostomis.**—*P. mandibularis* is northern.

**Narthecius.**—*N. grandiceps* is very rare; I once took about a dozen specimens from the burrows of *Pityophthorus pubipennis*, San Bernardino Mountains, in August. Casey has described *simulator* from Santa Cruz County.

**Pediacus.**—*P. depressus*, not common, under bark, San Bernardino Mountains: *fuscus* has been taken near San Francisco (Marin County) by Van Dyke.

**Cucujus.**—*C. clavipes* var. *puniceus*, Lake Tahoe and north.

**Læmophlæus.**—*L. biguttatus*, usually not common; found flying late in the afternoon of warm days in early spring; occurs at times in large numbers under decaying and fungus-covered bark of dead sycamores (Blaisdell): *nitens*, one or two examples taken at Pasadena in March by Dr. Fényes: *horni*, sometimes found in large numbers under bark of dead sycamores, several hundred having been once taken in this way in the Ojai Valley in March: *cephalotes*, rare, under bark of various trees, Pasadena, Los Angeles, San Diego, etc.: *ferrugineus*, not common, Los Angeles: *pusillus*, Colorado River (Le Conte).

**Lathropus.**—*L. vernalis*, not common; about a dozen specimens once taken under bark at Pasadena in February (Fényes). A careful comparison of the type of *pubescens* satisfies me that this is merely a specimen of *vernalis* in which the pubescence of the upper surface is more distinct than usual. This pubescence is plainly visible in specimens of *vernalis* in my own and in the Le Conte Collection, and, I suspect, may always be seen in fresh specimens.

**Brontes.**—*B. dubius* var. *truncatus* is rather common in many localities, especially under oak bark.

#### CRYPTOPHAGIDÆ.

**Henoticus.**—*H. serratus*, not rare, Pomona, Ojai, etc.; taken most frequently on the wing about houses and gardens on warm evenings.

**Cryptophagus.**—A species quite common at Pomona is referred with confidence to *debilis*; the species was described from San Diego: *cellaris* is said to occur at San Diego; it must be closely allied to *debilis* if it is really distinct: *lecontei*, occurs at Yuma. Two other species in my collection, from Pomona and San Bernardino Mountains, are not identified.

**Atomaria.**—At least three species are more or less common under debris and about the roots of plants in damp places; one of these is probably *lætulus* (described from San Jose), which is somewhat variable in color according to the series in the Le Conte cabinet. *A. fus-cicollis* is said by Fauvel to occur in California.

#### MYCETOPHAGIDÆ.

**Mycetophagus.**—*M. californicus* and *M. pluriguttatus* are both uncommon, occurring in fungus; both species

have been taken in Pasadena rather early in spring, and I have found the former in the San Bernardino Mountains in August.

**Litargus.**—*L. balteatus*, common in decomposing vegetable matter; found in many localities.

**Typhæa.**—*T. fumata*, plentiful everywhere.

**Berginus.**—*B. pumilus*, Yuma, Pomona, Redondo, San Diego, Long Beach; sometimes abundant on certain composite flowers, especially near the coast.

**Myrmexhixenis.**—*M. latridioides*, taken not rarely about houses and gardens, Riverside, Pomona, Pasadena.

**Diplocælus.**—An undescribed species taken at Yuma (Wickham).

#### DERMESTIDÆ.

**Byturus.**—*B. grisescens* has been beaten in numbers from live-oak at Pomona and Santa Barbara, during January and February.

**Dermestes.**—*D. marmoratus*, *D. mannerheimii* and *D. vulpinus* are everywhere common: *talpinus* is less common: *tristis*, rather scarce: *carnivorus*, not rare, but occurs, so far as I know, only in the southeastern region: *rattus*, a species quite distinct from *caninus*; is found in the middle coast region and also in the middle Sierras: *lardarius*, has been taken at Lake Tahoe by Dr. Fényes.

**Perimegatoma.**—*P. variegatum*, rare, Pomona and Pasadena: *cylindricum* and *falsum* are also Californian, but I do not know from what locality.

**Attagenus.**—*A. piceus*, occurs everywhere, in houses and on flowers: *varicolor*, southeastern region (Horn):

*perplexus* has been taken at Castle Crag (Fényes): *hornii*, recorded from the "Pacific States" by Jayne, but I have seen no specimens from California; it is rather frequent in Arizona, and it is likely that it may occur in the adjacent desert region of California.

**Trogoderma.**—*T. ornatum*, common and widely dispersed: *sternale*, much rarer; I have taken a single example on Catalina Island.

**Anthrenus.**—*A. scrophulariæ* and *A. verbasci* are both abundant everywhere on flowers; the latter has on one occasion been reared from larvæ found in one of my duplicate boxes.

**Cryptorhopalum.**—*C. apicale*, common on blossoms of willow in April: *balteatum*, much scarcer; on flowers, Pomona: *ruficorne*, two examples; Pomona and Yuma. An undescribed species is not rare at Pomona and Pasadena.

**Orphilus.**—*O. niger* is common nearly everywhere on flowers.

### HISTERIDÆ.

**Hololepta.**—*H. yucateca* is said by Blaisdell to be rather plentiful at San Diego from May to November, in decaying fruit of *Cucurbita*, *Echinocactus viridescens*, and leaves and stalks of *Opuntia occidentalis*. I once found a dozen or more examples in a decaying flower-stalk of *Yucca whipplei*, near Pomona in June. These latter were apparently the form described by Blaisdell as *pervalida*, and are easily separable from Arizona specimens of the typical *yucateca*, and it is not unlikely that Blaisdell's name will be restored to specific standing. *H. cacti* is abundant at San Diego in decaying cacti, and is frequently taken from beneath bark of decaying and water-soaked wood of the willow (Blaisdell); I have never seen

the species near Los Angeles: *vicina*, common in decaying fruit of *Cucurbita*, also between the leaves and in the decaying flower-stalk of *Yucca whipplei*: *neglecta*, rare, in decaying squashes; San Diego (Blaisdell); unknown to me: *populnea*, not common; one example, under bark, Riverside, May; more common in the desert portions of San Bernardino and San Diego counties.

**Hister.**—*H. sellatus*, “not common, in spring and early summer flying about sandy places near streams; also found about the roots of plants” (Blaisdell); I have found it rarely in spring at Pasadena, and in small numbers under plants growing near the beach at Santa Monica, August: *lucanus*, Southern California (Horn): *simplicipes*, one example, San Diego County: *sexstriatus*, not common, occurring in early spring, flying or resting beneath boards or about the roots of plants in grassy places; Riverside, Redondo, Pomona, San Diego: *militaris*, Yuma (Horn); in some seasons quite common, frequenting the sandy banks of streams; San Diego (Blaisdell): *bimaculatus*, one example found dead near a small stream, Riverside: *lecontei*, specimens in the Horn Collection from the southeastern portion of the State: *punctiger*, common under pine bark in the Sierras: *remotus*, recorded from Tejon (Horn): *fractifrons*, Lake Tahoe: *umbilicatus*, Marin County: *immunis*, from “California,” in Horn Collection.

**Tribalister.**—*T. marginellus*, “rare, taken from beneath rocks in moist places” (Blaisdell); I cannot help thinking there is some mistake here, as *marginellus* is a rare eastern species, the only examples known to me being the type described from Maryland, and a single example taken by myself in Rhode Island.

**Tribalus.**—*T. californicus* is common under leaves and stones in moist places near water courses.



**Epierus.**—*E. regularis*, San Diego, San Bernardino: *planulus*, San Diego, Yuma (Horn): *nasutus*, Yuma, under cottonwood bark (Horn). All these are doubtless more common in the southeastern part of the State, and are probably all to be found under the bark of various trees, but especially cottonwood.

**Hetærius.**—No species of this genus is yet recorded from Southern California, but *californicus* has been found at Sonoma, *tristriatus* at Calaveras, and *morsus* at Fort Tejon.

**Onthophilus.**—*O. lecontei*, very rare; three examples taken at Pasadena, in December, under rotten squash.

**Paromalus.**—*P. difficilis*, Mojave (Horn); San Diego (Fuchs): *maneus* is described from Humboldt County by Casey.

**Carcinops.**—*C. opuntiae*, not common at Pomona; more abundant in San Diego County and the southeast, in decaying *Opuntia*: *tejonicus*, a single specimen taken under pine bark in the San Bernardino Mountains: *gilensis*, one example, under chips on a freshly cut oak stump, Pomona; common in Arizona in decomposing cacti (*Cereus*): *consors*, said by Blaisdell to be common in decaying vegetable matter in San Diego County. A species found in numbers in decaying *Yucca whipplei*, near San Bernardino, was identified by Horn as *consors*, but a comparison with the type of this species makes this reference doubtful: *tenellus*, Colorado River, under bark. Casey mentions seeing a specimen of *14-striatus* from Lake County, and I have recently seen the same species from Lower California; it will probably be found in our region.

**Anapleus.**—*A. marginatus*, San Diego County. The species was described by Casey under the name *compactus*, but is considered identical with *marginatus* by authorities.



**Saprinus.**—Numerous species inhabit Southern California, occurring in or about decomposing animal or vegetable matter. *S. discoidalis*, moderately frequent at Redondo beach, especially in spring; occurs also in Owens Valley and in Arizona: *interstitialis*, rare at Pomona, February to March; Catalina Island, and Placer County, early spring: *pectoralis*, not rare from the mountains to the sea, in early spring (February to March); there can scarcely be a doubt that *behrensi* is nothing more than an unusual form of *pectoralis*, in which the prosternal striæ have become united in front. I have seen intermediate forms. *S. obscurus* occurs with *pectoralis* and is about equally common; they are certainly very closely allied, if, indeed, they be distinct: *pæminosus*, moderately rare; Pomona, Ojai, Pasadena, San Diego: *alienus*, occurs only in the desert regions from Owens Valley to San Diego County: *lugens*, excessively abundant at times and always common, widely distributed: *oregonensis*, not rare; many localities: *litticolus*, rather common on Redondo beach, December to April: *scissus*, common on sea beaches; I have never seen it inland: *laridus*, numerous examples taken on the beaches at Redondo and Santa Monica; Wickham reports it from Needles—possibly an error: *insertus*, two examples, Riverside, April: *ciliatus*, Colorado River, rare: *vitiosus*, found once at Yuma; occurs also in Owens Valley and is common in Arizona: *lubricus*, abundant everywhere: *plenus*, desert regions of the southeast: *fimbriatus*, very common and wide-spread: *cærulescens*, found once at Riverside in April, about decaying fish; Blaisdell says, “quite common in summer about the dead bodies of snakes and small animals,” San Diego: *intritus*, San Diego (Casey): *consobrinus*, one example, Yuma, July; *bigemmeus*, common on all beaches; specimens received from Wickham are labeled Los Angeles:

*estriatus*, several examples taken near San Diego in the bed of the San Diego River, August; also on margin of salt pool at Santa Barbara, August: *lucidulus*, plentiful in the sand dunes at various beaches: *propensus*, San Diego (Casey): *gaudens*, San Diego, Ventura; uncommon on sea beaches: *serrulatus*, rare; sea beaches, San Diego, San Pedro: *sulcifrons*, common on beaches. The following species are also Californian: *rotundatus*, near San Francisco and in Placer County (Van Dyke); *aequipunctatus*, *ciliatus* and *vestitus*, from near San Francisco; *lentus*, Truckee; *opacellus*, northern.

**Plegaderus.**—*P. fraternus* and *P. nitidus* are usually common under pine bark in the Sierras: *consors* I have seen labeled "San Diego County:" *molestus*, described from Lake Tahoe.

**Teretrius.**—*T. placitus*, one example found flying early in March at Pasadena (Fényes): *obliquulus* has been taken at Santa Monica by Mr. Albright.

**Abræus.**—*A. bolteri*, San Bernardino (Le Conte).

**Bacanius.**—*B. globulinus*, one example taken at Pasadena is thus referred; it is not typical, but as northern specimens show considerable variation, notably in the development of the elytral striae, it is likely that the differences shown are not specific.

**Acritus.**—*A. maritimus*, rare, on sea beaches; San Diego and Santa Barbara: *volitans*, several specimens taken flying in the early evening; Pomona, August; others at electric light, Pasadena (Fényes).

**Æletes.**—*Æ. basalis* occurs near Yuma.

## NITIDULIDÆ.

**Brachypterus.**—*B. troglodytes* is occasionally numerous on nettles; Pasadena, Pomona; April, May.

**Cercus.**—*C. sericans*, plentiful on flowers of elder (*Sambucus glaucus*) throughout Southern California and the adjacent islands; very variable in size and color.

**Amartus.**—*A. tinctus* and *A. rufipes* are found on flowers, in March; the former is the more common and has been taken most often from poppies (*Eschscholtzia*).

**Athonæus.**—*A. agavensis*, San Diego and Los Angeles; on flowers of *Agave*.

**Carpophilus.**—*C. yuccæ*, Mojave desert, in unopened flower heads of yucca (Crotch): *hemipterus*, abundant: *dimidiatus*, *decipiens* and *discoideus*, less common, in decaying fruit in orchards; September to November: *pallipennis*, plentiful in cactus blooms in all localities: *brachypterus*, northern part of the State: *niger*, recorded from "California" (Santa Cruz, Fuchs).

**Colastus.**—*C. truncatus*, frequent under bark of freshly cut trees; most specimens are of the form *limbalis*.

**Conotelus.**—*C. mexicanus*, southeastern California.

**Epuræa.**—One example (a female) of a species near *rufa* was taken in the San Bernardino Mountains: *terminalis* was plentiful in the same locality under rubbish along streams: *avara*, one example, same locality: *ovata*, not common at Pomona, in decomposing mushrooms; February. According to Crotch, *monogama* occurs "throughout the Sierra Nevada, in a small white globular fungus growing on dead pines, each fungus having a pair of the species." *E. ambigua* and *E. truncatella* are northern: *festiva*, I have taken on flowers at Lake Tahoe in July.

**Nitidula.**—*N. ziczac*, not very common at Pomona and in the southern Sierras, about dried animal remains, old bones, etc.; more common farther east.

**Omosita.**—*O. discoidea*, not common, in decomposing organic matter, Pomona, San Bernardino Mountains.

**Perthalykra.**—*P. murrayi*, San Francisco and north, also at Lake Tahoe (Fényes). Several specimens taken in the San Bernardino Mountains are not quite typical but doubtless belong here.

**Pocadius.**—*P. dorsalis*, "California" (Ulke).

**Meligethes.**—*M. brassicae*, common on willows in early spring; Pomona, Pasadena, Ventura, etc.

**Cybocephalus.**—*C. californicus* is not rare on live-oak, taken by beating; Pomona, Pasadena.

**Cryptarcha.**—*C. concinna*, one example, by sweeping near brook at Pasadena, August.

**Ips.**—*I. vitattus* and *I. cylindricus* are northern.

**Pityophagus.**—*P. rufipennis*, not common in the Sierras from San Bernardino north.

**Smicrips.**—*S. hypocoproides*, Palm Springs (Hubbard).

**Rhizophagus.**—*R. sculpturatus*, Placer County (Van Dyke).

#### LATHRIDIIDÆ.

**Holoparamesus.**—*H. kunzei*, one example taken on the wing in the early evening, at Riverside, July; others have been taken in "Los Angeles County" by Coquillett: *pacificus*, one example, under bark of decaying log at Pomona, April; also taken by Coquillett in "Los Angeles County:" *caularum*, taken in flight, also under bark, by Dr. Fényes, at Pasadena in February and again

in September; found about grass roots at Palm Springs, and in debris along the Colorado River by Mr. Hubbard.

**Dasycerus.**—Specimens of *angulicollis* have been taken by Dr. Fényes at Monterey, under the fungus-grown bark of a decaying stump.

**Metophthalmus.**—*M. rudis*, Ojai Valley, Los Angeles, and Panamint Valley; specimens were taken by me in March at the former locality, occurring in the interior debris of a woody fungus: *trux*, one specimen taken on the under side of a log in a moist situation in the San Bernardino Mountains, July: *parviceps*, found in the region about and to the south (San Jose) of San Francisco Bay.

**Lathridius.**—*L. armatulus*, a rather common species in Southern California from the foot-hills of the Sierras to the coast and as far north as San Francisco, commonly obtained by beating and sweeping; it is not confined to any one plant, but I have taken it most commonly near Pomona on *Solanum nigrum*, in October and November. *L. costicollis* is plentiful near San Francisco.

**Coninomus.**—*C. constrictus* and *C. australicus* are not rare in our region and probably occur nearly everywhere in the State; the former is a well known cosmopolitan species; the latter was described from Australia. *C. nodifer* has been taken in some numbers near San Francisco.

**Enicmus.**—*E. suspectus*, Pomona; Panamint Mountains; not common: *desertus*, Palm Springs and Yuma (Hubbard): *minutus*, occurs everywhere, but is not common in Southern California: *crenatus*, plentiful about the roots of herbage and in vegetable debris throughout Southern California: *tenuicornis*, Pomona, rare;

commoner further north: *nigritus* and *ventralis* have both been taken at Los Gatos and in the middle Sierras: *sulcatulus* and *fictus* are Californian without more definite locality.

**Cartodere.**—*C. argus*, several examples on drying yucca pith at Pomona: *quadrioveolata*, a single specimen, Los Gatos.

**Revelieria.**—*R. californica* was taken in company with *Metopthalmus rudis*, in woody fungus, Ojai Valley, March. I have seen specimens from "Los Angeles County" and Los Gatos.

**Corticaria.**—The various species occur under bark, and are frequently taken on the wing at sunset or in the early evening, and occasionally come in numbers to electric lights. None of the species appear to be very abundant, but several would doubtless prove common enough if properly looked for. I have seen specimens of *planula* from Pomona, Pasadena, Ojai Valley, and also from several locations in the central and northern part of the State; the majority of specimens have been taken flying, in early spring evenings. *C. serrata*, *C. elongata* and *C. ferruginea* are cosmopolitan species; a moderate number of the last two have been seen from Los Angeles County: *serrata* is apparently rare in this region: *occidua* and *tenuipes* have thus far occurred only in the Sierras, the latter being known only by the type, which was taken in the San Bernardino Mountains: *inopia* is known from the Tahoe region; *prionodera* is from the middle coast region.

**Melanophthalma.**—The species of this genus occur on vegetation, and may usually be taken plentifully by beating and sweeping. *M. casta* has been taken at Pomona, Pasadena and Redondo, by sweeping weeds:



*simplex* occurs on both sides of the mountains, but not very abundantly: *insularis* is known only from San Clemente Island: *distinguenda*, *gibbosa* and *americana* are all very common all over the State (I have not seen any *gibbosa* that were taken in the desert but it may occur there): *incompta* is represented by only two examples taken at electric light in Pasadena by Dr. Fényes: *similata* is plentiful near the coast from San Diego to San Francisco: *tenella* is based upon two examples in the Le Conte Collection that were found at San Jose: *villosa* has occurred in the Lake Tahoe region; other examples in the Horn and Le Conte collections are probably from the desert, though this is not certain.

**Fuchsina.**—*F. occulta*, “Los Angeles County;” one example seen, collected probably by Coquillett. The specimen in my collection was taken by Mr. Fuchs in Mill Valley, near San Francisco, by sifting earth about the roots of a redwood. A specimen in the Hubbard and Schwarz Collection is labeled Los Gatos.

### TROGOSITIDÆ.

**Alindria.**—*A. teres* is rare; found under Pine bark, San Bernardino Mountains.

**Trogosita.**—*T. virescens*, not rare under bark throughout Southern California in and near towns; it is not unusual under the loose layers of bark of eucalyptus: *yuccæ*, Mojave desert, under bark of tree yucca.

**Tenebrioides.**—*T. sinuata*, moderately common under bark in the Sierras. The following species or varieties are recorded from this State: *crassicornis*, *pleuralis*, *intermedia* and *californica*. With the exception of *intermedia*, from San Jose, I am unable to ascertain exact



localities. It is practically impossible to identify specimens with any degree of certainty with existing literature.

**Peltis.**—*P. pippingskoeldi*, Sierras; under bark.

**Grynocharis.**—*G. pilosula*, Sierras, Mt. Wilson, elevation 6,000 feet.

### MONOTOMIDÆ.

**Monotoma.**—*M. picipes*, not rare on the wing, and is taken by sifting in spring: *mucida*, not common, under leaves, etc., along the Colorado River at Yuma.

**Phyconomus.**—*P. marinus* is rather common under rubbish along the Southern California beaches above tide-water.

**Hesperobænus.**—*H. abbreviatus*, common under bark of freshly cut deciduous trees, flower-stalks of yucca, etc.

**Europs.**—*E. longicollis*, near San Francisco; under bark.

**Bactridium.**—*B. striatum*, one specimen from Pomona, November; more frequent under bark in the desert region of the southeast. Numerous specimens of a species closely allied to, if not identical with *striolatum* were found under the bark of sycamore, in the Ojai Valley, in March.

### BYRRHIDÆ.

**Nosodendron.**—*N. californicum*, one example from Pine Flats, elevation 6,000 feet, Los Angeles County; commoner in the Sierras farther north. .

**Amphicyrta.**—*A. dentipes*, one example under freshly cut live-oak chips; Pasadena, January 1. Specimens scarcely distinguishable are in my collection from northern California, under the name *chrysomelina*.

**Cytilus.**—*C. sericeus*, Lake Tahoe and north.

**Limnichus.**—*L. californicus*, common by washing the margins of streams; Pomona, Riverside, Palm Springs, San Bernardino Mountains, etc.: *perpolitus*, less common, or at least more local; I once took it in some numbers along the Santa Ana River at Riverside and again near Pomona, April to June: *nebulosus*, not common; Pomona, Riverside, Pasadena: *tenuicornis*, once found in numbers on a wet, springy hillside in the San Bernardino Mountains, elevation 6,000 feet: *naviculatus*, not rare at Los Angeles and Riverside: *analisis* will probably be found in Southern California, as it is recorded from "California and Arizona;" my specimens are from the middle coast region: *perforatus* is described from Santa Clara County.

**Bothriophorus.**—*B. minutus*, Riverside and Yuma, along the margin of streams.

#### GEORYSSIDÆ.

**Georyssus.**—*G. californicus* was taken once along the Truckee River, near Lake Tahoe; probably common enough in the middle and northern parts of the State, and may very likely yet be found in the mountains in the south.

#### PARNIDÆ.

**Psephenus.**—*P. falli*, rare; a small colony once found running over the rocks lying unsubmerged in the middle of a shallow stream near Pomona, June 4; a single specimen taken along a mountain stream at Arrowhead Springs, May 30: *veluticollis* occurs in Mendocino County.

**Lara.**—*L. avara* is northern.

**Throscinus.**—*T. crotchii*, several specimens found on mud-flats which are covered at high tide; Coronado, February, July.

**Dryops.**—*D. productus* and *D. suturalis* are both abundant in all streams: *striatus* occurs in middle California.

**Narpus.**—*N. angustus*, northern California (Mendocino County).

**Elmis.**—*E. divergens*, two examples taken from a small reservoir at Pomona, May 11; originally described from Ventura (Le Conte): *foveatus*, several examples at Pomona, June: *quadrinaculatus* is northern: *seriatus* occurs at the Geysers.

**Stenelmis.**—A small series of *nubifer* was found in a little mountain stream near Pasadena, October 31.

**Macronychus.**—*M. parvulus*, Pomona and Pasadena, in mountain streams.

### HETEROCERIDÆ.

**Heterocerus.**—The species, except *pusillus*, are about equally common west of the mountains, on muddy margins of streams and ponds. *H. pusillus* probably occurs only in the southeast. I once took *gnatho* in numbers, at electric lights in Riverside. *H. brunneus* occurs in the north.

### DASYLLIDÆ.

**Macropogon.**—*M. testaceipennis*, Mariposa district, rare (Horn); Alameda (Fuchs).

**Eurypogon.**—*E. confusus*, one example; Pasadena, May (Fényes): *californicus*, described from a single specimen taken at Geysers.

**Allopogon.**—*A. villosus*, not common at Pasadena and Pomona; taken in May, by sweeping weeds by the roadside.

**Stenocolus.**—*S. scutellaris*, rare; Sacramento and Yosemite Valley.

**Dascyllus.**—*D. davidsonii* is not uncommon in the country to the north of San Francisco (Lake and Sonoma counties, etc.): *plumbeus* occurs in the middle Sierras.

**Anorus.**—*A. piceus* is taken rather frequently in May and June, flying at dusk; Pomona, Pasadena, Arrowhead Springs; San Diego to Owens Valley (Horn).

**Anchycteis.**—*A. velutina* occurs in the Shasta region.

**Eucinetus.**—*E. infumatus*, one example, Santa Barbara, February; taken by sweeping; "occurs near San Francisco, under bark, in February" (Horn).

**Acneus.**—*A. quadrimaculatus* occurs north of San Francisco.

**Helodes.**—*H. apicalis*, not common; found in the Sierras from Los Angeles County north. A single example of an undescribed species has occurred at Pomona.

**Cyphon.**—*C. exiguus*, rare at Pasadena; taken by sweeping near water: *concinus*, rather plentiful on coniferous trees in the Sierras: *variabilis*, specimens so referred are quite common on willows at Pasadena in May: *brevicollis* occurs in the northern portions of the State.

**Placonycha.**—*P. edwardsii* occurs along mountain streams during May and June. The males are not rare, flying about and resting on rocks and overhanging foliage (the females are less common).

In its earlier stages, *Placonycha* is found on rocks under water; the insect rests in a limpet-like covering till the pupa is developed, after which time the beetle leaves the cover and lives out of water (Fuchs).

### RHIPICERIDÆ.

**Sandalus.**—*S. californicus*, "California and Nevada;" occurs at San Diego and Los Angeles (fide Fuchs).

### ELATERIDÆ.

**Melasis.**—*M. rufipennis*, Sonoma County; probably confined to the redwood belt (Ricksecker *in litt.*).

**Eucnemis.**—*E. americana* occurs near Mt. Shasta.

**Dromæolus.**—*D. californicus*, "California:" *nitens*, northern: *basalis*, Lake Tahoe and in the mountains of the more northern portions of the State.

**Xylobius.**—*X. cylindriciformis*, "California and Nevada" (Horn); Mendocino County (Van Dyke).

**Sarpedon.**—*S. scabrosus*, one example, Pasadena (Fényes).

**Palæoxenus.**—*P. dohrnii*, a species said to have been found near San Diego, but this is probably an error, as is suspected by Horn; the species is quite surely tropical.

**Anelastes.**—*A. druryi* is common in the higher parts of the Sierras; frequently taken on the wing about camp-fires, in the San Bernardino Mountains, July and August.

**Adelocera.**—*A. sparsa*, rather rare in Los Angeles County: *profusa* and *rorulenta* occur in the Lake Tahoe region, under bark.

**Meristhus.**—*M. cristatus* occurs at Yuma, under debris along the river bank; one example was also taken at Riverside; found flying in the evening, in April, at Pasadena (Fényes).

**Chalcolepedius.**—Le Conte records *C. rubripennis* from Cajon Pass; there is possibly some mistake in the locality: *webbii* is common on willows along the Colorado River: *tartarus*, rare in the same locality.

**Alaus.**—*A. melanops*, southern and middle Sierras.

**Cardiophorus.**—*C. amplipennis*, not rare on willows, Pomona, Ojai, etc.; March and April: *gemmifer*, one example in the Horn Collection bears a label indicating its occurrence in the southeast: *luridipes*, common throughout Southern California, December to June: *edwardsii*, rare, Los Angeles County (Van Dyke): *latiusculus*, said to occur in "Southern California:" *tenebrosus* var. (?) *fulvipes*, rather common; Pomona, Santa Monica, Ventura, etc.; January to May. It is probable that *fulvipes* is distinct from *tenebrosus* by the differently formed genitalia, as pointed out by Mr. Blanchard in his revision. I have not seen specimens of typical *tenebrosus* from Southern California, but they are common in the mountains farther north. *C. aeneus* is apparently not very common; I have seen specimens from San Diego: *seniculus*, rare; Pomona and Pasadena, February and March. In addition to the above, *bifasciatus* and *obscurus* occur about San Francisco; *fenestratus*, *coxalis* and *stigmaticus* are northern; *tumidicollis*, middle Sierras; *dispar*, eastern part of the State (Owens Valley ?); *abbreviatus*, *crinitus* and *carbonatus* are said to be from "California."

**Horistonatus.**—*H. inanus* and *H. sufflatus* are not rare at Pomona, Pasadena, etc., during May and June; taken



by sweeping: *basalis*, not common; Riverside, February to May: *simplex* occurs at Yuma: *flavidus*, several examples taken at light, at Palm Springs (Fényes): *transfugus*, central and northern California: *gracilis*, about Lake Tahoe.

**Esthesopus.**—*E. dispersus*, "Southern California" (Horn). Unquestionably a desert species.

**Cryptohypnus.**—*C. squalidus*, rare; found by washing the gravelly margins of mountain streams near Pomona: *nocturnus*, found in the north: *funeris*, recorded by Horn from "California and western Nevada;" it is probably confined to the middle Sierras in this State.

**Hypnoideus.**—All the species occur along streams in the mountains and foot-hills. *H. striatulus* has occurred but once near Pomona, in May: *gradarius* is not common; Pomona, Riverside, San Bernardino; April and May: *ornata*, common at times and quite widely dispersed: *pectoralis*, plentiful almost everywhere.

**Anchastus.**—*A. cinereipennis*, a rather common species, occurring under sticks and stones, about the roots of grass, at foot of trees, etc., near or within towns and cities, from San Diego to the northern portions of the State: *bicolor* var. *desertus* has been taken at Yuma: *tantillus* and *regularis* are said to be from California, but I have never seen specimens; they are doubtless northern, and one or both may be nothing more than a variety of *cinereipennis*, which is somewhat variable.

**Elater.**—*E. hepaticus*, a single example taken by sweeping, near Pomona, June: *fastus*, "Southern California" (Le Conte); I am unable to ascertain the exact locality of a pair collected by Morrison, but very likely they are from the southeast: *cordifer*, not common, under bark of



decaying stumps; Pomona, Ojai Valley, March and April: *ater*, a single specimen obtained in the San Bernardino Mountains, under bark, in July: *longicornis*, Santa Ana River (Le Conte); I have seen only the type: *carbonicolor*, a single specimen so referred was taken by sweeping, in the foot-hills near Pomona, in June: *dimidiatus*, this species under the name *affinis*, which was given by Le Conte to a form which differs in no respect except in the black portion of the elytra involving the apical third instead of half, is reported from "Washington Territory and Southern California;" I have found it in the Lake Tahoe region, and its occurrence in the mountains further south is not surprising. *E. rhodopus* is not rare under bark of decaying logs, in northern part of the State and Oregon: *morrens* is also northern: *melinus*, *phelpsii*, *cordatus*, *behrensii*, and *apicatus* are found in the middle Sierras and extend their range more or less to the north: *horni* and *atripennis* are given as Californian, but without indication of locality. There is little doubt that in collections more than one species is now included under the name *apicatus*, and it would not be surprising if the Californian specimens now so called were distinct.

**Elatrinus.**—*E. anthrax*, "California." A specimen in Mr. Ulke's collections is without definite locality.

**Drasterius.**—*D. liveus*, not very common; Yuma, San Bernardino (Wright), Santa Monica: *grandicollis* and *elegans* are both reported from "California," and it is not unlikely that both occur in the south.

**Megapenthes.**—*M. tartareus*, rare; a single example taken near Pomona in June, by beating; several specimens taken at Pasadena, in April, on thistle bloom (Fényes): *turbulentus*, Pomona, San Diego, Pasadena;

once taken in some numbers at the latter place on flowers of "greasewood," in June: *aterrimus*, beaten from pines in the San Bernardino Mountains, but not common: *elegans*, "occurs in the southern Coast Range" (Horn); Monterey (Rivers): *stigmatosus*, common on pines in the San Bernardino Mountains: *nigri-ventris*, northern part of the State: *4-maculatus* and *lepidus*, central part of the State.

**Ludius.**—*L. lecontei*, not common, under bark of logs or stumps; Riverside, Pomona, San Diego, Santa Barbara, etc.: *ater* occurs at Santa Barbara.

**Agriotes.**—*A. imperfectus*, "Southern Coast Range" (Le Conte): *hispidus*, not common; Pasadena, in May, by sweeping; San Bernardino Mountains, July; San Bernardino, under bark of eucalyptus within the city, December: *fucosus*, *apicalis*, *thevenetii* and *torquatus* occur in the mountains from Yosemite Valley north.

**Dolopius.**—*D. lateralis* is everywhere abundant and excessively variable.

**Melanotus.**—*M. longulus*, San Diego (Le Conte): *fissilis*, Yuma (Wickham): *cribricollis*, "Southern California and Arizona:" *variolatus*, San Pedro and Catalina Island; uncommon. An undescribed species is rather plentiful at times, especially on willows in spring; Pomona, Riverside, Pasadena. *M. oregonensis* appears to be a rather common species from San Francisco north.

**Limoni.**—*L. mirus*, San Diego (Le Conte): *crotchii*, one specimen found crawling on the sidewalk at Pasadena, April 4: *occidentalis*, several specimens obtained by beating, in the foot-hills near Pomona, May. A species once determined for me by Dr. Horn as *infus-catus*, but which on comparison with a typical specimen

of the latter in the Le Conte Collection seems to be distinct, is found quite frequently on willows in spring at Pomona, Riverside, Pasadena, etc. *L. ornatulus*, San Bernardino Mountains, June (Daggett); much commoner farther north: *pilosus*, described from San Diego; my specimens are from Sonoma County: *californicus*, widely diffused and generally common; specimens were dug out of a dead log at Pomona in February: *canus*, described from San Diego; my specimens are from San Francisco; it seems quite uncommon, at least in the south. Of the other species of the genus occurring within the State,—*fulvipes* is rather rare about San Francisco; *maculicollis* is found in the middle Sierras and north; *consimilis* in Santa Clara County, *nitidulus* in the north, and a species near *discoideus*, and now passing as that, in Kern and Sonoma counties.

**Pityobius murrayi.**—This species occurs, but not over abundantly, from Marin County north.

**Athous.**—*A. excavatus*, not rare, by beating and sweeping in the foot-hills, and at higher altitudes of the southern Sierras: *limbatus*, rare in the San Bernardino Mountains; I have also taken it at Lake Tahoe. Two undescribed species have been taken in our district; one of them common in the San Bernardino Mountains; the other represented by two specimens taken near Pomona. *A. opalinus* is found in the middle Sierras: *excavatus*, in the Coast Range, south of San Francisco: *vittiger* and *nigripilis* in the north. There are besides in my collection six unidentified and probably undescribed species from the middle and northern regions.

**Leptoschema.**—Specimens of *protractum* have been seen from Santa Clara County.

**Sericosomus.**—*S. debilis* is rather common; taken by beating and sweeping, in the foot-hills near Pomona, April. I am unable to separate *flavipennis* Mots from *debilis*. The former name is in general use for the smaller, paler specimens, and the latter for larger specimens with the thorax more or less dark; there are, however, all degrees of size and color between the extremes.

**Corymbites.**—This genus is numerously represented in the State, but very few species occur in our district. *C. maurus* is rare in the foot-hills near Pomona in June: *jaculus* is described from San Diego, but also occurs in the north (Sylvania): of *fallax* I have seen specimens labeled "So. Cal.," but have never taken the species here myself; it is certainly more common in the north. I have seen specimens of *pruininus* and *leucaspis* labeled southeastern California in the Horn Collection. Of the other species occurring within the State, *cribrosus*, *semivittatus*, *rotundicollis* and *conjungens* have been taken about San Francisco, the latter also in the middle Sierras; *anthrax* and *colossus* are from the central region; *monticola*, *fraternus*, *aripennis*, *carbo* and *obversus* from the middle or northern Sierras; *nigricollis*, one example from Lake Tahoe; *edwardsii* from Santa Cruz and Placer counties and north; *breweri* from Visalia and north; *fuscus*, *caricinus*, *xanthomus*, and *opaculus* from the northern region only; *obscurus*, *teres* and *rupestris* from "Cal." without definite locality. Besides these there are several undescribed species in collections.

**Oxygonus.**—*O. ater* is described from Amador Valley, central California.

**Asaphes.**—*A. morio* and *A. dilaticollis* are rare in the San Bernardino Mountains in July and August:

*tumescens* is recorded from Santa Cruz Island: *hirtus* and *carbonatus* occur near San Francisco.

**Melanactes**.—*M. densus* is not common at San Bernardino, Riverside, Pomona, etc.; found under bark and in decaying trees and stumps, in April and May.

**Aphricus**.—*A. californicus* is occasionally found at Pasadena and Pomona, by sweeping weeds and herbage by roadsides, May to August. According to Horn it has been found from San Diego to Owens Valley.

**Aplastus**.—*A. angusticollis* was many years ago taken in numbers at San Diego by Mr. Crotch; it has since been taken sparingly at the same place, in April, by Mr. Dunn: *corymbitoides* is a rare species which has been found at Riverside and Los Angeles: *speratus* is rather common at Pomona, where on warm evenings in May and June the males may be taken on the wing or at light; the females are wingless and are very seldom discovered: *tenuiformis* is described from "Cal.," and is probably from the central region: *optatus* has been found at Fresno, and in Alameda and Santa Cruz counties: *molestus* is recorded from near San Francisco.

**Plastocerus**.—*P. schaumii* is rather widely diffused and rather common at Pomona, Pasadena, etc., in June, coming often to light; the females are not apterous, but are very rarely taken.

**Euthysanius**.—*E. lautus* is found at same places and under same conditions as *P. schaumii*, but is less frequent at Pomona in my experience. Judging from a small collection sent me by a friend, it is not rare in the vicinity of Long Beach. *E. pretiosus* is certainly rare in our district, though specimens have been seen labeled "So. Cal.;" it is more common in the central valley.

**Perothops.**—*P. witticki* is rare in the central region, and has been taken sparingly at Kaweah, in Tulare County, by Mr. Hopping.

**Cerophytum.**—*C. convexicollis* is described from Fort Tejon; it is evidently a very scarce insect.

### THROSCIDÆ.

**Aulanothroscus.**—*A. validus* occurs in the middle Sierras.

**Throscus.**—*T. sejunctus* was described from Mojave; a specimen in my collection from Long Beach is so referred: *parvulus* is recorded from "Mojave and Yuma" (Horn); *sericeus* is common and quite widely dispersed: *mendax* is found in the Sacramento Valley.

**Pactopus.**—*P. hornii* has been found in some numbers flying at twilight about board piles in the San Bernardino Mountains, July.

### BUPRESTIDÆ.

**Gyascutus.**—*G. planicosta* was found in abundance at Needles by Mr. Wickham: *obliteratus* has occurred at San Bernardino in June (Fényes), at Banning, on flowers of mesquite, July 3 (Van Dyke), and is reported as being plentiful on low willows in Owens Valley (Horn): *cuneatus* was described from a single specimen taken with the preceding species in Owens Valley.

**Hippomelas.**—*H. californicus*, Banning, July 3 (Van Dyke); described from San Joaquin Valley.

**Chalcophora.**—*C. angulicollis* is not rare at Lake Tahoe and north.

**Dicerca.**—*D. californica* has been found on Echo Mountain and on Mt. Wilson (Los Angeles County), and



according to Van Dyke depredates on Douglas Spruce: *hornii* is not rare in the foot-hill cañons of the southern Sierras and at Santa Monica; I have taken the beetles from their burrows in alder stumps: *pectorosa* has been taken at Acton, Los Angeles County, by Coquillett (fide Van Dyke).

**Pæcilonota.**—A specimen referred with some doubt to *ferrea* was taken by sweeping the herbage along a small brook near Pasadena in August.

**Buprestis.**—*B. gibbsii* has been found in San Diego County (fide Van Dyke); specimens in Mr. Rivers' collection were taken in Yolo County: *leviventris*, *maculiventris* var., and *aurulenta* occur in the San Bernardino Mountains in July; the latter is the most common and ranges north to Vancouver: *fasciata* is represented by a single specimen taken by sweeping beside a brook in the San Bernardino Mountains, at an altitude of 5000 feet: *connexa* occurs in Owens Valley: *adjecta* is found in the middle Sierras (Lake Tahoe) and north.

**Melanophila.**—Of *M. consputa* several examples have been taken by Dr. Fényes and myself about the streets and on windows at Pasadena, September to November: *acuminata*, which name is now used for *longipes* Say, is reported from Santa Monica by Van Dyke: *gentilis* and *intrusa* may be found in the coniferous belt of the San Bernardino Mountains; the former I have taken from their burrows in pine twigs, and the latter I have beaten from *Pinus ponderosa*.

**Anthaxia.**—*A. æneogaster* is widely distributed, but is more common in the mountains; I have taken it on various flowers, from May to August; it is said to breed in white oak (Van Dyke), and in the California walnut, *Juglans californica* (Coquillett): *deleta* is less common,



and so far as I know is found only at altitudes of 4000 feet and upward in the southern Sierras; it is found at lower altitudes further north. There are in Mr. Ulke's collection two examples of *quercata* said to have been taken in California.

**Chrysobothris.**—*C. octocola* has been taken at Indio and along the Colorado River, and is likely to be found wherever there is mesquite, in which it breeds: *debilis* also breeds in mesquite (Van Dyke); I have taken it at Palm Springs in April, and Dr. Van Dyke has found it at Banning from May to July: *femorata* has been taken in the San Bernardino Mountains by myself, and Dr. Blaisdell reports finding both pupæ and beetles in their burrows in the bark of the trunk of the live-oak (*Quercus agrifolia*), at San Diego. Dr. Blaisdell also records the breeding of "*semisculpta*" (now placed with *contigua*) from the half dead limbs of apple and live-oak; San Diego. *C. cuprascens* is very abundant on pines in the San Bernardino Mountains, in July and August: *speculifer* has been taken at Pasadena, by Dr. Fényes, in May: *californica* is not common, on pines in the San Bernardino Mountains, August; specimens have been taken by me from their burrows in the smaller branches of *Pinus ponderosa* (?); Dr. Blaisdell reports this species as being very injurious to apple trees at or near San Diego. *C. mali* is widely distributed and seemingly variable in habits. Horn says in his monograph of the genus, "Specimens from the Sacramento Valley were sent me by Mr. L. E. Ricksecker as infesting apple trees; others collected in Owens Valley could not possibly have had that habit." Dr. Van Dyke writes me that he has found it at Banning on mesquite, and at Santa Monica Cañon on *Ceanothus*; while I have myself taken it at Pomona and

Riverside, beating it on one occasion from live-oak. *C. deserta* occurs on the Mojave Desert and near San Diego (Horn): *merkelii*, Banning, May to July, on mesquite (Van Dyke): *prasina*, one example (Fényes), Pasadena: *lucana*, Pasadena, one specimen taken on sidewalk (Daggett); also at Santa Monica, on flowers (Van Dyke). Other species that are known from the State are as follows: *viridicyanea*, *dolata*, *ludificata*, *trinervia*, *nixa* and *ceneola* are from the middle Sierras; *deleta* from Owens Valley; *cyarella* from Shasta region; *texana* from "Cal.," probably the southeast.

**Actenodes.**—*A. mendax* occurs at Yuma, and *calcarata* must also be from the desert region of the southeast. The species which has stood on our lists as *Belionota californica* Mots. is now referred to *Actenodes acornis* Say; I am not able to name the locality from which specimens have been obtained.

**Glyptoscelimorpha.**—*G. marmorata* has thus far been taken only at Lancaster, on the border of the Mojave Desert. We owe its discovery to Mr. Coquillett.

**Dystaxia.**—*D. murrayi* is an uncommon insect which has, however, been found at numerous localities. Mr. Coquillett has taken it at Lancaster, Dr. Van Dyke at Banning, on live-oak, and at Santa Monica Cañon, and Mr. W. G. Wright has a considerable series taken in San Bernardino County.

**Schizopus.**—*S. latus* is a rare species which has been found at San Diego. *S. sallei* is described from Mariposa.

**Polycesta.**—*P. velasco* is from the Colorado Desert (Indio and Yuma); Dr. Van Dyke reports it as occurring on mesquite at the former place: *P. californica* has

been taken at Pasadena in May and June, on oak wood-piles; it is reported by Mr. Coquillett as breeding in oak.

**Acmaeodera.**—This genus is very numerous represented in California, and of the thirty-two species recorded from the State all but three—*mariposa* from the middle Sierras, and *robusta* and *pubiventris* from Owens Valley—are found in our district. It is more than likely that these three species will eventually be found here. The various species are taken on flowers in May and June, but their breeding habits are almost unknown. According to Van Dyke, *dohrnii* breeds in *Ceanothus* and *gibbula* in mesquite. The following species have thus far been taken only in the Desert region of the east and southeast: *flavomarginata*, *lanata* (Los Angeles and San Bernardino counties), *tuta* (Death Valley), *variegata*, *quadrivittata* (Mojave). Of the remaining species, *fenyesi* is not rare at various localities in Los Angeles, Riverside, and San Bernardino counties; *plagiaticauda*, rare, Mariposa and Los Angeles counties; *jocosa*, foot-hills of the southern Sierras, Pomona and Pasadena; *coquilletti*, "Los Angeles County" (Coquillett); *angelica*, foot-hills of Sierras in Los Angeles County and north to Lake County; *hepburnii*, rather common west of the Sierras nearly throughout the State; *morbosa*, Pomona and Pasadena, foot-hills, rare; *flavosticta*, Riverside and San Diego County; *acuta*, *labyrinthica*, and *connexa* are all common and widely dispersed in and to the west of the Sierras, the last named, however, is seldom found in our district except at considerable altitudes in the Sierras; *vandykei*, scarce in Los Angeles County, more frequent further north; *prorsa*, Los Angeles and San Bernardino counties; *dohrnii*, rare in the southern Sierras (Los Angeles

County); *dolorosa* and *postica* are known only from Los Angeles County, the latter being represented by a single specimen taken by Coquillett; *alicia*, one example, Los Angeles County (Coquillett); *versuta*, rare in Los Angeles and Mariposa counties; I have taken the species from yucca blooms in June; *guttifera*, one example taken in Los Angeles County by Van Dyke; *gemina*, Los Angeles and San Bernardino counties to Siskiyou County, but not common; *comata* and *alacris* are described as from "near Fort Yuma."

**Ptosima.**—One example of *P. walshii* has been taken by Mr. Coquillett in Los Angeles County.

**Crysophana.**—*C. placida* is rare in the San Bernardino Mountains, where I have beaten it from yellow pines in August.

**Agrilus.**—*A. angelicus* is a rare species described from one specimen taken in Los Angeles County by Coquillett, and a short series from the Santa Cruz Mountains (National Museum Collection). I have taken one specimen in the San Bernardino Mountains in July, at an elevation of 5000 feet, and have seen several specimens taken by Mr. Daggett at Deep Creek, a point a little farther east in the same mountains, at an altitude of 6000 feet. *A. niveiventris* is common on willows throughout the State; specimens taken at Yuma by myself are of a dull green color, while those from the southern maritime region are almost invariably cupreous; I have seen specimens of a brilliant blue color taken by Dr. Fényes in the Shasta region. *A. obolinus* is in Mr. Ulke's collection from "Southern California;" *walsinghami* is found in the extreme northern part of the State: *jacobinus*, San Diego: *politus*, San Bernardino Mountains, on willow: *blundus*,

on flowers at Pasadena in June and July, at Santa Monica (Van Dyke), and at Tehachapi (Wickham): *gibbicollis* is known by a single example taken near San Bernardino in June by Dr. Fényes, and by a single specimen without definite locality in the Horn Collection: *illectus* is found at Pomona and Pasadena by sweeping late in June: *lacustris*, I am told by Dr. Van Dyke, has occurred on white oak at Newhall (Los Angeles County).

**Taphrocerus.**—*T. gracilis* is introduced on the authority of Van Dyke; Los Angeles County.

#### LAMPYRIDÆ.

**Calopteron.**—*C. reticulatum* is recorded from "California" by Le Conte.

**Eros.**—*E. latus* is rarely taken in the San Bernardino Mountains.

**Calochromus.**—*C. dimidiata* occurs in the Mariposa region and in Siskiyou County.

**Matheteus.**—*M. thevenetii* is from the Mariposa region.

**Ellychnia.**—*E. californica* is a common and widely dispersed species: *corrusca* is less common than the last, specimens in my collection having been found at Riverside in May.

**Pyropyga.**—*P. fenestralis* is not rare; specimens are known to me from Riverside, Pomona, Needles and Yuma: *indicta* is described from Lake Tahoe.

**Lamprorhiza.**—*L. riversi* occurs in Sonoma County.

**Microphotus.**—*M. angustus* is from central California.

**Pterotus.**—*P. obscuripennis* is exceedingly scarce in our district, and I have seen but one example from near Pomona; it is more common further north.

**Zarhipis.**—I have seen one example of *integripennis* from Palm Springs (April 10), and one taken at light, at Pasadena, April 14; this and *ruficollis*, *piciventris*, and *riversi* are all taken about San Francisco; but it is very doubtful if we have as many species as names.

**Mastinocerus.**—One example taken at light, at Riverside, June 7, is referred to *opacus*.

**Cenophengis.**—*C. debilis* has been found at Los Angeles and at Pasadena (July); it is very rare.

**Podabrus.**—*P. comes* is a rather common species in the southern sierras and in the foot-hills near Pomona and Pasadena. An apparently undescribed species is not rare at Pasadena in May, occurring usually on willows. *P. latimanus* and *P. tomentosus* are both recorded from "Cal.," but it is not likely that they will be found with us as the genus is as a rule northern in distribution: *binotatus* is said to occur at San Mateo: *xanthoderus*, Lake Tahoe: *macer*, San Mateo and Sylvania: *tejonicus*, Tejon: *bolteri*, Lake Tahoe: *mellitus*, Sylvania, Geysers, and Lake Tahoe: *cavicollis*, San Francisco to Lake Tahoe: *corneus* is described from "Cal." without mention of exact locality. *P. lutosus* is the female of *cavicollis*. There are several undescribed species in my collection.

**Silis.**—*S. cava* occurs not rarely in the foot-hills and mountains near Pomona: *filigera* is scarce; I have found it at Riverside in early spring: *flavida* occurs at Lake Tahoe: *lutea* is found about San Francisco.

**Telephorus.**—*T. consors* is very abundant at Pomona, Riverside, Pasadena, etc., flying in the early evenings of April and May, and often coming to light. A form which is with some doubt referred to *notatus* is not rare at



Pomona and Pasadena, by beating willows in May: *ingenuus* is not common; Pasadena, May: *ochropus* is described from San Diego: *lautus* is reported from various localities in our region: *fraxini* is northern: *transmarinus* and *grandicollis* occur in the middle and northern Sierras.

**Polemnius.**—*P. languidus* is described by Horn from the vicinity of San Diego.

**Ditemnus.**—*D. obtusus* occurs at Riverside in May, but is uncommon.

**Malthodes.**—*M. laticollis* and *M. fuscus* are common at Pomona, Riverside, Pasadena, etc., on flowers and various trees and shrubs: *fragilis* is described from "Cal.;" I have not yet recognized it.

#### MALACHIDÆ.

**Collops.**—*C. marginicollis* was described from San Diego, in which vicinity it appears to be commoner than elsewhere; I have taken it sparingly at Pomona in April, May and September: *cribrosus* occurs on the ocean beaches and on the adjacent sand dunes; it is especially common during April and May at Redondo, and I have also taken it in August at Santa Monica: *argutus* is not rare at Pomona and Pasadena, and I have seen specimens from San Diego and also from Sonoma and Lake counties: *punctulatus* is perhaps our rarest species, single specimens from Pomona and San Diego being all I have seen from our district: *marginellus* is not common; I have taken it at Pomona and San Bernardino in August: *insulatus* is described from California, but I have seen no specimens from our section. Two females of an apparently undescribed species have been taken at Pomona.

**Endeodes.**—The species of this genus are found only on the sea beaches. *E. abdominalis* is at times quite common at Redondo (April and May): *collaris* is much rarer, in the south at least, and I have found but a single example on Catalina Island: *basalis* has not yet been recorded from our district.

**Malachius.**—*M. auritus* and *M. thevenetii* are not rare on willow blossoms in early spring; they are generally diffused: *inornatus* occurs at Pomona from March to May: *acutipennis* is not rare at the same place a little later in the season: *prolixicornis* has occurred only at Pasadena, June and July: *directus* is known from Los Angeles County and from the San Bernardino Mountains: *nigrinus* and *pristinus* are, so far as known, found only in the Sierras, at an altitude from 5000 to 7000 feet; the latter has been taken only in Los Angeles County; *nigrinus* ranges from near San Bernardino to Lake Tahoe. The species are all found on flowers, and are to be obtained by sweeping and beating. In addition to the above, the following are found in other parts of the State: *biguttulus*, *mirandus* and *spinipennis* at Fort Tejon; *macer* at Lake Tahoe; and *mixtus* is from "Cal." without more definite statement of locality.

**Tanaops.**—*T. abdominalis* is rather common on flowers at Pomona, Pasadena, etc., in June: *longiceps* is said by Horn to occur with the preceding, but I have not yet detected it in our district; it is not rare near Lake Tahoe and at Sylvania.

**Microlipus.**—*M. laticeps* occurs from San Francisco to San Diego; I have seen specimens from San Diego, Pasadena and Long Beach in our district; it is not common: *longicollis* is still scarcer, and I have met with it

but once, at Redondo in April: *maerens* is described from "California."

**Pseudebæus.**—*P. bicolor*, or a closely allied species, is generally distributed and not uncommon.

**Attalus.**—The various species occur on flowers, and when found may usually be taken in numbers. *A. trimaculatus* is not common at Pomona, but seems more so farther north: *rufomarginatus* is common throughout the foot-hills of the Sierras: *oregonensis* occurs in the same localities as the last, but is rare with us: *basalis* is rather common at Pomona, Pasadena and Yuma: *cinctus* and *difficilis* are described from Yuma: *lobulatus* is common at various places both east and west of the mountains: *transmarinus* is known only from San Clemente Island. There are in my collection several apparently undescribed species from various parts of the State.

**Pristoscelis.**—Specimens of *P. grandiceps* have been taken at Pasadena (July) by Dr. Fényes; it is a rare insect.

**Eudasytes.**—Casey described *E. ursinus* from "Southern California," and *amplus* from California simply.

**Asydates.**—*A. rufiventris* is described from Santa Barbara: *explanatus* is not rare in cactus blooms; Palm Springs (April), and at Yuma in March.

**Trichochrous.**—According to Casey this genus should be used for the greater number of the species now referred in our list to *Pristoscelis*. He describes very many new species in his late revision; but there are not a few in my collection still evidently without names. They occur abundantly on flowers, as do the most of the species of allied genera. *T. antennatus* occurs at Riverside

(May) if specimens in my collection were properly identified by Casey. *T. griseus*, *T. subcalvus*, *T. suturalis*, and *T. politus* have been announced from San Diego only: *fuscus* occurs in San Diego County, and has been once taken by me at Pomona (April 30): *umbratus* is common in cactus blossoms, at Palm Springs (April). A large series taken by me makes it certain that *fulvovestitus* is the male of *propinquus*, and that both are synonyms of *umbratus*. This species was described from Fort Crook in northern California, and has been collected in the Tempe Desert of Arizona by the late Dr. Griffith of Philadelphia. *T. suffusus*, *T. lobatus* and *T. barbaræ* are described from Santa Barbara; specimens taken by Mr. Daggett in Bear Valley (San Bernardino Mountains), at an elevation of 6400 feet, are identified by Casey as his *seriellus*, described from Utah: *brevicornis* is found at Pomona, Pasadena, Redondo, etc., in spring, being especially common near the coast: *vilis* is not different: *conspersus* is common at Pomona, Riverside, San Diego, etc., March to May: *squalidus*, *sordidus* and *tejonicus* are more or less common at San Diego, Riverside, Pomona and Pasadena; they appear to run into one another, and it is not unlikely that they represent but a single species: *crinifer* occurs at Riverside (May): *fulvescens*, San Diego and Pomona: *nigrinus*, Pomona and Pasadena: *pedalis* and *punctipennis*, Santa Catalina Island, and *ænescens*, "San Diego and the islands off Santa Barbara." The following are described from Southern California without definite locality: *compactus*, *apicalis*, *testaceus*, *prominens*, and *cuspidatus*; *insignis* is from the southeastern region. Of the species occurring in the State outside of our district, *quadricollis* is from Fort Tejon; *brevipilosus* from the central region; *innocens* from the middle Sierras; *fraternus*,

*cylindricus*, *laticollis*, *transversus*, *fulvotarsis*, *separatus*, *indigens*, and *hystrix* from the middle coast region; *oregonensis*, *nubilatus*, *fallax*, *sexualis*, and *sonomæ* from the north. The following, recently described by Casey, have no locality other than California: *brevis*, *agrestis*, *sinuosus*, *discipulus*, *femoralis*, *fimbriatus*, *pruinosis*, *sobrinus*, *mucidos*, *curticollis*, *remotus*, *tectus*, *villosus*, *irrasus*, *rusticus* and *stricticollis*.

**Adasytes.**—*A. luciniatus* is described from San Diego.

**Listrus.**—*E. interruptus* is recorded by Le Conte as occurring on Santa Cruz Island, but I am inclined to believe the reference incorrect; the species is common about Lake Tahoe: *difficilis*, *luteipes* and *famelicus* are common and generally dispersed in our district west of the Sierras: *obscurus*, *extricatus* and *balteellus* are recorded only from San Diego: *definitus* is thus far known only from Pasadena, where a short series was taken in the sweeping net in April by Dr. Fényes: *motschulskii* and *canescens* occur in the central region and north: *rotundicollis*, *ampliocollis*, *maculosus*, *pardalis* and *incertus* are recorded from the middle coast region: *interstitialis*, *densicollis*, *subcaneus*, *tritus*, *variegatus* and *montanus* are from the north: *fidelis* is from "California." Several other forms in my collection are apparently undescribed.

**Dasytellus.**—*D. inconspicuus* is plentiful at Palm Springs (April).

**Dasytes.**—*D. dissimilis* is found in the higher parts of the Sierras from San Bernardino to Lake Tahoe: *clementæ* is known only from San Clemente Island: *macræ* is described from "So. Cal.:" *pusillus*, from San Diego: *musculus* is not rare at Pomona and Pasadena (June):

*lineellus* is rare, mountains of Los Angeles County: *obtusus*, *breviusculus*, and *seminudus* are found in middle California; the first named I have taken from coniferous trees at Lake Tahoe: *expansus*, *nitens* and *cruralis* are northern: *fastidiosus* and *minutus* are described without definite locality.

**Dasytastes.**—*D. catalinae* and *D. insularis* have thus far been found only on Santa Catalina Island. The former is quite common on various flowers in July; the latter has been taken but once (Aug. 1). *D. bicolor* is a common species at Pomona in May and June. A small species found by me on oaks in the San Bernardino Mountains is referred to *remissus*. *D. dispar* occurs north of San Francisco, and *otiosus* is without specific locality.

**Eschatocrepis.**—*E. constrictus* is everywhere common.

**Allonyx.**—*A. sculptilis* is an uncommon species; I have taken it at Pomona in June, and at Pasadena early in April. Casey describes *denudatus* from north of San Francisco, and *disjunctus* from "California."

**Vectura.**—*V. longiceps* is described from Yuma, Arizona; it is probable that the specimens were taken on the California side of the river.

**Eurelymis.**—*E. flavipes*, San Bernardino Mountains, elevation 6,000 feet, July 5 (Daggett); two examples only were taken. This species occurs also in the middle Sierras.

#### CLERIDÆ.

**Elasmocerus.**—*E. californicus* is an uncommon insect of which I have seen specimens taken from Sonoma to Los Angeles counties.



**Cymatodera.**—*C. puncticollis*, San Bernardino Mountains, on pines; not common; Yuma and country along the Colorado River: *californica*, a rare species, of which I have seen only two examples from Southern California, one from the San Bernardino Mountains, the other from the mountains near Pomona, September: *morosa*, specimens so referred have been taken in Los Angeles County: *punctata*, along the Colorado River: *fuscata*, occurs at Yuma: *undulata* var. *balteata*, Yuma (fide Fuchs): *angustata*, Yuma and various other localities in Southern California: *ovipennis*, Pomona, November; San Bernardino Mountains, July; Catalina Island, July; var. *pilosella*, San Diego.

**Aulicus.**—*A. nero*, "Collected by Gabb in the South Coast Range" (Horn).

**Trogodendron.**—*T. edwardsii*, San Diego County (Fuchs).

**Trichodes.**—*T. ornatus*, rather common in most every locality; the variety *tenellus* is also widely diffused, but aside from Yuma I know of no record of its capture elsewhere in our district: *bimaculatus* is found in the north.

**Clerus.**—*C. quadrisignatus* occurs along the Colorado River: *eximius* is rare at Pasadena, April; found on willows (Fényes): *abruptus*, specimens which seem to be a slight variety of this species have been seen from Owens Valley: *mæstus*, not rare in the San Bernardino Mountains, on coniferous trees: *sphegeus*, occurs at Lake Tahoe and north.

**Thanasimus.**—*T. repandus*, Mendocino County (Van Dyke).

**Hydnocera.**—*H. robusta*, Owens Valley (Horn): *scabra*, rather common throughout the southern Sierras and the adjacent region: *discoidea*, Palm Springs, April; Colorado River (Le Conte): sp. near *pallipennis*, one example, Pomona: *bicolor*, Colorado River (Le Conte).

**Chariessa.**—*C. elegans*, one example taken at an altitude of over 6,000 feet in the Sierras, in Los Angeles County, by Mr. Daggett, is the only instance known to me of the capture of this beautiful insect in Southern California. *C. dichroa*, Los Angeles County (Coquillett), is described from Sacramento.

**Cregya.**—*C. fasciata* is described from San Diego. A specimen with entirely piceous elytra taken by me in the San Bernardino Mountains is probably a mere color variety. The species also occurs in northern California.

**Lebasiella.**—*L. maculicollis* occurs sparingly at Pasadena and Pomona late in May and early in June.

**Corynetes.**—*C. rufipes* occurs everywhere: *ruficollis*, the only specimens seen from our district were taken early in June on San Clemente Island.

## PTINIDÆ.

**Trigonogenius.**—*T. farctus* is not rare at San Francisco, where it has distinguished itself as an herbarium pest.

**Sphæricus.**—*S. gibbioides* also occurs at San Francisco.

**Ptinus.**—The species of this genus are in some confusion. Specimens of two species which have been considered to be (and probably rightly so) *interruptus* and *verticalis* are found quite commonly by beating live-oaks and occasionally other trees or shrubs, October to

March. Specimens of *brunneus* are labeled "Cal." in the Le Conte Collection, and Horn cites *pygmaeus* Gorb. from California in his list of the Coleoptera of Lower California.\*

**Hedobia.**—*H. granosa*, not rare, Pomona, Pasadena, Ojai Valley; March to April; chiefly found on live-oak.

**Ernobius.**—*E. debilis*, Santa Barbara and Santa Cruz Island (Le Conte). An apparently undescribed species near *punctulatus* has been taken abundantly on pines in the San Bernardino Mountains. Another undescribed species occurs also on pines about Lake Tahoe. *E. punctulatus* is described from the northern coast region, and *alutaceus* from "Cal."

**Ozognathus.**—*O. cornutus*, Pomona, Pasadena, Riverside, February to May; *misellus*, San Diego. The original specimens of the former sent to Dr. Le Conte were said to have been hatched from galls.

**Xestobium.**—One example of an undescribed species, taken at Pomona in May: *affine* is recorded from "Vancouver and California."

**Oligomerus.**—Three species doubtfully congeneric are for the present referred here. One is not rare on Catalina Island; found by beating *Heteromeles arbutifolia*; a second occurs on *Quercus agrifolia* at Pasadena in June; the third is represented by a solitary example found dead among the sand-dunes near the beach at Santa Monica.

**Sitodrepa.**—*S. panicea* is more or less frequent everywhere in houses. I raised a nice crop once from a lot of European Coleoptera received from France.

---

\* Proc. Cal. Acad. Sci., 2d Ser., Vol. IV, 1894, p. 332.

**Ctenobium.**—One example of an undescribed species taken at Riverside at electric light.

**Ptinodes.**—*P. setifer*, San Diego; the type alone is known to me.

**Hadrobregmus.**—*H. gibbicollis*, Pomona, San Bernardino Mountains; June and July; not common. One specimen of an undescribed species near *foveatus* was taken at Lake Tahoe.

**Trichodesma.**—*T. cristata*, middle California.

**Anobium.**—*A. quadrulum*, Lake Tahoe and north.

**Trypopitys.**—*T. punctatus*, Los Angeles, Riverside, Pomona, Pasadena, etc.; July and August: *temulineata*, one example, Catalina Island.

**Petalium.**—*P. bistriatum* is not rare in the southern Sierras.

**Theca.**—*T. striatopunctata*, "Cal." (Le Conte).

**Vrilletta.**—The three species on our list—*convexa*, *expansa* and *murrayi*—are described by Le Conte from California without definite locality; I have seen only *convexa* from our district, but all may occur there; *expansa* is not uncommon near San Francisco.

**Xyletinus.**—Specimens both black and brown, identified by Dr. Horn as *lugubris*, are not rarely taken by beating in the foot-hills near Pomona in May and June. Le Conte mentions the taking by Crotch of a species at Calaveras which scarcely differs from *fucatus*, and is so referred by him.

**Catorama.**—*C. frontalis* may be taken in moderation by beating *Quercus agrifolia*; Pasadena, May and June.

**Hemiptychus.**—*H. latus* occurs on the Southern California sea beaches, but not plentifully: *pusillus*, Yuma

(Le Conte): *luteolectus*, Riverside and Yuma, July; Palm Springs, April: *palliatu*s, not rare on a species of *Dalea* at Yuma early in June. Our commonest species, which occurs pretty nearly everywhere, and most frequently on oaks, is not yet identified; it is possibly not different from *obsoletu*s, which was described from Cape San Lucas, but it would not be proper to so call it without careful comparison with the type. There appear to be two other allied species in my collection from Pomona and Pasadena, but they are not sufficiently strongly characterized to warrant their description in advance of a general synopsis of the genus. *H. integer* is represented by two examples taken by me at Lake Tahoe.

**Cœnocara**.—*C. californica*, one example from the Sierras near Pomona, September: *occidens*, described by Casey from central California.

**Ptilinus**.—*P. basalis*, San Diego (Le Conte): *ramicornis* has been beaten from *Quercus engelmanni* at Pasadena late in May, and also occurs in the San Bernardino Mountains: *flavipennis* occurs in the southern Sierras, in Los Angeles and San Bernardino counties: *acuminatus* is described from the Santa Cruz Mountains.

**Euceratocerus**.—*E. pleuralis*, *E. macer*, and *E. saginatus* are described by Casey from California; the first mentioned from the Santa Cruz Mountains, the others without definite locality.

**Sinoxylon**.—*S. sericans*, Colorado River: *sextuberculatum*, Yuma; "probably depredates on mesquite" (Horn): *declive*, Pomona, August and September; numerous specimens taken on the windows of a winery; one specimen was taken on San Nicolas Island, May 23: *suturale*, rare at Pasadena (Fényes).

**Bostrychus.**—A single specimen of *B. californicus* was secured in the San Bernardino Mountains in July.

**Amphicerus.**—*A. fortis*, Colorado River: *punctipennis*, Riverside, Redlands, Pomona, Long Beach; June to October; often taken at electric lights: *teres*, Yuma; unknown to me.

**Dinapate.**—*D. wrightii* is a mammoth Bostrychide known only from the cañons leading up from the western borders of the Colorado Desert, at and near Palm Springs, where it depredates on the Washingtonia palms (*Neowashingtonia filifera*). It was originally discovered by Mr. W. G. Wright, and has since been found by Mr. H. G. Hubbard, whose most interesting letters to Mr. E. A. Schwarz, describing the rediscovery, with much information concerning the larval habits, have been recently published by Mr. Schwarz.

**Dinoderus.**—*D. pacificus*, San Bernardino Mountains, not rare: *sobrinus*, from the same locality; scarce: *truncatus*, "California" (Horn).

**Polycaon.**—*P. stoutii* is common throughout maritime Southern California; once taken in numbers from beneath the loose bark of eucalyptus in early summer: *confertus* is also widely diffused; it depredates on grape vines. Of *megalops* two examples only have been seen, one of which was taken at light at Pomona, September 25; the other is in the collection of Dr. E. C. Van Dyke.

**Psoa.**—*P. maculata* and *P. 4-signata* are both rare insects in Southern California.

**Lyctus.**—*L. planicollis*, common throughout our district from Yuma to the Coast: *parvulus*, rare, Pomona, June: *californicus*, Yuma (Casey): *curtulus*, described from "California."



## CUPESIDÆ.

**Priacma.**—I have seen one example of *serrata* in the collection of Professor Rivers, said to have been taken at Grass Valley in the Middle Sierras.

**Cupes.**—*C. lobiceps*, San Diego; apparently very rare.

## CIIDÆ.

**Cis.**—*C. versicolor*, *C. vitulus*, and *C. duplex* are all common in dry, woody fungi at Pomona, Pasadena, Ojai Valley, and probably throughout maritime Southern California. Casey also describes *illustris*, *impressa*, and *soror* from northern California, and *macilenta* and *hystricula* from Lake Tahoe.

**Orthocis.**—*O. aterrima*, Alameda County.

**Plesiocis.**—*P. cribrum* is not uncommon in the Sierras from Calaveras County to San Bernardino County.

**Ennearthron.**—*E. grossulum* and *E. convergens* are described from Southern California. Specimens of what is doubtless one of these species occur plentifully in various localities near Los Angeles. *E. californicum* is said to especially frequent the northern coast region, while *discolor* is described from Sonoma County.

**Octotemnus.**—*O. denudatus*, San Francisco to Vancouver.

## SPHINDIDÆ.

**Odontosphindus.**—*O. clavicornis*, one example, San Bernardino Mountains, July.

## LUCANIDÆ.

**Sinodendron.**—*S. rugosum* is a rather common species from the Santa Cruz Mountains, north, but I have

no other record of its occurrence within our district; El Monte (Los Angeles County) teste Rivers. I know of no record of any other species of this family having occurred in our district. In the genus *Platycerus*, *oregonensis*, *agassii*, *californicus*, *parvicollis*, *pacificus*, *thoracicus* and *chalybæus* are all Californian. *P. oregonensis* has occurred as far south as Tejon, but the true habitat of the genus is more northern. *Ceruchus punctatus* and *C. striatus* occur in middle and northern California.

### SCARABÆIDÆ.

**Canthon.**—*C. simplex* is widely diffused and generally common; the variety *humeralis* is rare. It is probable that the varieties *corvinus* and *militaris* occur in our region but I have no specific reference to the fact. *C. lævis* and *C. perplexus* are recorded respectively from "So. Cal." and "Cal.," but there can be little doubt that they were taken in the desert regions of the south-east.

**Copris.**—*C. mæchus*, Los Angeles; one example taken by Dr. W. J. Karlsioe (fide Linell).

**Oniticellus.**—*O. californicus*, Shasta region; very scarce.

**Ægialia.**—*A. conferta*, Pomona, Riverside, etc., especially in early spring, when it may frequently be taken on the wing: *latispina* occurs with the preceding and is of about equal frequency: *crassa*, not rare along the seacoast (Santa Monica, Redondo, Monterey): *cylindrica* occurs in Marin County: *lacustris*, middle Sierras: *blanchardi*, Mendocino County.

**Psammodius.**—*P. nanus*, a little species sometimes seen flying about the streets of Pasadena in large numbers on warm evenings; September to May; it is widely

diffused: *cælatus* occurs only along the sea-shore; it is tolerably common at Redondo and Santa Monica.

**Pleurophorus.**—*P. cæsus*, Lake County (Van Dyke).

**Rhyssemus.**—*R. californicus*, Riverside, April; not common.

**Atænius.**—*A. desertus*, Yuma: *abditus*, everywhere abundant: *gracilis*, Pomona, Palm Springs, Yuma; widely diffused but much less common in California than the preceding: *californicus*, San Bernardino, Yuma; common at the latter place in July: *lobatus*, Palm Springs; not common. *A. oblongus* is described from "California," but I am unable to ascertain the exact locality.

**Aphodius.**—*A. granarius* is quite common at Pomona from January to April; it is generally diffused: *lividus*, Yuma, Pomona, Los Angeles; not frequent: *rugifrons*, common under rubbish at Pomona after the first soaking rain in Nov. or Dec.; seldom seen later than Feb.: *consociatus*, Bear Valley (San Bernardino Mountains), June (Daggett): *subcæneus*, Los Angeles County; not common: *alternatus*, Yuma (Horn): *cribratus*, one example taken at or near Santa Monica by Mr. Max Albright: *ochreipennis*, Owens Valley (Horn): *rubidus*, rather common at Riverside, Pomona, Pasadena, etc., from Dec. to May: *militaris*, rare at Pomona and Pasadena, May: *coquilletti*, Los Angeles County (Coquillett): *luxatus*, Santa Barbara and Long Beach, Feb. to April; rare: *ungulatus*, taken thus far only during Dec. and Jan. at Pomona and Pasadena; not common: *pardalis*, Ojai Valley, March; rare in Southern California but commoner farther north. Numerous other species occur within the State limits. Of these, *conspersus*, *congregatus*,

*deutus*, *pectoralis*, *nevadensis* and *opacus* inhabit the region to the north of San Francisco: *gentilis* and *inutilis* occur about San Francisco: *vittatus* at San Francisco and at Lake Tahoe: *sparsus* in the Mariposa region, and *ovipennis* at Fort Tejon. Since the above was written a specimen of *vittatus* has been taken at Pasadena by Dr. Fényes (February).

**Ochodæus.**—*O. californicus*, Pomona and Pasadena; very scarce; four specimens only are known, taken on the wing in April and May.

**Pachyplectrus.**—*P. laevis*, described from Santa Barbara. I have seen no examples from California except the type, but have seen two or three specimens which were taken at Phoenix, Arizona, by Dr. H. G. Griffith of Philadelphia.

**Bradycinetus.**—*B. serratus*, Yuma: *horni* is rare in the country about San Francisco.

**Odontæus.**—*O. obesus* occurs near San Francisco.

**Geotrupes.**—*G. occidentalis*, exceedingly scarce, occurring in the region to the east of Visalia.

**Pleocoma.**—*P. puncticollis*, San Diego County (Julian?): *fimbriata*, *behrensii*, *hirticollis*, *staff*, *rickseckeri* and *conjungens* all inhabit the central part of the State from San Francisco to the Sierras, but are confined to hilly or mountainous districts.

**Trox.**—*T. suberosus*, one example, San Bernardino: *punctatus*, one example taken at Yuma, July: *gemma-latus*, San Diego County, Pomona; not common: *atrox*, Pasadena, Santa Monica, Riverside; occasionally occurs in numbers, but generally rarely seen: *fascifer*, taken by Mr. Fuchs near San Francisco.

**Amphicoma.**—The only specimens of *edwardsii* from Southern California that I have seen were taken by Dr. E. C. Van Dyke in the Santa Monica Mountains. *A. canina* is not infrequent in the Southern Sierras. Professor Rivers reports finding a single example of *ursina* on the sand-dunes at Santa Monica; it is not rare in similar situations near San Francisco: *cooperi* occurs at Sacramento in early July: *rathvoni*, the middle Sierras.

**Oncerus.**—*O. floralis*, San Diego County, Palm Springs; occurs on flowers during April at the latter locality.

**Hoplia.**—*H. sackenii*, San Diego (Le Conte); Sylvania (Ricksecker): *callipyge*, not abundant at Pomona, on flowers of greasewood; May and June: *pubicollis*, not rare at Riverside during April and May, on willows along the Santa Ana River: *dispar*, common from Lake Tahoe to Siskiyou County. Specimens of two apparently undescribed species have been taken by Dr. Fényes and myself at Castle Crag and Lake Tahoe respectively.

**Gymnopyge.**—*G. hoplieformis*, Mojave Desert (Coquillett).

**Dichelonycha.**—*D. crotchii*, San Bernardino Mountains, July; more common in the middle Sierras: *fuscata*, plentiful in the foot-hills near Pomona, etc., on flowers of greasewood; June: *truncata*, Pomona, Palm Springs, San Bernardino Mountains; April to July: *pusilla* occurs with the two preceding species in the foot-hills near Pomona, but unlike them is also frequently taken within the town itself, especially about roses: *pallens* and *clypeata* are northern and appear to be not often taken: *fulgida* and *valida* are more or less

common throughout the middle Sierras. There are also in my collection three seemingly undescribed species, one of which comes from San Diego and is the smallest species in our fauna.

**Cænonycha.**—*C. rotundata*, Yuma, Riverside, two examples in March; a rare species: *socialis*, specimens I have seen were said to have been taken in San Diego County.

**Serica.**—*S. fimbriata* and *S. mixta* are both more or less abundant throughout Southern California in early summer, when they are often seen on the wing in the early evening: *elongatula*, several specimens taken under dead leaves in Ojai Valley, March: *alternata*, Los Angeles and San Diego counties; less frequent than *mixta* or *fimbriata*: *anthracina* occurs in middle and northern California: *valida* is probably not different.

**Plectrodes.**—*P. riversi*, generally not common but occasionally attracted in numbers to light; Pomona, July: *carpenteri*, Los Angeles and Mojave Desert: *palpalis*, specimens so called are occasional in the vicinity of Los Angeles: *blaisdelli* is not uncommon near San Diego: *squamosa*, from San Bernardino: *fossiger* and *pistoria* are described by Casey from Los Angeles County; the latter has been taken in some numbers at Santa Monica by Mr. Rivers: *pubescens* was described by Horn from specimens taken at Visalia; it is possible that we have more names than species.

**Orsonyx.**—*O. anxius*, Needles and Yuma (Wickham).

**Diplotaxis.**—*D. mærens* and *D. tenuis* are described from the desert regions of San Diego County: *subangulata* is occasionally abundant at Pomona, at light. Several specimens of an undescribed species were taken at



light, at Palm Springs. I have seen a few specimens of a larger species taken in Placer County that seems to be undescribed.

**Lachnosterna.**—*L. errans*, middle California; described from Contra Costa: *lenis*, desert regions of the southeast.

**Listrochelus.**—*L. mucoreus*, Yuma.

**Polyphylla.**—*P. decemlineata* and *P. crinita* are both fairly common; the former at lights in the towns, the latter at altitudes of 4,000 feet and over in the Sierras.

**Thyce.**—*T. marginata*, San Diego County. I have seen one specimen taken at Oceanside by Mr. F. D. Twogood.

**Phobetrus.**—*P. comatus* is generally scarce; once taken in some quantity at electric lights, in Pasadena, April; occurs also on Catalina Island, San Diego County, and in the San Joaquin Valley.

**Anomala.**—One example of *centralis* was taken at Palm Springs.

**Cotalpa.**—*C. ursina*, very abundant at Pomona, San Bernardino, etc., in spring, flying by day and often seen in quantities clinging to the cypress hedges about the city. *C. granicollis* is reported as having been taken in the Argus Mountains by the Death Valley Expedition.

**Cyclocephala.**—The species of this genus are to be seen flying in the evening about the city streets, and are sometimes attracted to lights in large numbers. *C. immaculata*, *C. longula*, and *C. villosa* are most common: *hirta* is scarce: *dimidiata* while not frequent at light is not rare in the blossoms of *Datura metelloides*.

**Ligyris.**—*L. gibbosus* is abundant and widespread.

**Euphoria.**—*E. verticalis*, from southeastern region; not common.

**Cremastochilus.**—*C. wheeleri*, described from Eldorado County, but said by Fuchs to have been taken by Dunn in San Diego County: *ineptus*, Yuma (fide Fuchs): *schaumii*, San Diego, Pasadena, not frequent: *westwoodi*, Owens Valley: *pilosicollis*, San Bernardino, April; more common farther north: *crinitus*, San Diego, collected by Dunn (fide Fuchs): *planatus* occurs in the Coast Range south of San Francisco: *angularis*, about San Francisco and in the middle Sierras.

**Valgus.**—*V. californicus*, central and northern portions of the State; rare.

### SPONDYLIDÆ.

**Parandra.**—I have not yet seen native living specimens of *brunnea*, but once found the remains of several about sycamore stumps at Pasadena.

**Spondylis.**—*S. upiformis* is at times not uncommon about pines in the San Bernardino Mountains.

### CERAMBYCIDÆ.

**Ergates.**—*E. spiculatus* is quite common in the southern Sierras above 4,000 feet, where the larva infests the roots and stumps of various coniferous trees; it is occasionally taken at light in the valleys.

**Mallodon.**—Specimens of *melanopus* have been taken under mesquite bark at Indio, and on the wing at Yuma, on July evenings.

**Derobrachys.**—One example of *geminatus* was seen at Palm Springs.

**Prionus.**—*P. californicus* is rather widely diffused; it is not rare at electric lights in the towns, also in the

foot-hills and at moderate altitudes in the Sierras. The larva lives in decaying roots of live-oak (*Quercus agrifolia*).

**Trogosma.**—*T. harrisii* is not uncommon in the middle Sierras. The larva is reported as depredating on *Pinus contorta*. *T. pilosicornis* is described from Mt. Diablo, near San Francisco.

**Aseum.**—Specimens referred to *nitidum* have been taken at an altitude of 6,000 feet in the Sierras in June; larva has been found in decayed *Pinus insignis* (Rivers); *atrum* has been taken in the middle Sierras and on Mt. Whitney, at 11,000 feet altitude (Daggett).

**Tetropium.**—*T. velutinum*, labeled "Cal.," must be northern.

**Decentrus.**—*D. blüthneri*, redwood forests, north of San Francisco.

**Hylotrupes.**—*H. amethystinus*, San Bernardino Mountains, Bear Valley; rare (Van Dyke); *ligneus*, common; depredates on pine. A nearly black variety has been taken on wood-piles in the middle Sierras (Placer County) by Mr. Van Dyke, who writes that this melanotic form is found only on fir wood-piles that are situated in the forests and are more or less in the shade, while the type-form occurs on pine and spruce wood-piles in sunny situations.

**Phymatodes.**—One example of *blandus* was beaten from pine in the San Bernardino Mountains: *obscurus*, San Diego, on live-oak from the branches of which it has been bred by Dr. F. E. Blaisdell; *decussatus*, one example beaten from *Quercus engelmanni* at Pasadena, May 31; *juglandis*, Los Angeles County; depredates on *Juglans*

*californica*: *aneus*, *nitidus*, and *vulneratus* occur in the central and northern parts of the State: *obliquus* in Santa Clara County.

**Callidium.**—*C. antennatum*, Mt. Wilson, Los Angeles County (Van Dyke): *hirtellum*, San Bernardino Mountains; not very common: *vile*, described from "Cal.;" it is doubtless from the north. Le Conte mentions a race of *janthinum* from California.

**Xylocrius.**—*X. agassizii* and *X. cribratus* are found in the middle and northern Sierras.

**Malacopterus.**—*M. lineatus*, Colorado River, above Yuma.

**Ceme.**—I have seen one example of *gracilis* from Ontario (Los Angeles County) and one from Santa Catalina Island; bred from the dead wood of *Quercus agrifolia* at Poway (Blaisdell): *costata* is recorded from "Cal."

**Eucrossus.**—One example of *villicornis* was taken in the San Bernardino Mountains.

**Haplidus.**—*H. testaceus*, "Cal."

**Brothylus.**—One example of *gemmulatus* was taken on Mt. Wilson (Van Dyke).

**Romaleum.**—The only specimen of *simplicicollis* seen by me was taken on Santa Catalina Island; it is a widely diffused species and should occur elsewhere: *seminitidum*, one example in my collection, from the Colorado River.

**Elaphidion.**—*E. albofasciatum*, taken in the foot-hills along the southern border of the Mojave Desert (Coquillett): *imbelle*, Poway (San Diego County); bred from the dead wood of live-oak by Dr. Blaisdell, who says that the beetles are common beneath bark in August.

**Aneflus.**—One example of *linearis* taken on Echo Mountain, Los Angeles County.

**Pæcilobrium.**—*P. chalybæum* was taken on flowers of *Ceanothus*, Pasadena, April. Linell has described *rugosipenne* from "Cal."

**Hybodera.**—*H. debilis*, one example, Echo Mountain, Los Angeles County.

**Callimus.**—*C. cyanipennis*, Los Angeles County, on *Ceanothus* bloom. *C. ruficollis* bears a label in the Horn Collection indicating its occurrence in Southern California; it is not rare about San Francisco.

**Megobrium.**—*M. edwardsii* is described from Santa Rosa Island. I have one example beaten from live-oak near Pomona, Oct. 27.

**Molorchus.**—*M. longicollis*, Los Angeles County, on flowers.

**Callimoxys.**—*C. fuscipennis*, Los Angeles County (Van Dyke); very abundant further north.

**Holopleura.**—*H. helena* is very rare in Southern California, but has been bred in quantity from dead twigs of *Umbellularia californica* from near San Francisco, by Professor Rivers; Santa Monica (Albright).

**Rosalia.**—One example of *funebis* was sent me from San Diego County. Mr. W. G. Wright of San Bernardino writes me that he has taken it locally, but it is certainly much more common further north. Professor Rivers records the larva from "decaying *Umbellularia californica* among the mycelia of some fungus."

**Dendrobius.**—*D. mandibularis* is common on the wing in early evening, about the willows along the Colorado River, at Yuma, in July.

**Lissonotus.**—*L. multifasciatus*, "Cal., Ariz."

**Tragidion.**—*T. annulatum*, Pasadena, June; Pomona, September; San Bernardino Mountains (Van Dyke); Santa Monica (Albright); rare. *T. armatum* has been taken at Newhall (Van Dyke), while sucking sap from stalks of flowering *Yucca whipplei*.

**Purpuricen**us.—*P. dimidiatus*, northern California.

**Metaleptus.**—*M. angulatus*, "So. Cal.;" doubtless from the southeastern region.

**Amannus.**—*A. pectoralis*, Yuma (Le Conte.)

**Batyle.**—*B. suturalis*, San Diego County (Fuchs).

**Oxoplus.**—*O. jocosus* occurs rather rarely in the foothill cañons near Pomona; found flying by day along the streams.

**Crossidius.**—*C. hirtipes*, *C. ater*, *C. testaceus*, and *C. punctatus* bear Californian labels in one collection or another, but with the exception of *testaceus*, which, according to Fuchs, has been taken in San Diego County, I have not been able to obtain exact localities. There is not much doubt, however, that they occur only in the eastern and southeastern portions of the State.

**Ischnonemis.**—*I. bivittatus*, Indio (Van Dyke).

**Stenosphenus.**—*S. debilis*, several examples, beaten from willows at Yuma, July.

**Cyllene.**—*C. antennatus*, common in the desert and along the Colorado River; breeds in mesquite: *crinicornis* also occurs in the southeast.

**Calloides.**—*C. lorquini*, Mt. Lowe (Los Angeles County), one example; Lower Soda Springs, Siskiyou County, July (Van Dyke).

**Clytus.**—*C. lanifer*, Santa Barbara; more frequent farther north.



**Xylotrechus.**—*X. nauticus*, San Diego, Los Angeles, Pomona (August). According to Blaisdell, "Abundant about live-oak groves during July and August. The larvæ and pupæ have been taken from the dead branches and trunk of the oak. The beetle is both diurnal and nocturnal in habits, becoming active near the middle of the afternoon and continuing so until late in the evening." The larvæ of this insect have also been found destructive to logs of *Eucalyptus globulus*.\* *X. oblitteratus* is not infrequent about willows, in various localities in Southern California and on Santa Catalina Island, June: *insignis* is the female of this species, as I am informed by Mr. Fuchs, a fact I had suspected, but of which I had no positive evidence: *undulatus* and *annosus* have been found in the north: *plumifrons* occurs near San Francisco, where the larvæ may be found in dead branches of willow (Rivers).

**Neoclytus.**—Several examples of *irroratus* have been picked up on the streets of Pasadena during early September by Mr. Daggett, and I have similarly found it at Santa Barbara late in August: *conjunctus*, *muricatus*, *balteatus* and *interruptus* inhabit the middle or northern parts of the State.

**Atimia.**—*A. dorsalis*, Riverside, May and October; uncommon; occurs also in the Sierras, Placer County, April (Van Dyke): *confusa* has been found in California also, according to Henshaw.

**Desmocerus.**—*D. cribripennis*, "So. Cal.;" the species must be exceedingly rare: *californicus* has been taken in Los Angeles County, on elder, but is, like the preceding, very scarce: *auripennis* has not occurred in our

---

\* See Blaisdell's account, "Insect Life," Vol. V, p. 34.

district so far as I am able to learn, but is not so exceedingly rare farther north; it has been bred from the dead wood of *Sambucus glaucus* (Blaisdell).

**Necydalis.**—*N. barbare*, Santa Barbara; rare (Rivers): *lavicollis*, middle California; taken by Van Dyke and Rivers near San Francisco, where it is found to breed in decayed *Quercus agrifolia*, and in dead *Eucalyptus globulus* (Rivers): *cavipennis* occurs (in the north) at Piedmont, Alameda County, May (Fuchs).

**Ulochætes.**—*U. leoninus*, San Bernardino Mountains; Glenbrook, Lake County, June (Fuchs); Sisson, Shasta County, June, on yellow pine (Van Dyke). The dried remains of the beetles in their burrows in dead *Pinus ponderosa* have been seen by the writer; the beetles emerge in July, but are scarce or at least seldom taken.

**Pyrotrichus.**—*P. vitticollis* was taken by Mr. Fuchs at Laundry Farm, Alameda County, in April.

**Leptalia.**—*L. macilentæ*, northern California; Sonoma County (Ricksecker); on willows, in Mill Valley, Marin County, May (Fuchs).

**Rhagium.**—*R. lineatum* is central or northern; not rare about Lake Tahoe.

**Centrodera.**—One example of *nevadica* from Santa Monica is in the collection of Van Dyke.

**Toxotus.**—*T. vestitus* is not uncommon in the southern Sierras (Van Dyke): *flavolimbatus* is recorded from "Cal.," *nubifer* from Tejon, *lateralis* from near San Francisco.

**Pachyta.**—*P. spurca*, one example from Echo Mountain, Los Angeles County: *liturata*, one example from Mt. Whitney, elevation 8,000 feet (Daggett).

**Anthophilax.**—*A. tenebrosus*, from the southeastern region.

**Acmaeops.**—*A. tumida* is very rare in Southern California; I secured a single specimen on flowers at Pasadena, in April: *falsa*, Pomona, May; Pasadena, May, from between leaves of *Yucca whipplei* (Daggett); Dr. Van Dyke finds it in June on flowers of wild holly. The following species of the genus are found in the middle and northern portions of the State: *subcænea*, *punguis*, *viola*, *longicornis*, *vineta*, *basalis*, *militaris*, *subpilosa*, *variipes*, *prutensis*; the latter from Mt. Whitney, elevation 8,000 to 9,000 feet (Daggett).

**Strangalia.**—*S. delicata*, San Bernardino, Pomona, Pasadena, late May to July, on various flowers; not common.

**Leptura.**—*L. molybdica* is not rare on flowers of *Ceanothus*, Pasadena in April; all specimens found here have the red humeral spot: *læta*, not rare on flowers in the foot-hills of the Sierras, at Pomona, Pasadena, etc., in June; breeds in dead *Quercus agrifolia* (Rivers): *tribalteata*, Mt. Wilson, Los Angeles County; uncommon: *coquilletti*, Los Angeles County: *instabilis*, rare, Pomona, June, in the foot-hills; var. *convexa*, Los Angeles (Fuchs): *sexspilota*, abundant usually on the flowers of greasewood in May and June throughout Southern California: *pernigra*, Mt. Wilson, Los Angeles County (Coquillett): *crassipes*, not common in the San Bernardino Mountains, 5,000 feet elevation, in August; it has been bred from decayed wood of *Umbellularia californica* (Rivers): *valida*, Bear Valley, San Bernardino Mountains; very scarce. Aside from the above, there are numerous species occurring within the State,

all of which—except as noted below—are from the more northern parts: *obliterata*, *soror*, *plagifera*, *rubida*, *subargentata*, *impura*, *vexatrix*, *matthewsii*, *grossa*, central and northern Sierras; *propinqua*, Kern River; *carbonata*, *dehiscens*, *sanguinea*, *latifica* (these last two rare at Kaweah, Tulare County) (Hopping), common farther north; *quadrillum*, *chrysocoma*, *dolorosa*, *behrensii*, Alameda County; *scripta*, *gnathoides*, *cubitalis*. Several undescribed species exist in collections.

**Ophistomis.**—*O. ventralis*, “So. Cal.,” El Taste, Lower California.

**Ipochnus.**—*I. fasciatus* is not uncommon under bark, and by beating, especially *Rhus*. Blaisdell reports larvæ taken from the wood of *Rhus integrifolia* at Coronado, and *Rhus laurina* at Poway. Larvæ have been found by Coquillett in dead and dry apple twigs. The beetles are abroad from February to Sept.

**Monilema.**—*M. spoliatum* has been taken by Mr. W. G. Wright, exact locality not stated.

**Monohammus.**—*M. maculosus* is not rare at Lake Tahoe (Fényes) and north.

**Synaphœta.**—*S. guexi*, rare at Pomona and Pasadena, March and April; it has been bred from dead limbs of *Æsculus californica*, by Rivers.

**Cænopœus.**—*C. palmeri* has been taken in numbers at San Bernardino by Mr. W. G. Wright; breeds in *Opuntia*.

**Hyperplatys.**—The form of *aspersus* described by Casey as *californica*, and which according to Leng is only a variety of Say’s species, is not unfrequently taken about San Francisco. I have myself found it in Marin County in August.

**Acanthocinus.**—*A. obliquus* and *A. spectabilis* are both found in the higher parts of the southern Sierras, but are more frequent in the middle and northern portions of the range.

**Pogonocherus.**—Several examples of *crinitus* have been taken at Pasadena by Dr. Fényes, on live-oak, in June. Professor Rivers reports the larva as occurring in dead branches of *Quercus agrifolia*. *P. mixtus* occurs in the middle or northern regions, and I have found *oregonus* at Lake Tahoe.

**Lypsimena.**—*L. californica* is rare at Pasadena (Fényes); Santa Monica, August (Van Dyke); Yuba County (Fuchs).

**Saperda.**—A form which passes as *mæsta* is occasionally found in Southern California, on cottonwood trees; I have taken it at Riverside, in May.

**Oberea.**—*O. schaumii* is rather uncommon; found on willows in the foot-hills near Pomona in June; it has been called *quadricollis*, a name now regarded as a synonym of the one given above.

**Tetraopes.**—One of the numerous color varieties of *femoratus* is plentiful on milkweed throughout our region in May and June.

**Idæmea.**—One example of *californica* was beaten from live-oak, in June, and others were secured at electric light, at Pasadena, in April.

**Methia.**—Several examples of an unidentified species were taken at electric light, at Pasadena, in September (Fényes); this is possibly *mormona* Linell.

#### CHRYSOMELIDÆ.

**Aulacoscelis.**—*A. purpurea*, Needles (Wickham).

**Donacia.**—No specimens of this genus have yet been found in our region, but the following six species are

known from the central and northern portions of the State: *hirticollis* and *pubescens* from Lake County (Van Dyke); *cincticornis* var. *proxima* and *subtilis*, from Marin County; *pusilla* and *emarginata*.

**Orsodacna.**—*O. atra* comes as far south as Lake Tahoe in the more elevated regions of the Sierras.

**Zeugophora.**—One example of *californica* taken at Pasadena in April (Fényes): *abnormis* I found in some numbers on the leaves of aspens, at Lake Tahoe, in July.

**Syneta.**—One example of *carinata*, Lake Tahoe: *hamata*, northern: *albida*, occurs as far south as Alameda County.

**Thricolema.**—*T. anomala*, Calaveras County.

**Lema.**—*L. nigrovittata* depredates in all stages on *Datura metelloides*; very common.

**Euryscopa.**—*E. subtilis*, *E. vittata* and *E. lecontei* are all on record as having been found in Southern California; they are probably found only in the southeast.

**Coscinoptera.**—*C. canella* and *C. mucorea* are not common in the southeast; I have seen specimens of the latter taken at Yuma, in March, by Mr. Daggett; *aneipennis* is not rare at Pomona, Pasadena, etc.

**Babia.**—*B. 4-guttata*, var., Needles (Wickham).

**Saxinis.**—*S. saucia* is common nearly everywhere west of the mountains, occurring most plentifully on flowers in May: *politula*, Santa Barbara (Horn): *speculifera*, Owens Valley (Horn), San Bernardino Mountains, July and August; not common: *hornii*, one example from San Diego in my collection is so referred; I have seen others from the same locality.



**Chlamys.**—A specimen taken at Palm Springs in April, it has not been possible to place with certainty; it is very probably undescribed.

**Exema.**—*E. conspersa* is common in most localities; taken by sweeping.

**Cryptocephalus.**—*C. sanguinicollis* is rather common from March to May; Pomona, Pasadena, San Diego, etc.; the variety *nigerrimus* has not yet been seen from our region, but it occurs near San Francisco, though rarely: *castaneus* is not rare at Pomona during May and June: *spureus* is rather rare; I have seen specimens from Pomona in July, and San Diego in August.

**Pachybrachys.**—*P. analis*, rare; Arrowhead Springs and San Diego, late in May: *pubescens* (of which *circumcinctus* and *viduatus* are synonyms) is quite common on willows during March and April: *hybridus*, moderately common; Pomona, Pasadena, San Bernardino, San Diego, etc.; May and June; taken by sweeping weeds and bushes near streams and by roadsides: *celatus*, San Diego, Yuma, Pasadena, Pomona; June: *livens*, Yuma, Pomona, San Bernardino Mountains; May to October; on *Salix*: *lustrans* is said to inhabit Southern California: *melanostictus* and *signatifrons* are from the State, without definite locality: *donneri*, recorded from Utah and Lower California; it is, therefore, probable that it is to be found in our district; I have seen specimens from Yreka in this State.

There are many undescribed species in my collection, from various parts of the State, but most of them from the south.

**Diachus.**—*D. auratus* is everywhere plentiful: *erosus*, not rare about San Francisco (Marin County).

**Adoxus.**—*A. obscurus*, "Cal. and Nev." (Horn); Lake Tahoe (Fényes).

**Myochrous.**—*M. longulus*, from Long Beach and Bakersfield, May; rather common along the Colorado River in July.

**Glyptoscelis.**—*G. illustris*, rare on pines in the San Bernardino Mountains, July; taken in moderate numbers by Van Dyke in Placer County, also on pines: *squamulata*, not rare at Riverside and Long Beach; April and May; occurs also in the San Joaquin Valley and in Arizona: *alternata*, Owens Valley and San Diego (fide Fuchs): *albida*, specimens have been seen from middle California.

**Colaspidea.**—*C. subvittata* is common on Santa Catalina and San Clemente islands during June and July; this species exhibits but little variation, while the three following are exceedingly variable in color, size and form. *C. cuprascens* is thus far known only from San Diego, where it is common in May: *varicolor* is abundant along the western base of the Sierras, near Pomona, Pasadena, etc.: *smaragdula* is found at higher altitudes, being abundant on pines in the San Bernardino Mountains in July, at an altitude of from 4,000 to 5,000 feet.

**Typophorus.**—*T. viridicyaneus*, "So. Cal." (Horn): *canellus*, Yuma.

**Metachroma.**—*M. californica* is occasionally not rare at San Diego and Long Beach in Spring; found also near Yuma.

**Chrysochus.**—*C. cobaltinus* is very common on milkweed, during May and June, nearly everywhere in Southern California.

**Colaspis.**—*C. oregonensis* is northern.

**Timarcha.**—*T. intricata* is northern. I have seen specimens collected by Van Dyke in Mendocino County.

**Leptinotarsa.**—*L. behrensi* and *L. 11-lineata* are quoted from "So. Cal." in Linell's recent table. I have not seen specimens, but they will be found only in the southeast, if really inhabitants of our region.

**Calligrapha.**—*C. elegans* var. *californica* is a rare species in Southern California; I have seen small series secured at Pomona and Pasadena, in June; Los Angeles (Van Dyke): *sigmoidea* and *serpentina* are more or less common in the middle and northern parts of the State, the former having occurred as far south as Santa Barbara (Fuchs).

**Chrysomela.**—*C. inornata*, "Cal."

**Plagiodera.**—One example of *prasinella*, San Bernardino Mountains; it is more common in Tulare County and about San Francisco.

**Gastroidea.**—*G. cyanea*, not rare at Pomona in May; probably found throughout the State; the variety *casia* occurs near San Francisco.

**Lina.**—*L. californica*, *L. lapponica*, and *L. scripta* are all common in various localities of middle and northern California.

**Gonioctena.**—I have a specimen of *pallida* bearing a "Cal." label. If the specimen is really from the State it must be from the north.

**Trirhabda.**—I have a specimen of *canadensis* from Yosemite Valley; it is probably confined to high altitudes or to the extreme north in this State: *geminata*, San Diego: *caduca*, Owens Valley, Arrowhead Springs, Pasadena, during May and June: *luteocincta* and *flavo-marginata* are found both in the larval and mature form

on *Artemisia californica*; the former is very abundant everywhere, the latter generally less common: *diducta* is found in Marin County, also (according to Horn) in western Nevada and adjacent regions: *attenuata* is also occasionally taken in the middle Sierras, Truckee (Fényes).

**Galerucella.**—*G. nymphæ*, Owens Valley (Horn): *notulata* has occurred in Alameda County: *tuberculata* is quite common in various localities in the middle and northern parts of the State.

**Monoxia.**—*M. puncticollis* is not common; found along the Southern California seacoast—Santa Monica and Santa Barbara—in August: *conspuata* and *sordida*, both quite variable, are plentiful and widely diffused: *debilis* is said to be from "Cal." and doubtless is found in the south.

**Diabrotica.**—*D. 12-punctata* is found only in the southeast, as is also its variety *tenella*; the latter has been taken at Yuma: *soror* and *trivittata* are everywhere found west of the mountains, the former being one of the commonest of our beetles.

**Phyllobrotica.**—*P. viridipennis* is not common; taken in the San Bernardino Mountains, in July, by sweeping herbage on a damp hillside: *luperina*, reported from Santa Barbara: *nigripes*, Los Angeles.

**Scelolyperus.**—*S. flavicollis*, "Tejon and various places in So. Cal." (Horn): *maculicollis*, San Diego, April: *graptoderoides*, Santa Barbara, Ventura, Los Angeles: *tejonicus*, described from Tejon: *loripes* occurs in Tulare County, *schwarzii* and *longulus* are found in the north, and *decipiens* at Yreka.

**Trachyscelida.**—*T. bicolor*, Yuma.

**Luperodes.**—*L. bivittatus* is moderately common at San Diego, less so at Pomona; May and June: *transitus*, Santa Barbara: *laticeps*, occasionally common at Pomona, April and May: *torquatus*, quite common at Pomona, San Diego and Los Angeles: *smaragdinus*, Pasadena and Pomona, April and May; not very common: *varipes*, rather rare in the San Bernardino Mountains, more common farther north.

**Androlyperus.**—*A. fulvus*, "Coast region south of San Francisco."

**Malocorhinus.**—*M. maculatus*, "So. Cal."

**Ædionychis.**—*Æ. violascens* is exceedingly rare; I have only once met with it, a series of about a dozen specimens having been found beneath stones along the margin of a small stream in the foot-hills near Pomona in November.

**Disonycha.**—*D. 5-vittata* is abundant everywhere on willows, near water; April to July: *maritima*, one example taken at Pomona in October. I have seen specimens of *pennsylvanica* from Lake County, and Mr. Fuchs writes me that it has been found near Los Angeles. .

**Haltica.**—*H. bimarginata* is widely diffused and common, at times being so abundant as to nearly strip the alders of their foliage: *carinata* is sometimes numerous, especially on grape-vines, and widely diffused: *californica*, not common at Pomona and Pasadena, June to October: *aruginosa*, Pasadena, August; Long Beach (Daggett): *obolina*, common along the western and southern base of the Sierras, in Los Angeles County, May to September: *punctipennis*, specimens so referred are quite common at Redondo in spring, on low plants above the beach. A single specimen of an apparently undescribed species has been taken in the San Bernardino Mountains.

**Hemiglyptus.**—A small series of *basalis* was obtained by sweeping, near Pasadena, in April.

**Crepidodera.**—*C. helxines* is found everywhere, on willows.

**Epitrix.**—*E. cucumeris* is widely diffused and usually common: *suberinita*, not rare, especially near the sea-coast; Santa Monica, February and March: *parvula*, common at Pomona on various weeds about orchards, etc.; April to October.

**Leptotrix.**—*L. reticulata* is found always in February; Pasadena, Santa Monica, San Diego.

**Chætocnema.**—*C. opacula*, Los Angeles, August; not common: *subviridis*, Owens Valley: *opulenta*, Owens Valley (Horn), San Bernardino Mountains; not rare, by sweeping, in a meadow at some 6,000 feet elevation: *ectypa* is our commonest species in Southern California—Los Angeles, Pomona, Pasadena, etc., April to November; also at Yuma (Wickham): *cribrata* occurs at Lake Tahoe, July: *irregularis*, San Jose and northward: *subcylindrica*, Lake Tahoe (Fényes): *cribrifrons*, "Cal." (fide Ulke): *denticulata*, "Cal." (fide Horn): *confinis*, middle and northern parts of the State.

**Systema.**—*S. taniata* is abundant nearly everywhere, and very variable; it may be taken from October to June, by sweeping weeds in gardens and by roadsides: *subaenea*, "Cal. and Nev."

**Longitarsus.**—*L. repandus*, Pomona, Pasadena, San Diego; abundant at Pomona on *Senecio californica*: *livens* is also common, though somewhat less so than the preceding; found throughout maritime Southern California: *montivagus*, foot-hills near Pomona, May to October; not common: *mancus*, a few specimens taken



by Dr. Fényes at Pasadena, February: *rufescens* occurs in the northern coast region—Mendocino County. An undescribed species was taken at Yuma, Colorado River, in July, and at Pasadena.

**Glyptina.**—*G. cerina* and *G. atriventris* are more or less plentiful in most localities west of the mountains; taken by sweeping.

**Phyllotreta.**—*P. lepidula*, Los Angeles (Fuchs): *vittata*, specimens so referred have been taken on two occasions at Pomona: *albionica*, San Bernardino Mountains, August; Pomona, October and January; not rare, though less frequent than *pusilla*: *ramosa*, rare, Pasadena, June: *pusilla*, common and widespread: *oregonensis*, *denticornis* and *lewisii* inhabit the mountainous regions of the middle and northern parts of the State. One example of an undescribed species has been taken in the San Bernardino Mountains.

**Dibolia.**—Specimens referred to *ovata* have been found by Mr. Daggett, in Bear Valley, in June.

**Psylliodes.**—*P. punctulata*, rare, Pasadena, June: *conversior*, rare, Pomona, November.

**Microrhopala.**—*M. rubrolineata* is common everywhere in our district, at least west of the mountains, on *Heterotheca grandiflora*: *melsheimeri* has been collected in some numbers, by beating *Ceanothus* (?); San Bernardino Mountains, June to August.

**Odontota.**—*O. californica* is abundant on *Ceanothus*; San Bernardino Mountains, in midsummer; the beetles have been bred by Coquillett from larvæ mining the leaves of *Ceanothus integerrimus*: *rubra*, specimens, doubtless from the north, have been seen.

**Stenopodius.**—*S. flavidus*, "California (San Diego) and Arizona."

**Cassida.**—Two examples of *texana* were taken at Pomona, May 5. Coquillett records the finding of both larvæ and beetles on *Solanum xanti*, May 12.

**Coptocycla.**—*C. aurichalcea*, Pasadena, August.

### BRUCHIDÆ.

**Bruchus.**—*B. pisorum* is not often seen; the writer has taken a few specimens under bark of eucalyptus at Pomona, in winter: *sordidus*, from desert regions of the east or southeast: *ramicornis*, Owens Valley: *limbatus*, Yuma, March (Daggett): *pruininus*, taken sparingly at Pasadena, March (Fényes): *aureolus*, desert regions "on flowers of *Astragalus*;" *pauperculus*, common, taken by sweeping flowers; Pomona, Pasadena, Ojai Valley, etc, March to May: *prosopis*, breeds abundantly in the pods of mesquite (*Prosopis juliflora*), in the desert regions: *protractus*, occurs in the same region as the preceding, also on mesquite: *uniformis*, "pods of *Prosopis* and *Strombocarpus*, desert of Colorado River;" *obtectus*, one example taken in a Los Angeles hotel; it is likely to prove more common than is desirable: *exiguus*, Sierra Madre Mountains (Daggett); I have also seen specimens from the neighborhood of San Francisco: *seminulum*, recorded from California by Horn, but I have not seen specimens. A small species closely resembling *seminulum* but apparently distinct, is very plentiful throughout Southern California, and is probably the species regarded as *seminulum* by Horn. Several undescribed species have been taken by various collectors, there being three or four in my own collection.

**Zabrotes.**—*Z. spectabilis* is especially found on flowers of greasewood, though not common; San Bernardino

Mountains, July; Pomona, September; Pasadena, June: *densus* is credited by its describer to "California." A single example of an undescribed species was taken in the San Bernardino Mountains.

### TENEBRIONIDÆ.

**Craniotus.**—According to Mr. Fuchs, a specimen of *pubescens* was secured by Dr. Blaisdell from near Vallecito, San Diego County.

**Edrotes.**—*E. ventricosus* is not uncommon on the Colorado Desert; I have taken it under stones at Palm Springs, on the western border of the desert, in April. With typical specimens of *ventricosus* occurred others of duller surface lustre, with coarser punctuation and pubescence less evidently condensed in lines along the elytra; these Dr. Horn considered as mere varieties of *ventricosus*, and if his opinion is well founded probably *nitidus*, described from the Mojave Desert, should be similarly disposed of.

**Triorophus.**—*T. punctatus* was described from a single dead specimen found at Vallecito: *lævis* is common throughout the Colorado Desert: *subpubescens*, Antelope Valley, Los Angeles County.

**Stibia.**—*S. ovipennis* and *S. maritima* both occur only near San Diego, so far as is known. According to Dr. Blaisdell, the former "inhabits the seashore and is found beneath the beach-berry (*Mesembryanthemum æquilaterale*) growing upon the sand-dunes," while *maritima* inhabits the Coronado peninsula, and is not confined to the seashore.

**Auchmobius.**—*A. sublævis* is widely diffused in California, but everywhere very rare; I have taken a single

specimen from under a stone in San Antonio Cañon, near Pomona. Horn gives as localities Owens Valley, Tejon, and the southern Coast Range.

**Eurymetopon.**—*E. rufipes* is plentiful in the eastern or desert portions of Southern California: *fusculum*, Yuma, March (Daggett): *convexicolle*, very abundant throughout the region to the west of the mountains, occurring under bark, stones, and rubbish: *sodalis*, Owens Valley, Yuma: *serratum*, Palm Springs, April: *inflatum*, at times obtainable in some numbers beneath the vegetation growing on the sand-dunes along the shore from Santa Monica to San Diego: *bicolor* and *cylindricum* have been taken in Kern County; of the former I have specimens taken at or near Bakersfield in May.

**Emmenastus.**—*E. longulus* is rather common, though less plentiful than *Eurymetopon convexicolle* with which it occurs (see above): *piceus*, San Diego and San Bernardino counties; Blaisdell records typical specimens from Julian, at an altitude of 6,000 feet: *obesus*, San Diego; "plentiful under dry cow manure, boards, rags, etc." (Blaisdell): *thoracicus*, San Geronimo Pass (Casey): *obtusus* and *ater* belong to middle California; I have seen numerous specimens of the latter from San Francisco, and Casey mentions Napa and Sonoma counties for the former: *crassicornis* is described from Humboldt County.

**Epitragus.**—*E. pruinosus*, Owens Valley (Horn); Pomona, two examples obtained by beating late in July, uncommon; San Diego, "taken in net from *Rhus laurina* and also beneath bark at the base of rotten trees."

**Chilometopon.**—*C. abnorme*, Palm Springs: *helopioides*, recorded by Horn from "Cal." simply, but without doubt from the desert country to the southeast.

**Cnemodus.**—*C. testaceus*, Yuma; apparently scarce.

**Zopherus.**—*Z. tristis*, San Bernardino County: *granicollis*, San Bernardino Mountains, July; Pomona, May, and Echo Mountain: *induratus*, Julian, San Diego County. None of the species are at all common.

**Phlæodes.**—*P. diabolicus* is common and widely dispersed in our district, occurring under bark and in decaying stumps, especially of the live-oak. According to Blaisdell, the adult insect feeds upon a large and tough species of fungus growing upon the oak.

**Noserus.**—*N. plicatus* is found under the bark of dead trees, at Pomona and at Julian, San Diego County (Blaisdell); scarce.

**Phellopsis.**—*P. obcordata* var. *porcata*, mountain regions from Lake Tahoe to Shasta.

**Usechus.**—*U. lucerta*, middle and northern regions.

**Dacoderus.**—*D. striaticeps*, Vallecito, San Diego County.

**Aræoschizus.**—*A. sulcicollis*, Owens Valley: *costipennis*, Vallecito (Le Conte); one example under bark, Indio, December; Palm Springs, April (Fényes): *armatus*, Owens Valley.

**Anepsius.**—*A. delicatulus*, Pomona and Palm Springs; found under stones, in dry places; rare west of the mountains, more common on the desert; according to Blaisdell the species is not uncommon in the maritime portions of San Diego County.

**Typhlusechus.**—*T. singularis* is a minute, blind species of which two specimens were found by Koebele in Los Angeles County.

**Nyctoporis.**—*N. carinata* is widely distributed in our district, occurring under bark, logs and debris: *galeata*, *cristata* and *aquicollis* occur in the State, but farther north.

**Cryptoglossa.**—*C. verrucosa* is not rare in the Mojave and Colorado deserts; I have seen specimens from Antelope Valley and from Needles, also from other points farther south, along the Colorado River: *laevis* occurs at Yuma, "under logs and stones" (Horn).

**Centrioptera.**—*C. muricata* is not rare at Palm Springs, in April. The beetles remain concealed beneath stones and rubbish during the heat of the day, appearing about sundown, when numbers may be seen walking slowly about the dry washes and amid the sagebrush, a custom common to many of the larger Tenebrionidæ of the southwest. *C. seriata* is found in "So. Cal." according to Horn, and Blaisdell mentions *asperata* as occurring rarely at Poway, San Diego County.

**Schizillus.**—*S. laticeps*, Mojave Desert (Crotch). I do not know of any specimens having been taken by our later collectors.

**Microschatia.**—*M. inaequalis*, San Diego and Riverside in February and March; not rare. Specimens of what appears to be an undescribed species have been taken by Mr. George H. Field in San Diego County.

**Asida.**—*A. ægrota* is recorded by Blaisdell from the desert parts of San Diego County; it is otherwise only reported from Lower California: *actuosa*, Owens Valley, in winter (Horn); Antelope Valley, Los Angeles County, in early spring: *carinata*, San Diego County, Colorado Desert: *confluens*, *parallela*, *hirsuta* and *hispidula* are all from the Colorado Desert: *impestrata*, San



Diego and Yuma: *obsoleta*, "Borders of Col. Desert;" San Bernardino Mountains, July; Riverside, Julian, Pasadena, October: *muricatula*, San Diego and Los Angeles: *luctata*, rare in Owens Valley: *angulata*, San Diego and Riverside, winter and early spring: *lecontei*, Kern and Tulare counties: *captiosa*, Tulare region and Monterey County: *marginata* and *gabbii*, San Diego County (fide Fuchs).

**Coniontis.**—The species of this genus occur under stones and rubbish, usually in dry places, though this is not always the case. They are difficult to discriminate, and it is doubtful if all the species named below are well founded; if so, then there are surely several others awaiting description. *C. abdominalis* is our largest species, and one of the rarest; it has been taken at Santa Barbara, Los Angeles, and Pasadena: *robusta* occurs at Santa Barbara: *elliptica*, specimens so identified by Casey are common at Riverside, Pomona, San Diego, Catalina Island, etc.; it is difficult to separate from *robusta*, and was not considered distinct by Horn: *lata*, San Clemente and Santa Barbara islands; var. *insularis*, Santa Cruz and Santa Rosa islands: *opaca*, Owens Valley: *punctipes*, San Bernardino County: *subpubescens*, not rare at various localities from Los Angeles to San Diego: *globulina*, Tehachapi Pass: *pallidicornis*, "So. Cal.:" *parviceps*, San Diego, Pomona, Pasadena. Numerous other species are found farther north: *inaequalis*, "Cal.," without definite locality: *elongata*, *genitiva*, and *parallela* are from the northern part of the State: *viatica*, *eschschooltzii* and *nemoralis* are from San Francisco: *puncticollis*, from San Francisco and the central valley: *farallonica*, from the Farallon Islands: *montana*, from Lake Tahoe.

**Cœlomorpha.**—*C. maritima* was described from Lower California, but Horn also cites San Diego; it is similar to *Cœlus* in habits.

**Cœlus.**—The individuals of this genus are very abundant at all seasons in the sand-dunes along the coast from San Francisco to San Diego; they vary exceedingly in size, form and thoracic sculpture, and I am unable to recognize more than two species along our southern coast, viz., *globosus* and *arenarius*, the former having a deeply, and the latter a feebly, sinuate epistoma. Typical *grossus* is very large, with unevenly punctuate thorax; but there is every gradation between this and typical *globosus*, in fact, the majority of specimens are intermediate. Both *latus* and *curtulus* are founded chiefly on form and punctuation of the prothorax, both of which appear to me to be too unstable to be of use in specific division, and I have little hesitation in placing the former as a synonym of *arenarius* and the latter as a synonym of *ciliatus*. Two other species are found on the Southern California islands: *pacificus*, on Santa Barbara and Santa Rosa, and *remotus*, on San Clemente; it is doubtful if these two species share the burrowing habit of their congeners on the mainland.

**Eusattus.**—*E. robustus* has thus far occurred only on San Clemente Island: *politus*, Santa Barbara and Santa Rosa islands; apparently not common: *coquilletti*, Los Angeles County (Coquillett): *lavis*, Southern California (teste Horn)\*: *dubius*, Colorado and Mojave deserts, Antelope Valley: *productus*, "Ari. and Lower Cal." (Horn); I have an example labeled "Cal.," doubtless from the southeastern region: *convexus*, taken at

---

\* "The Coleoptera of Baja California." Proc. Cal. Acad. Sci., 2nd Ser. Vol. IV, 1894, p. 349

Vallecito, San Diego County (Le Conte): *difficilis*, not rare at San Diego, where I have taken it in some numbers about the roots of plants on the Coronado peninsula: *muricatus*, Colorado Desert and Owens Valley. Casey has described *nanus* from Kern County.

**Eleodes.**—A goodly number of representatives of this common western genus are found in Southern California. Their habits are practically the same as those of the greater part of the larger wingless Tenebrionidæ. While they are occasionally abroad by day, as a rule they remain concealed beneath boards, stones, rubbish, etc., till toward evening, when they become active. *E. quadricollis* is a common species about San Diego, and is found near the seacoast as far north as San Francisco; it is not uncommon on Catalina Island: *armata* is probably the most abundant species of the genus along the line of the Southern Pacific Railroad, through the Colorado Desert; I have found it common at Palm Springs, in April, on the western border of this desert: *femorata* is decidedly rare in the vicinity of San Diego, the only locality from which it is known: *gentilis* is said by Blaisdell to be common at San Diego, but specimens coming to me from that place and bearing this name proved to be only *quadricollis*; the type, however, comes from that vicinity, and Blaisdell's reference is perhaps correct: *interrupta* is rare about San Diego (Blaisdell): *gracilis* is represented by one example taken by me at Palm Springs, and so identified by Dr. Horn; it is not typical and may possibly prove distinct: *grandicollis* and *gigantea* are found at various points from Los Angeles to San Diego, on the western side of the mountains, and both are said to be more or less common about San Francisco; of the two, *grandicollis* is the scarcer in Southern California: *acuticauda* is our most

abundant species throughout the southern part of the State; at Pomona, Pasadena and vicinity, it finds shelter in cellars, and is, in fact, the only species of the genus that occurs commonly about houses: *pilosa* and *hirsuta* inhabit Owens Valley: *scabripennis* is known from Santa Barbara and Santa Rosa islands and Fort Tejon; it does not appear to be common: *consobrina* is not rare in the foot-hills and up to altitudes of 5,000 to 6,000 feet in the southern Sierras: *tenebrosa* occurs in Owens Valley: *parvicollis* and its varieties, *producta* and *marginata*, are given on the authority of Dr. Blaisdell, who names them in a list of species observed by him in San Diego County; both occur about San Francisco, and I have never seen specimens from further south: *subnitens* has been taken in San Diego County (fide Fuchs). The following species occur in those parts of the State to the north of our district: *E. obscura* var. *sulcipennis*, desert region in the northeast; *humeralis* and *elegans*, northern coast region; *estriatus*, *clavicornis* and *parvicollis*, about San Francisco, the second named species being very abundant in the sand-dunes on the ocean front; *dentipes*, common to the south of San Francisco; *prominens*, San Luis Obispo County; *granulata*, *granosa* and *cuneaticollis*, exact locality not named, though it is probable that the first two occupy the desert region of the northeast and possibly come as far south as Owens Valley; *cordata*, middle region, from San Francisco to Lake Tahoe, also ranging both north and south of this latitude in the central valley; *scabricula*, Sacramento and Lake Tahoe.

**Embaphion.**—*E. depressum*, Vallecito (Le Conte).

**Trogloderus.**—*T. costatus*, San Diego County (Fuchs).

**Eulabis.**—*E. grossa* is not rare on San Clemente Island, and has also been taken on Santa Barbara and

San Nicolas islands: *rufipes*, *pubescens* and *laticornis* are more or less common from San Diego north through the western portions of our district, especially under stones and rubbish in dry places in spring: *crassicornis* is described from "So. Cal.," and must resemble the preceding in habits: *obscura* is plentiful along the sea beaches: *bicarinata* occurs in the San Joaquin Valley.

**Cerenopus.**—*C. concolor* and *C. costulatus* are taken in the Colorado Desert; the former is moderately common at Palm Springs, the latter is said by Blaisdell to be rare in San Diego County.

**Argoporis.**—*A. costipennis* and *A. bicolor*, Colorado Desert: *inconstans*, San Diego (Horn).

**Amphidora.**—*A. littoralis* is common everywhere west of the mountains; found beneath logs, rocks, bark, etc.; as remarked by Blaisdell, the name is not at all appropriate: *nigropilosa* occurs along the seashore and may usually be found in numbers about the roots of plants on and adjacent to the sand-dunes, or concealed beneath any object that will offer shelter.

**Cratidus.**—*C. osculans* is common and widely diffused: *fuscipilosus* occurs, according to Blaisdell, on the summits of the mountains about Julian; it is, perhaps, not specifically distinct from the preceding.

**Stenotrichus.**—*S. rufipes*, San Diego, Riverside, Pomona, Santa Monica; not very common.

**Alobates.**—*A. pennsylvanicus* is northern.

**Ipthimus.**—*I. serratus* and its variety, *sublaevis*, occur under bark, in the Sierras; the latter has been taken in the San Bernardino Mountains.

**Cælocnemis.**—*C. dilaticollis* is found in the higher parts of the southern Sierras, under bark, logs, etc.:

*obesa* is not rare in the foot-hills near Pomona, Pasadena, etc.: *magna*, Central Valley: *rugosa*, Los Angeles County (Coquillett).

**Centronopus.**—*C. parallelus*, middle Sierras, under bark.

**Cibdelis.**—*C. blaschkii*, San Diego County, "mountainous districts, under bark" (Blaisdell); commoner farther north: *bachei*, Santa Barbara, San Clemente and Santa Catalina islands: *laevigata*, Santa Barbara.

**Tenebrio.**—Specimens of *obscurus* have been found in grain at Pasadena; perhaps a mere sporadic importation: *tenebrioides* is rare, under bark, in the San Bernardino Mountains.

**Bius.**—One example of *estriatus* was found under pine bark in the San Bernardino Mountains, July; it is one of our rarest species.

**Doliema.**—*D. plana* is generally rare, but is occasionally found in large numbers under the bark of dead sycamores; I have taken it thus in the Ojai Valley, in March, and Blaisdell reports similar captures in San Diego County; occurs also at Yuma (Wickham).

**Alæphus.**—*A. pallidus*, Palm Springs (Fényes), Fort Tejon (Horn). Mr. Wickham reports a species of *Alæphus* "near *pallidus*" from Needles, on the Colorado River.

**Eupsophus.**—*E. castaneus*, Palm Springs, not rare in spring; Owens Valley, flying by night (Horn).

**Mecysmus.**—*M. angustus* is found rather abundantly flying at night at Yuma (Horn), Mojave Desert, Pomona and Riverside; under stones and on sidewalks at the latter places (never on the wing in my experience): *tenuis* is described from "So. Cal."



**Trichoton.**—*T. sordidum* occurs at Palm Springs, in April, under boards, stones, etc., as is generally true of the *Blapstini*.

**Ulus.**—*U. crassus* is common in many localities: *latus*, along the San Diego River (Blaisdell).

**Blapstinus.**—*B. longulus*, Yuma, San Diego (Blaisdell): *validus*, "So. Cal." (Casey); probably from the Colorado Desert: *dilatatus*, common everywhere: *histricus*, Los Angeles County (Newhall): *coronadensis*, rather common at Coronado (Blaisdell): *rufipes*, our commonest species: *aqualis*, San Bernardino: *funnebris*, Southern California (Casey): *brevicollis*, Riverside, Santa Monica, Pomona; not very abundant: *pubescens*, San Bernardino, Pomona, San Diego: *sulcatus*, Riverside, San Diego, Ojai Valley; rather common. The following species are described from the middle and northern parts of the State; their identification is a matter of much difficulty: *fuliginosus*, *discolor*, *gregalis*, *pulverulentus*, *parallelus*, and *inquisitus*.

**Conibius.**—*C. parallelus* is said by Blaisdell to be rather common about San Diego and Poway, but I have not yet detected it in the vicinity of Los Angeles; it was described from San Jose: *seriatus* is common in Los Angeles County, and occurs as far north as Sylvania; it was described from the Colorado Desert. The distinctness of this and the preceding species I believe may fairly be questioned. *C. elongatus* is said by Horn to be rather common under stones, in Owens Valley: *crassipes* is described by Casey from "So. Cal."

**Notibius.**—*N. puberulus*, Colorado Desert, Needles (Wickham), not rare at Palm Springs in April: *puncticollis*, rather frequent at Riverside, Pomona, and Pasadena; often seen on city streets and walks: *granulatus*,

Palm Springs, Yuma: *gracilis*, Indio (Casey): *sulcatus*, Riverside, San Diego, San Clemente Island; not common.

**Cnemeplatia.**—*C. sericea* is generally rare; it was once taken in some numbers, under boards, amid stable refuse at Pomona; it is occasionally taken on the wing toward evening, and at San Francisco comes to electric lights, at times in great numbers. Horn speaks of the species as being rare in Owens Valley, under stones.

**Alaudes.**—*A. singularis* is a very rare species originally taken by Dr. Horn in Owens Valley, where it was found living under stones, with a small black ant. A few specimens were next taken by Wickham—some twenty-five years later—in eastern Oregon, then a single example by Koebele, in the Argus Mountains of southeastern California, not far south of the original habitat. In November and December of 1893 and 1895, I secured about a dozen examples from under boards in my garden at Pomona, under conditions which apparently indicated no connection with ants; and since then Dr. Fényes has taken about an equal number at Azusa, Los Angeles County, this time with ants. I know of no other instances of its capture.

**Tribolium.**—*T. ferrugineum* and *T. confusum* are both found in ground cereals, the former being less common in my experience. Blaisdell speaks of *ferrugineum* as being rare, under the bark of trees, at San Diego.

**Phthora.**—*P. americana* occurs at Lake Tahoe.

**Gnathocerus.**—*G. cornutus* occurs with *Tribolium* in ground cereals.

**Echocerus.**—Several specimens of *maxillosus* have been taken at Pasadena in March.

**Ulosonia.**—*U. marginata* occurs along the Colorado River, under cottonwood bark.

**Merotemnus.**—*M. elongatus* labeled "Cal." is probably from the southeast.

**Aphanotus.**—*A. brevicornis* is rather common in the eastern portion of San Diego County (Blaisdell); very rare at Pomona, a single dead specimen being all that I secured in seven years' collecting.

**Alphitobius.**—*A. ovatus* and *A. piceus* have both found their way into our territory, though neither is as yet common. Of the former species, I secured a specimen under bark at Indio, in the Colorado Desert, and several examples of the latter have been taken at Pasadena by Dr. Fényes.

**Cynæus.**—*C. angustus* is reported from the Colorado Desert: *depressus* is rather common in and upon the decaying base and roots of *Yucca whipplei* in various localities; I have found it near Pomona in May.

**Metaclisa.**—*M. marginalis* is rather rare, occurring under bark of pines in the San Bernardino Mountains; also found at Julian (San Diego County).

**Uloma.**—*U. longula* is not rare under bark, in the Sierras.

**Phaleria.**—*P. rotundata* is very abundant everywhere on our seacoast from San Diego to San Francisco, occurring beneath kelp. I have no hesitation in uniting *limbalis* with the preceding; a long series shows every grade in coloration from uniform testaceous to typical *limbalis*, which is in great part black; a precisely analogous variation is observable in the common *testacea* of the Atlantic coast. *P. debilis* is reported from Yuma:

*globosa* is common on the seashore, near San Francisco: *humeralis* is said to be from "Cal."

**Anemia.**—*A. californica*, Owens Valley.

**Platydemia.**—*P. subquadratum*, Colorado River. Mr. Wickham reports it as not rare in August at East Bridge, just across the river from Needles. *P. oregonensis* is rather common about fungus on old stumps, under bark, etc., in the Sierras from San Diego County north.

**Alphitophagus.**—*A. bifasciatus* is not especially uncommon of late at Pasadena and in the neighboring mountains.

**Hypophlæus.**—*H. substriatus* occurs in the Sierras, under bark: *opaculus* is very scarce; two examples were taken in March from the burrows of *Monarthrum scutellare*, in live-oak.

**Pentaphyllus.**—*P. californicus* is northern.

**Eleates.**—*E. explanatus* and *E. occidentalis* both occur on fungus growing on logs in the region about Lake Tahoe; the former also inhabits the forests to the northward of San Francisco.

**Megeleates.**—*M. sequoiarum*, middle Sierras—Lake Tahoe and Calaveras. The larva has been taken by Dr. Blaisdell in a species of woody fungus, and has been described by Wickham.

**Apocrypha.**—*A. anthicoides*, Pomona, Pasadena, San Diego; uncommon. All specimens seen have been taken in early spring. Dr. Fényes once found it associated with ants, but perhaps accidentally, as this habit has not been elsewhere noticed. *A. dyschirioides* is said by Blaisdell to occur rarely near San Diego: *clivinoides* is found in Owens Valley.

**Helops.**—*H. edwardsii* is not rare, under bark, in the San Bernardino Mountains: *ovipennis*, Mojave Desert: *rugicollis*, Owens Valley and Tejon: *strigicollis*, "So. Cal.," probably near San Diego (Horn): *attenuatus*, Vallecito (Colorado Desert) and Owens Valley. *H. bachei* and *H. blaisdelli* are said by Blaisdell to occur under maritime plants growing on the sand-dunes near San Diego, and one of these or a closely allied species I have found very common in similar situations at Redondo Beach; *bachei* was described from the Santa Barbara Islands, and typical examples are perhaps found only on the islands: *discipula* is described from the vicinity of San Diego.

The following species are more northern: *opacus*, San Francisco and north, also Sacramento; *punctatus*, "Cal.;" *rugulosus*, San Francisco; *angustus*, Tejon; *californicus*, San Jose, Lake Tahoe (beaten from pines in July); *tumescens*, "Cal., Coll. of Ulke."

#### ÆGIALITIDÆ.

**Ægialites.**—The exact locality of *californicus* is not stated; it is doubtless northern, if really from the State: *fuchsii* has been obtained in some numbers from Mendocino County and Farallon Islands, by Mr. Fuchs.

#### CISTELIDÆ.

**Xystropus.**—Horn expresses a belief, which I think well founded, that *X. californicus* is an accidental importation from Mexico or further south. The original specimen was found at Martinez, on the shore of San Francisco Bay, and has never since been duplicated.

**Stenochidus.**—I have taken specimens that are referable to *gracilis* and *cyanescens*, at Pomona and Pasadena, during May and June, but have not yet succeeded

in separating them to my satisfaction. It is certain that the color of the legs is not a specific character, these being usually red in the males and black in the females.

**Hymenorus.**—*H. infuscatus*, Los Angeles, Pomona, June; Catalina Island, July: *grandicollis*, specimens occurring at light on Echo Mountain were so identified by Casey; a different species in the collection of Dr. Fényes, and taken at Palm Springs, fits the description of *grandicollis* still better: *fusicornis*, Riverside and Pasadena, April and May; rare: *fuscus*, Coronado, in decaying sunflower blossoms (Blaisdell): *macer*, Poway, common, (Blaisdell): *punctatissimus*, Riverside, Pomona; June to October; not rare. Blaisdell reports *inquilinus* as occurring in the nests of the agricultural ant, in Calaveras County, in September. *H. punctulatus*, *H. discrepans*, and *H. uniseriatus* are Californian, but without definite locality.

**Mycetochares.**—*M. longipennis* is rare at Pasadena, March (Fényes): *procera*, Los Angeles (Casey); Pasadena, April (Fényes): *pubipennis*, Pasadena and Redondo, March. All the above are more or less rare. *M. nevadensis* I have beaten from pines at Lake Tahoe, in July: *crassulipes* and *pacifica* are also described by Casey from California, the former from Humboldt County, the latter without precise locality.

**Isomira.**—*I. variabilis* is exceedingly common on flowers, especially of greasewood, in the foot-hills, and up to moderate elevations in the Sierras; it occurs also on Catalina Island and is probably found throughout the region to the west of the mountains. *I. luscitiosa* is also quite common in the Sierras, in company with the preceding: *monticola* is described from Lake Tahoe,



and *discolor* from "Cal." Hopping alludes to the latter as being common at from 1,000 to 7,000 feet elevation along the Kaweah River, in Tulare County.

**Cistela.**—*C. opaca* is common on flowers in the southern Sierras from April to June. The color of the legs is not a specific character, the femora being about as often red as black. The specimens taken by Blaisdell in San Diego County and referred to *theveneti*, are doubtless the red-legged form of *opaca*, *theveneti* occurring in the middle Sierras, so far as known.

#### OTHNIIDÆ.

**Othnius.**—*O. longicornis*, Yuma.

#### MONOMMIDÆ.

**Hyporhagus.**—*H. gilensis*, San Bernardino County (Fuchs).

#### MELANDRYIDÆ.

**Carebara.**—*C. longula* is rather plentiful on pines, in the San Bernardino Mountains, in July.

**Phlæotrya.**—*P. riversi*, *P. bicincta*, and *P. vaudoueri* are all northern, the two former being from Sonoma County.

**Eustrophus.**—*E. repandus* is from the extreme north.

**Nothus.**—*N. luteus*, Pomona and Pasadena in June; taken on oaks.

**Lacconotus.**—*L. pinicolus*, Pomona and Pasadena; rare during May and June; found always on oaks, notwithstanding its name.

**Mycterus.**—*M. concolor* is not rare in the San Bernardino and other ranges of the southern Sierras, in June and July; the variety *flavipennis* occurs rarely with the type: *quadricollis* is said by Crotch to occur on flowers of Agave, at Temescal: *canescens* is found also in the Sierras, but farther north.

#### PYTHIDÆ.

**Priognathus.**—*P. monilicornis*, Tallac (Lake Tahoe), under bark (Fényes).

**Cononotus.**—*C. macer*, Owens Valley: *punctatus*, San Jose: *sericans*, San Jose and San Diego.

**Salpingus.**—*S. alternatus* is northern.

**Rhinosimus.**—*R. ceneirostris*, about San Francisco.

#### CEDEMERIDÆ.

**Calopus.**—*C. angustus*, "Cal.," probably northern.

**Ditylus.**—*D. quadricollis*, from the northern and middle Sierras.

**Nacerdes.**—*N. melanurus*, Long Beach; a single dead specimen washed up by the waves. I have not thought it wise to place this species in the foregoing list on so slender a basis.

**Xanthochroa.**—*X. californica*, middle Sierras: *centralis*, Sylvania (Ricksecker): *marina*, Marin County.

**Copidita.**—*C. quadrimaculata* is rather common, under rubbish along the sea beaches at Santa Monica and San Francisco: *bicolor* is northern: *cyanipennis*, "Coast Range below San Francisco."

**Asclera.**—*A. nigra*, San Bernardino County: *excavata*, foot-hills near Pomona, April and May: *discolor* is northern.

**Chrysanthia.**—*C. repanda*, Yuma.

**Oxacis.**—*O. fragilis*, San Diego: *bicolor*, not rare on the wing at dusk in the San Bernardino Mountains, altitude 5,000 feet: *O. sericea*, Owens Valley: *lucana* and *debilis* are from Yuma.

**Rhinoplatia.**—*R. ruficollis*, Owens Valley.

### CEPHALOIDÆ.

**Cephaloon.**—*C. tenuicorne*, northern: *bicolor*, Sonoma and Placer counties.

**Drachylis.**—*D. simulans* was recently described by Casey from Placer County.

### MORDELLIDÆ.

**Pentaria.**—*P. nubila* is abundant everywhere: *hirsuta*, Southern California: *pusio*, common and widely dispersed, frequently beaten from oaks, but nearly always obtainable by promiscuous beating and sweeping; this species has been wrongly included with *Anaspis*.

**Anaspis.**—*A. atra* and *A. collaris* are both frequent on flowers, though the former is more plentiful farther north: *militaris* and *sericea* are quoted from "Cal.," but I am unable to ascertain precise localities. Two specimens of an undescribed species were taken by me in the San Bernardino Mountains, in July.

**Mordella.**—*M. scutellaris* is very common and widely dispersed; it is extremely variable in size. A single specimen of an undescribed species was secured at Pomona some years ago.

**Mordellistena.**—The species of this genus are rather numerous in our district and many are undescribed. The greater number are black without ornamentation and will prove difficult to define. The following are

now credited to the State: *vilis*, San Diego, Los Angeles, etc., common; *comata*, San Diego; *aspersa*, San Diego (teste Smith); *tosta*, Yuma (Horn); *nubila*, San Diego (Le Conte), Riverside, one example identified by Smith; *unicolor*, Pomona, not common; *infima* and *æqualis* also inhabit the State according to Smith.

#### ANTHICIDÆ.

In a recent revision of this family Casey has established a large number of new genera, especially for the reception of the species now included under *Anthicus* and *Xylophilus*. It may be that some of these will stand but the writer is convinced that many of them will not, and until proper discrimination is possible, it has seemed best to employ the genera recognized by Le Conte and Horn in the "Classification."

**Eurygenius.**—*E. constrictus* is not very common; it is taken by beating and sweeping, in the foot-hills near Pomona and Pasadena, during May and June.

**Stereopalpus.**—*S. pruinus* is moderately common on willows, near Pomona, in June: *nimius* and *incanus* are described from "So. Cal." and Los Angeles County, respectively: *variipes*, *indutus*, and *impressicollis* are described from the northern part of the State.

**Corphyra.**—*C. distinguenda* is not rare, Pomona during April and May: *bardii*, Ventura, Santa Barbara, Pasadena; not common: *inconspicua*, Riverside, Pomona, Pasadena; not rare. The males of all these species are much rarer than the females, and as the latter sex are mutually nearly indistinguishable in many cases, it is quite likely that we shall find other species are inhabitants of our territory when the males are known. The following species are also Californian:

*flabellata*, Lake Tahoe and north; *crotchii*, middle coast region; *abnormis*, Lake Tahoe; *funnebris*, "widely diffused in Cal.;" *punctulata*, middle and northern; *lewisii* var. *monticola*, "Cal.;" *vittata*, "Cal."

**Formicomus.**—*F. mundus* is found under rubbish along the banks of the Colorado River, at Yuma, in July.

**Anthicus.**—*A. tenuis* is not uncommon at Yuma, in similar situations to the preceding: *pinalicus*, *canonicus*, *bipartitus*, and *turgidicollis* belong to the genus *Lappus* of Casey; they are arboreal in habit, *canonicus* being an especially abundant species in the foot-hills above Pomona, Pasadena, etc., occurring on many trees and shrubs. Specimens taken at Arrowhead Springs are referred with some doubt to *pinalicus*: *bipartitus* occurs at Yuma: *turgidicollis* is from "So. Cal.:" *californicus* is widely diffused, being known to me from Santa Monica, Long Beach, Riverside, Santa Catalina and San Clemente islands, Bakersfield and San Francisco: *confinis* is plentiful throughout our district; it is found about the roots of grass, on the borders of lawns, and often comes to light: *formicarius* is not frequent; I have seen specimens from Pomona and Pasadena, the only specimens taken by me being found in stable refuse: *rufulus* occurs at San Diego (Le Conte): *cribratus*, Riverside and Pomona, in May and June; not common: *hecate*, margins of streams, in the foot-hills near Pomona: *luteolus*, common along the Colorado River: *pinguescens*, common and widely dispersed from Southern California to the Columbia River: *ovicollis*, Los Angeles (Casey): *biguttulus*, Riverside, May and June; also middle California and Vancouver: *punctulatus*, abundant and widely dispersed in the true Pacific region: *seminotatus*, Los Angeles and Kern counties: *nanus*, San

Diego (Le Conte): *obliquus*, very common in Los Angeles County, coming often to light in swarms during May and June: *militaris*, mountains near Pomona, but not common: *bellulus*, Pasadena in March and San Diego; not common: *herifuga*, San Bernardino, Riverside, Pomona, Pasadena, etc.; rather common: *maritimus*, not abundant on the sea beaches; Long Beach, June: *corticalis*, Yuma: *obesus* is rather plentiful, and in much variety of size and color; various beaches of Southern California and at San Francisco. *A. albicans* and *A. parviceps* are merely varieties which would occur in any considerable series taken at any one locality. The following are either from the middle or northern regions or are mentioned without definite locality: *vigilans*, *nitidulus*, *franciscanus*, *nigritulus*, *protectus*, *mercurialis*, *præceps*, *auriger*, *inscitus*, *amænus*, *obscurellus* and *helvinus*.

**Tanarthrus.**—The writer has not yet met with *salinus*, which is said to occur around the southern salt lakes: *alutaceus* is not rare at San Diego and Redondo; I have taken specimens at the latter locality in March and April, on the muddy border of a small salt lake.

**Notoxus.**—*N. spatulifer*, Los Angeles County (Casey): *denudatus*, rare, Pomona and Los Angeles: *debilitans*, San Diego: *conformis*, Southern California (Horn): *sparsus*, not rare in the foot-hills near Pomona: *constrictus*, exceedingly abundant nearly everywhere: *robustus*, Los Angeles County: *alameda*, at times plentiful on the sand beneath and about the roots of plants, above the beach at Redondo: *cavicornis*, rare; found under bark and on the wing, at Pomona, in November and December: *calcaratus*, common and widely dispersed in



our region: *lustrellus* is described from San Francisco: *humboldti*, described from Humboldt County: *nevadensis* has been found at Tallac, by Dr. Fényes: *serratus* and *apicalis* are both credited to California by Dr. Horn, but there is reason to suspect that his specimens were not properly referred. Three other species of doubtful standing, but probably undescribed, are known to me from Southern California.

**Mecynotarsus.**—*M. delicatulus*, Yuma and Pasadena, April; several examples taken at electric light.

**Xylophilus.**—*X. brunnipennis* is not common; I have seen examples from San Bernardino, Pomona, and Pasadena. This is the species described by Casey as *saginatus*, but there appears to be no need for a new name. *X. brunnescens*, *X. nucleus* and *X. constrictus* are known only from one or two specimens each, and all are from the higher parts of the southern Sierras; *constrictus* has also been found at Lake Tahoe.

#### PYROCHROIDÆ.

**Pyrochroa.**—*P. californica* was recently described by Dr. Horn, from specimens taken in Los Angeles County by Coquillett; it is undoubtedly a rare insect.

#### MELOIDÆ.

**Cysteodemus.**—*C. armatus* occurs at many places on the Colorado Desert; taken in numbers at Yuma, in March, by Mr. Daggett.

**Meloe.**—*M. barbarus*, Santa Barbara Islands (Le Conte): *strigulosus*, a specimen so referred was taken by me at Palm Springs: *opacus* is northern.

**Nomaspis.**—*N. sublaevis*, from Fort Tejon.

**Poreospasta.**—*P. polita*, from "Southern Coast Range of Cal." (Gabb).

**Nemognatha.**—The species of this genus are moderately numerous on flowers, chiefly Compositæ, from June until October. There is much variation within specific limits, which renders the proper separation of species difficult. *N. apicalis* and *N. dubia* are common on sun-flowers, in September, at Pomona, Pasadena, etc.; two undescribed species occur with them, but more sparingly: *lutea* probably inhabits the Colorado Desert, but I have not yet recognized specimens with certainty: *nigripennis* is not common; I have seen specimens from Los Angeles, Pomona (May), and the San Bernardino Mountains (5,000 feet elevation), July: *scutellaris* is very common on low Compositæ, near Pomona, Pasadena, etc., in June and July: *dichroa* and *piezata* are reported from the State, but I have been unable to determine the exact localities.

**Gnathium.**—*G. nitidum*, Owens Valley.

**Zonitis.**—One example from Barstow is referred to *flavida*. I have seen a specimen in Dr. Van Dyke's collection of an apparently undescribed species from Southern California.

**Epicauta.**—I have specimens of *alphonsii* said to have been collected at or near San Diego; the species was described from Mariposa County: *puncticollis*, *oblita*, and *straba* are found on flowers in the foot-hills in Los Angeles County, from September to November; the first is always common, the second rare, the last is rather rare, though occasionally met with in abundance: *fallax* is found in Owens Valley: *maura*, San Diego: *maculata*, though no locality is named, doubtless occurs only in the deserts of the east or southeast.

**Cantharis.**—Three examples of *melæna* in my collection are from Southern California, one from Temescal, the other two without definite locality: *magister*, Colorado River, Owens Valley: *vulnerata*, common at times at Pomona and Pasadena, on flowers, in September; Owens Valley (Horn): *childii*, near Los Angeles, April (Van Dyke); in Compositæ; San Diego (Horn): *tenebrosa*, San Diego; Arrowhead Springs, April: *mærens*, Santa Monica, Owens Valley, Tejon, Sacramento: *insperata*, Mojave Desert and San Diego, in April: *occipitalis* and *incommoda* are recorded from "So. Cal.:" *stygica*, San Bernardino Mountains, at an altitude of 6,000 feet; "Oregon to the borders of the Colorado Desert:" *auriculata*, Riverside, Pomona, Pasadena, etc., not rare, on low Compositæ, in early spring: *aneipennis*, found in the same localities and at the same time as the preceding: *crotchii*, San Diego, March: *nitidicollis*, Pasadena, Riverside, San Diego in May; rare at the former places, more common at San Diego: *lugens*, not common at Pomona and San Diego in May: *compressicornis*, Owens Valley; Los Angeles County (Van Dyke): *cyanipennis*, *refulgens* and *rathroni* are found in middle California: *ulkei* is recorded simply from "Cal."

**Calospasta.**—*C. elegans*, Cariso Creek, western part of Colorado Desert, in San Diego County: *perpulchra*, rather rare on flowers, Pomona, in August and September: *mirabilis*, desert regions of Los Angeles and San Diego counties: *mæsta*, Southern California, exact locality not known: *nemognathoides*, Owens Valley and Yuma: *opaca*, desert or semidesert regions near Los Angeles: *fulleri* is simply reported from "Cal." but is probably, like nearly all the species of the genus, from the deserts of the east or southeast.

**Tegrodera.**—*T. erosa*, San Diego, Riverside, Owens Valley (race *latecincta*). Certainly very rare at Riverside, but common enough at times at various places on the Desert. Horn states that it occurs in Owens Valley in "the latter part of June, on a low plant bearing blue flowers."

**Phodaga.**—*P. alticeps*, Indio, Owens Valley.

#### RHIPIPHORIDÆ.

**Rhipiphorus.**—*R. flavipennis* and *R. cruentus* are rather uncommon, on flowers in the southern Sierras, at an altitude from 1,500 to 5,000 feet; I have taken them in the San Gabriel and San Bernardino ranges; var. *rufus* of the latter species has occurred at Los Angeles (Fuchs).

**Myodites.**—*M. californicus*, Los Angeles County (Van Dyke).

#### RHINOMACERIDÆ.

**Rhinomacer.**—*R. comptus* and *R. bombifrons* occur on conifers, in the Lake Tahoe region: *pilosus* is described from "Cal."

**Diodyrhynchus.**—*D. byturoides* is found in the middle Sierras (Placer County).

#### RHYNCHITIDÆ.

**Auletes.**—*A. nasalis* is found near San Diego, also in Los Angeles County; not common: *latifrons*, Southern California (Casey).

**Rhynchites.**—*R. bicolor*, Pasadena, May; San Bernardino Mountains, August; common wherever wild roses occur: *aureus*, Riverside, San Diego, and Ojai Valley, from February to May; San Clemente Island, June 1. A blue variety, supposedly of this species, has been taken in the San Bernardino Mountains. *R. ceratoides*

is abundant on *Eriogonum cinereum*, at Redondo, in April and May; it has also been found in the foot-hills near Pomona, but on what plant I am not now able to say: *velatus* occurs in the "Sierra Nevada Mts.—Morrison:" *naso* is described from "Cal."

**Deporaus.**—*D. glastinus* is common on live-oaks, especially when in blossom, throughout our region.

#### OTIORHYNCHIDÆ.

**Minyomeres.**—*M. languidus*, Fort Tejon (Horn).

**Stamoderes.**—*S. uniformis*, Sonoma County.

**Agasphærops.**—*A. nigra*, Mendocino County.

**Trigonoscuta.**—*T. pilosa* is abundant throughout the year in the sand-dunes along the ocean front.

**Eupagoderes.**—*E. argentatus*, *E. desertus* and *E. varius* inhabit the Colorado Desert, while *geminatus* and *plumbeus* occur farther north, in Owens Valley. It is almost certain that *plumbeus* is not distinct from *varius*. Specimens of a species of this genus have been found at Claremont, by Professor A. J. Cook; these are possibly *varius*, but so far as seen they are uniformly vittate and I am inclined to believe them different from anything described.

**Rhigopsis.**—*R. effracta* is common on and about the roots of various weeds, at Pomona, Pasadena, San Bernardino, etc. Horn reports it as having been found "feeding on yucca."

**Amotus.**—One example of *gracilior* was found on the beach at Santa Barbara, in February; it is not rare at Santa Monica (Albright): *longisternus* seems very close to the preceding, and like it was described from Los Angeles County.

**Dyslobus.**—*D. signis* is found only in the most northern parts of the State.

**Amnesia.**—*A. tumida*, Santa Cruz Mountains: *rauca*, *decidua*, *elongata*, and *tesselata*, from the vicinity of San Francisco: *sculptilis*, *discors*, and *granulata*, from the extreme north: *decorata*, middle Sierras.

**Adaleres.**—*A. ovipennis* and *A. humeralis* are quite common on live-oaks, especially in early spring. The differences named by Casey seem illusory and intercurrent in a large series.

**Nomidius.**—*N. abruptus*, "California."

**Orimodema.**—*O. protracta*, Mojave Desert (Horn).

**Nocheles.**—*N. vestitus*, Lake Tahoe (Fényes).

**Mimetes.**—*M. setulosus*, San Diego, April to June; Ventura: *seniculus* is probably from farther north, but the exact locality is not known to me.

**Phymatinus.**—*P. gemmatus*, "Cal. and Oreg."

**Miloderes.**—*M. setosus*, Kern County.

**Sciopithes.**—Specimens taken on San Clemente Island in June are identified by Casey as *S. setosus*, var.; the species was described from north of San Francisco (Marin and Napa counties). *S. obscurus* occurs from San Francisco north, and is quite common; *significans*, *brumalis*, *arcuatus*, and *angustulus* are described by Casey from the same region.

**Agronus.**—*A. cinerarius* is abundant on conifers, at Lake Tahoe: *deciduus*, San Francisco.

**Paraptochus.**—*P. sellatus*, foot-hills near Pomona, in June.



**Stenoptochus.**—*S. inconstans*, Riverside and Pasadena; swept from weeds, in April and May.

**Orthoptochus.**—*O. squamiger*, "Cal."

**Mylacus.**—*M. saccatus* is northern.

**Thricolepis.**—*T. inornata* is not rare in the San Bernardino Mountains, in July: *simulator* is described from Fort Tejon.

**Peritelopsis.**—*P. globiventris*, central California.

**Peritelodes.**—*P. obtectus*, Monterey County.

**Peritelinus.**—*P. variegatus*, Lake Tahoe.

**Geoderces.**—One example of *puncticollis* was taken at Pomona, in February. Two undescribed species, one of them quite common, have been taken at Pomona and Pasadena. *G. incomptus* occurs from San Francisco north.

**Geodercodes.**—*G. latipennis*, Monterey, in June. (Casey and Fényes).

**Aragnomus.**—*A. griseus*, Tulare County: *hispidulus*, Los Angeles County.

**Dysticheus.**—*D. insignis*, Southern California (Horn).

**Eucyllus.**—*E. vagans*, from the southeast.

**Thinoxenus.**—*T. squalens*, "Not rare near the seacoast of California."

**Rhyppodes.**—*R. dilatatus*, "near the seacoast at San Diego." I have an example taken near Pomona in June.

**Panormus.**—*P. setosus*, Monterey County.

**Dirotognathus.**—*D. sordidus*, Fort Mojave (Crotch).

**Elissa.**—*E. constricta*, Yuma.

**Aramigus.**—*A. fulleri* is rapidly spreading throughout Southern California and attacking a variety of trees and shrubs. I have found the beetles from September to November, most frequently on rose bushes.

**Scythropus.**—*S. californicus* is an abundant species in the middle Sierras. Specimens taken on pines in the San Bernardino Mountains are probably distinct, but are left here for the present: *lateralis* is described from Lake County, *ferrugineus* from Marin County, *cinereus* from Lake Tahoe.

### CURCULIONIDÆ.

**Sitones.**—*S. sordidus*, "San Francisco and San Diego;" it is unknown to the writer: *crinitus*, one example taken at Pomona is so referred: *nebulosus*, one example, Pomona, July: *hispidiceps* is the commonest species in the vicinity of Los Angeles; identified by Capt. Casey, who writes that his *prominens* and *angustulus* are to be considered synonymous with this species: *explicitus*, not rare at San Diego in February. I have taken one specimen of a very distinct and undescribed species in the San Bernardino Mountains in July. The following are credited to the State, but their recognition is practically impossible as our literature now stands: *californicus*, "Cal. and Oreg.;" *vittatus*, San Francisco; *varians*, *sparsus* and *osculans*, Humboldt County; *margaritosus*, Santa Cruz and Monterey; *procerus*, Napa; *occidentalis*, Sonoma; *montanus*, Placer County; *alternans*, "Cal.;" *apacheanus*, "So. Cal. or Ariz."

**Triglyphus.**—*T. ater*, middle Sierras; a fine species which appears to be decidedly rare.

**Apion.**—*A. pennsylvanicum* is one of our rarer species, occurring at Pasadena in May and August; it is widely

diffused throughout the United States, but much more common in the Atlantic district: *hesperum* has been found only by Coquillett in the desert portions of Los Angeles County: *sordidum* is an abundant species in the desert but very uncommon west of the mountains; specimens have been taken in May at Pomona and Pasadena; it has been bred from galls on several occasions—notably by Mr. Koebele from “Cecidomyid and dipterous galls on *Artemisia californica*,” and by Mr. Schwarz from “bud-like gall on stems of *Hymenoclea monogyra* in Arizona: *antennatum* is not rare, on pines in both the southern and middle Sierras, July; taken also at Pomona in January, Catalina in July, and Ventura in March: *ædorhynchum* is common throughout our region, April to October: *opacicolle*, Ojai Valley, March; uncommon: *troglydytes* is plentiful from San Diego to San Francisco, December to May: *proclive* is abundant nearly everywhere both east and west of the mountains; it has been beaten from willows at Pomona, March to October: *cribricolle* is widely distributed throughout the State, and is generally abundant; common on *Eriogonum cinereum* at Redondo in Spring, but occurs on a variety of plants and trees. *A. cordatum* is rare at Pomona, where specimens have been beaten from willows in October: *varicorne* is plentiful at Yuma, in July, on a species of *Dalea*: *ventricosum* is very abundant on the desert, occurring on mesquite: *attenuatum* is rather rare; specimens have been taken at Pomona, Pasadena and Riverside. Two other species—*protensum* and *walshii*—have been found within the State; the former is known by the unique type taken at San Francisco, while the latter has been found only in the north.

**Podapion.**—*P. gallicola*, middle and northern Sierras; not common.

**Lepyrus.**—*L. perforatus* is northern.

**Listronotus.**—Two undescribed species of this genus are known from our district. One of these is near *terestrois*; it has been taken at San Diego, Pomona, and Antelope Valley. Of the second species, a small form near *gracilis*, one example only has been found by me at Long Beach.

**Macrops.**—*M. hyperodes* and *M. californicus* are known from the central or northern parts of the State.

**Emphyastes.**—*E. fucicola* is not uncommon along the sea beaches, beneath sea-weed, etc.

**Lixus.**—*L. asper*, not frequent at San Bernardino, on *Rumex* (Wright): *perforatus*, not rare at Pomona, March to June: *semivittatus*, specimens which do not appear to differ from typical Arizona examples of this species are not rare on and about the roots of a plant growing along the sand-dunes near the beach at Santa Monica: *auctus* is northern: *parvus* is described from San Francisco.

**Dinocleus.**—One specimen of *pilosus* found at Coronado in July is scarcely different from the Le Conte type. The type is without exact locality but is doubtless from the south, and probably from the same region. *D. jacobinus* is described by Casey from San Diego: *fartus*, from "near the southern boundary:" *albovestitus* is not rare near Los Angeles in August and September: *molitor* is common on the desert at Indio and Yuma: *wickhami* was taken by Mr. Wickham at Indio.

**Cleonus.**—*C. inornatus*, Owens Valley: *pacificus*, one example at Redondo in April, on *Astragalus crotalaræ*: *erysimi*, not rare at Redondo on flowers of *Erysimum asperum*, March and April: *lobigerinus*, from the southeast: *vittatus*, Owens Valley, if Casey is correct in his

disposition of Le Conte's *virgatus*: *modestus*, common at Pomona, Ojai Valley, etc., in flowers of *Lupinus* in spring; specimens apparently identical have been collected in the San Bernardino Mountains: *pleuralis*, one example taken at Pomona in January.

**Dorytomus.**—*D. inaequalis*, Los Angeles (Casey); San Diego, March; probably affects cottonwood as do the closely related *mucidus* and *brevipilosus* of the East: *hystricula* is very abundant: *luridus* is tolerably common on willows when in blossom; Pomona, Pasadena, Riverside, etc.: *hirtus* was described from San Diego; this species is either a rarity or else an unusual form of the common thing which Casey has called *hystricula*: *rufus*, one example beaten from willows at Yuma: *cuneatulus* and *marginatus* are described by Casey from more northern parts of the State.

**Pachyphanes.**—*P. carus* is described from "Cal.;" it is probably not from our region: *corpulentus* is represented by a single specimen, so named for me by Dietz, taken near San Francisco in August.

**Desmoris.**—*D. constrictus* is abundant everywhere on sunflowers: *fulvus* and *sordidus* are reported from the State, and it is quite probable that both are found in the desert regions of the east or southeast. Dietz also describes *incertus* from the State.

**Smicronyx.**—*S. pusillus*, San Bernardino (collections of Horn and Dietz): *cinereus*, common and widely diffused: *instabilis*, rather rare at Pomona and Pasadena: *defricans*, not common, same localities as preceding: *nubilis*, Pasadena, Redondo, Palm Springs; not rare: *californicus*, Pomona, Pasadena, Palm Springs; April to August: *obtectus*, Pomona, Long Beach, San Diego:

especially abundant at the latter place in March. Two or three undescribed species are in my collection. All the species are obtained, and usually in numbers, by sweeping flowers in spring and early summer. Of the following, also from the State, *seriatus* was originally from Mariposa, while *perplexus*, *tardus*, *resplendens*, *pusio*, *scalator*, and *ornatipennis* are not definitely located.

**Synertha.**—*S. imbricata*, Mojave; San Bernardino Mountains, on pines.

**Promecotarsus.**—*P. maritimus*, San Diego.

**Phyllotrox.**—*P. nubifer*, Santa Barbara, February; Lake Tahoe, July.

**Endalus.**—One example of *limatulus* from Los Angeles County (Van Dyke): *ovalis*, Pomona; found about the roots of grasses in swampy places in winter: *aratus*, Yuba County.

**Stenopelmus.**—*S. rufinasus*, Santa Monica and Ventura, February and March; not common, San Diego (Crotch and Le Conte).

**Bagous.**—*B. californicus*, San Diego (Crotch).

**Phycocætes.**—*P. testaceus* is common at times under seaweed, from Santa Barbara to San Diego.

**Schizonotus.**—*S. cæcus*, San Francisco. The species was discovered by Mr. Fuchs, by sifting mouldy earth among the redwoods north of the Bay.

**Otidocephalus.**—*O. vittatus*, Owens Valley (Horn).

**Magdalis.**—*M. cuneiformis* and *M. lecontei* may be beaten from coniferous trees along the higher portions of the southern Sierras. The former is comparatively



scarce, the latter not rare in the San Bernardino Mountains in July. An undescribed species depredates upon alders in the locality just mentioned. *M. gracilis* is apparently not rare in central and northern parts of the State: *subtincta* is described from Gilroy (Crotch): *gentilis* occurs at Lake Tahoe.

**Magdalinops.**—*M. vittipennis*, "Cal."

**Macrorhoptus.**—*M. hispidus*, San Diego, June; Pomona, May: *estriatus* is recorded from Santa Barbara and Warner's Ranch (Crotch).

**Tachypterus.**—*T. quadrigibbus*, "Cal."

**Cionistes.**—One example of *insolens* was beaten from a sycamore at Pomona in October. I have seen numerous specimens collected near Bakersfield in April.

**Anthonomus.**—Several specimens of *peninsularis* were taken at Palm Springs in April by Coquillett: *confusus*, Los Angeles County; rare: *sycophanta* is common on willows in the vicinity of Los Angeles: *apertus*, Los Angeles, Riverside, Pasadena; not common; taken by sweeping weeds along roads in April and May: *albopilosus* is from the southeastern portions of the State: *ceneolus* is usually rare, but was taken in some abundance during October and November of several successive years, on *Solanum nigrum* in a cañon near Pomona: *ater*, one example, on sunflowers at Pomona, April: *ochreopilosus*, Riverside and San Diego; quite common near the latter place in May: *pauperculus*, common at Riverside, Pomona, San Diego, and many other places in our district; taken by sweeping in waste places: *ornatulus* is found in similar places but is less common than the preceding: *figuratus* is unknown to me; it is said to have been taken at Santa Monica: *inermis*,

southern Sierras from 3,000 to 5,000 feet altitude; not rare: *stolatus*, San Diego; several examples from Mr. George H. Field. The species following are also reputed to be Californian: *brunnipennis*, Geysers; *morulus*, near San Francisco; *effetus*, *squamosus*, *hirtus*, *subvittatus*, *canus*, and *affinis* are recorded from "Cal." simply; *tahoënsis* is described from a small series taken at Tallac (Lake Tahoe) by Dr. Fényes. Several of these doubtless were taken in the desert regions of the east and southeast, but I suspect that the identifications are not exact in all cases, and it would be well to await further evidence before placing the names in our list.

**Epimechus.**—*E. mimicus* is not common; taken by sweeping in the foot-hills near Pomona in May and June: *soriculus* is found in the same localities as the preceding but is still less common; January, April, June, also Long Beach in April: *nevadicus* occurs a little more frequently than either of the preceding; Pomona, April to July: *æmulus*, one example from San Diego: *adpersus* has been taken by Dr. Fényes at Monterey.

**Elleschus.**—*E. ephippiatus* is rather plentiful on willows in spring; Riverside, Ojai Valley, etc.

**Orchestes.**—*O. parvicollis* is common at Los Angeles, Pomona and San Diego from June to August: *puberulus* is not rare in the southern Sierras and the foot-hills near Pomona, Pasadena, etc: *salicis* is found near San Francisco: *ephippiatus* and *rufipes* are said to occur within the State; it is probable that they are northern or central.

**Tychius.**—*T. lineellus* is abundant in flowers of *Lupinus* at Pomona and in the Ojai Valley; March to

May. Of *T. prolixus* one specimen has been found, also on *Lupinus*, at Pomona in May: *setosus* is obtainable in numbers on mesquite in the Colorado Desert; and with it is what appears to be an allied, undescribed species: *semisquamosus* is reported from Tejon.

**Sibynes.**—*S. fulvus*, San Bernardino County.

**Paragoges.**—*P. maculatus* is an uncommon insect which was originally taken by Crotch at San Diego; I have seen specimens from Redondo, San Nicolas Island, and Alameda County.

**Conotrachelus.**—*C. duplex*, “California — probably southern” (Casey).

**Rhyssematus.**—*R. pubescens* is a rare species of which I have seen but two examples; one swept from flowers near Pomona in June, the other collected at electric light at Pasadena in April. Horn’s type was obtained in Owens Valley.

**Micromastus.**—*M. gracilis*, San Francisco.

**Tyloderma.**—*T. morbillosum*, San Francisco.

**Zascelis.**—*Z. irrorata*, “Cal., Nev., Col.”

**Piazurus.**—*P. californicus* is abundant on pines in the Sierras.

**Copturus.**—The type of *mamillatus* was probably from San Diego County, though the exact locality is not known. The only other examples seen by me were taken by sweeping weeds along the Colorado River at Yuma, and a single specimen from Ensenada, Lower California. *C. adspersus* is our commonest species; it is obtained by sweeping weeds in waste places, and is wide-spread: *longulus* is found only on coniferous trees in the higher parts of the Sierras. In the northern

and central parts of the State are found *koebeleri*, *lunatus*, *nubilatus* and *mucidus*. There are two undescribed species from our region in my collection.

**Gyrotus.**—*G. munitus*, "So. Cal." (Casey).

**Acanthoscelis.**—*A. californicus*, *A. frontalis* and *A. perplexus* are all given by Dietz as Californian, but without exact locality; they are probably not southern.

**Ceutorhynchus.**—*C. subpubescens*, "Los Angeles and Tejon" (Crotch): *angulatus*, *disturbatus* and *obliquus* are all found by sweeping low herbage—most often near streams—in or along the base of the Sierras; the first named is least common: *hornii* is described from "So. Cal.:" *nodipennis*, Los Angeles County, on *Ceanothus* (Coquillett): *mutabilis*, Little Bear Valley, June (Daggett): *convexicollis*, common at Pasadena on a cruciferous plant, February and March: *albopilosus*, Los Angeles: *pervestitus*, sp. nov., one example taken near Bakersfield: *isolatus*, "Montana, Elko, Nev. and Cal.:" *sericans*, middle Sierras: *ovipennis*, Dunsmuir: *decipiens*, "Cal.:" *cyanipennis*, northern: *pollinosus*, "Cal.:" *pusio*, San Francisco and north: *adspersulus*, "Cal. and Ariz.:" *pusillus*, northern: *puberulus*, near San Francisco.

**Cælogaster.**—One example of *zimmermanni* is from the San Bernardino Mountains.

**Pelenomus.**—*P. cavifrons*, Pomona, Riverside, etc., April to June; about the roots of weeds and under vegetable debris near water.

**Baris.**—*B. dilatata*, San Bernardino, Long Beach, Santa Monica, Yuba County: *rubripes*, San Bernardino, Pomona, Yosemite Valley, Santa Rosa: *futilis*, not very common; on willows at Riverside, Pomona, etc., March,

April and November: *heterodoxa*, one example, Los Angeles County: *monticola*, Bear Valley, altitude 6,400 feet (Daggett): *tenuestriata* and *macra* are found near San Francisco: *sparsa* is northern: *opacula* and *brunneipes* are without exact locality.

**Onychobaris.**—*O. densa* is abundant in the flowers of a low plant growing on the sand-dunes at Coronado, July and August: *depressa*, Santa Monica: *austera*, San Diego: *insidiosa* is rather common at Pomona, Pasadena and San Diego: *arguta*, less common than the preceding; foot-hills near Pomona and Pasadena: *audax*, Southern California: *seriata* occurs at San Francisco.

**Pseudobaris.**—*P. nigrina*, Lake County (Casey).

**Trichobaris.**—*T. mucorea* and *T. compacta* are found in the flowers of *Datura metilloides* in April and May. The former is less common.

**Orthoris.**—*O. crotchii*, "California."

**Centrinus.**—*C. lineellus*, Tejon.

**Centrinogyna.**—*C. procera*, San Francisco.

**Limnobaris.**—*L. nasuta*, Pasadena, April and May; on willows: *seclusa*, "So. Cal.;" I have one example from Kern County.

**Barinus.**—*B. difficilis*, Southern California.

**Barilepton.**—*B. falciger*, San Bernardino.

**Balaninus.**—*B. uniformis*, not rare at Pomona, Pasadena, etc.; also found on Santa Catalina Island affecting several species of oaks.

## CALANDRIDÆ.

**Scyphophorus.**—*S. acupunctatus* is not rare in San Diego County: *robustior* is also reported from San Diego County (Horn\*): *yucca* is abundant in the vicinity of Los Angeles. All depredate on yucca.

**Metamasius.**—*M. sericeus* is said to have been collected by Gabb in the southern Coast Range.

**Cactophagus.**—*C. validus*, San Diego, under decomposing *Opuntia*; rare.

**Rhodobænus.**—*R. tredecimpunctatus*, Riverside, May and June; not common.

**Sphenophorus.**—*S. simplex* is moderately common and generally dispersed throughout our district from Yuma to the coast: *vomerinus*, Yuma, July; San Diego, February: *pictus*, Santa Monica, Vallecito; not common: *pertinax*, Riverside, May; Santa Monica; abundant about the roots of grass or reeds in low ground: *sayi*, one example at Long Beach in May: *placidus*, specimens from Santa Monica are so referred: *gentilis*, Pomona; one example: *tardus*, not rare at San Bernardino.

**Calandra.**—*C. granaria* and *C. oryza* are both occasionally found in Los Angeles County depredating on cereals.

**Yuccaborus.**—*Y. frontalis*, Antelope Valley (Los Angeles County); on yucca.

**Dryophthorus.**—*D. bituberculatus* is said by Boheman to occur in California.

**Metopotoma.**—*M. repens* is northern.

---

\* "The Coleoptera of Baja California," Proc. Cal. Acad. Sci., 2nd Ser., Vol. IV, 1894, p. 359.



**Cossonus.**—*C. crenatus* is abundant under pine bark, in the southern Sierras: *piniphilus* is found in similar situations farther north; my specimens are from the middle regions.

**Macrorhyncholus.**—*M. protractus* is common in the foot-hills of the Sierra Madre Mountains; the larva lives in the flower stalks of *Yucca whipplei*.

**Elassoptes.**—*E. marinus* is found on the sea beach; Santa Monica and San Pedro.

**Rhyncholus.**—*R. dorsalis*, San Diego: *angularis*, Colorado Desert, under willow bark: *oregonensis*, San Bernardino Mountains: *R. spretus*, San Bernardino Mountains, on freshly cut alder. Casey has described *pallens* from the vicinity of San Francisco, and *dilatatus* from "Cal." Wollaston has described from California, *cylindricollis*, *californicus*, and *protensus*, all of which are unknown to American entomologists; but there is doubtless some synonymy involved in Casey's recent descriptions.

#### SCOLYTIDÆ.

**Monarthrum.**—*M. scutellare* and *M. dentigerum* have both been taken in March from burrows extending into the solid wood of live-oak; the latter species has been taken on the wing from April to June.

**Gnathotrichus.**—*G. retusus* is found under pine bark, San Bernardino Mountains, August.

**Pityophthorus.**—*P. pubipennis* affects alders; swarms have been seen flying in August in the San Bernardino Mountains: *carinulatus*, *confinis*, and *digestus* were all taken by Crotch in the Mojave region. The first-named species as well as *nitidulus* and *puncticollis* occur in the

middle Sierras. Le Conte describes *pilosulus* from "Middle California."

**Hypothenemus.**—*H. striatus* is recorded from both Lower and upper California, and is therefore undoubtedly from our district. An undescribed species is frequently taken in spring, at Pomona, by sweeping and beating in waste places.

**Xyloterus.**—*X. lineatus*, Sylvania, April (Ricksecker).

**Cryphalus.**—*C. terminalis* is described from California, but is not recognized by any of our authors.

**Xyleborus.**—Specimens of *xylographus* have been taken on the wing in spring at Pomona, Pasadena, and Ojai Valley: *celatus* is represented by a single example taken in the San Bernardino Mountains.

**Dryocætes.**—*D. septentrionis*, Sylvania, April (Ricksecker).

**Xylocleptes.**—*X. cucurbitæ* is widely diffused; the author has taken it at Pomona from April to November.

**Tomicus.**—*T. confusus* is moderately common under pine bark, San Bernardino Mountains: *plastographus* and *latidens* occur in the middle Sierras.

**Micracis.**—*M. hirtella*, Pomona; not common. An undescribed species occurs in Los Angeles County (Coquillett).

**Scolytus.**—Specimens referred to *ventralis* have been taken on Douglas Spruce, in the San Bernardino Mountains. Specimens have also been received from Mendocino County, where they affected the same tree according to Dr. Van Dyke. *S. praeceps* is reported from Calaveras (Crotch): *subscaber* is from the north: *californicus* is without other locality than that indicated by the name.

**Hylesinus.**—*H. aspericollis* is not rare in the southern Sierras: specimens referred to *imperialis* have been seen from the vicinity of San Francisco: *sericeus* is found in the middle and northern Sierras.

**Phlæosinus.**—*P. punctatus* is common throughout the Sierras: *cristatus* occurs farther north; it has proved injurious to cypress hedges near Sacramento.

**Chætophlæus** —*C. hystrix*, San Diego; rare.

**Carphoborus.**—*C. simplex*, Mojave Desert.

**Dendroctonus.**—*D. terebrans* is common and *D. frontalis* rather scarce; on pines in the Sierras: *similis* and *simplex* are said to occur in the State.

**Hylastes.**—*H. macer* and *H. longus* are common in the coniferous forests throughout the State: *gracilis* is found in the Tahoe region: *nigrinus* is common in the more northern parts of the State; I have seen one example in Mr. Fuchs' collection labeled San Diego County.

**Hylurgops.**—*H. subcostulatus* occurs throughout the Sierras: *granulatus* and *rugipennis* are thus far known only from the more northern parts of the State.

#### ANTHRIBIDÆ.

**Toxotropis.**—*T. approximatus* was beaten from dead twigs of live-oak, near Pomona, in June.

**Gonops.**—*G. fissunguis*, "Big Trees."

**Brachytarsus.**—I have seen four species from the State, none of which are identified with certainty; one of these is certainly exceedingly close to, if not identical with, the eastern *alternatus*; specimens have been taken at Redondo by sweeping. Two other species are from Southern California.

The following table shows by families the number of genera and species mentioned in the preceding pages, both for Southern California and for the entire State. Between these is introduced for comparative purposes similar data from Dr. Hamilton's List of the Coleoptera of Southwestern Pennsylvania. The number of species named by Dr. Hamilton being substantially equal to the number here recorded from Southern California, a comparison of figures will give a very good idea of the relative abundance of the various families on the Atlantic and Pacific slopes. A few numbers in the Pennsylvania list are marked with an asterisk (\*) to indicate that in these cases the record is incomplete:—

FAMILY.	SOUTHERN CALIFORNIA.		SOUTHWESTERN PENNSYLVANIA.		CALIFORNIA.	
	Genera.	Species.	Genera.	Species.	Genera.	Species.
CICINDELIDÆ.....	2	15	1	6	3	20
CARABIDÆ.....	49	188	61	263	63	318
AMPHIZOIDÆ.....	1	1	.....	.....	1	1
HALIPLIDÆ.....	2	3	2	2	2	3
DYTISCIDÆ.....	19	41	20	33	19	60
GYRINIDÆ.....	2	3	2	6	2	4
HYDROPHILIDÆ.....	19	52	14	38	19	70
PLATYPSYLLIDÆ.....	.....	.....	.....	.....	.....	.....
LEPTINIDÆ.....	.....	.....	1	1	.....	.....
SILPHIDÆ.....	10	20	10	30	17	37
SCYDMÆNIDÆ.....	5	6	6	21	9	17
PSELAPHIDÆ.....	13	20	23	56	21	61
STAPHYLINIDÆ.....	83	277	71	200*	107	462
TRICHOPTERYGIDÆ.....	5	10	3	3*	6	20
HYDROSCAPHIDÆ.....	1	1	.....	.....	1	1
SPHÆRIIDÆ.....	1	1	.....	.....	1	1
SCAPHIDIIDÆ.....	1	2	3	6	1	2
PHALACRIDÆ.....	4	11	3	7*	4	12
CORYLOPHIDÆ.....	4	7	5	7	4	7
COCCINELLIDÆ.....	18	56	15	34	21	71
ENDOMYCHIDÆ.....	2	3	8	10	4	7
EROTYLIDÆ.....	3	3	8	19	3	3
COLYDIDÆ.....	10	13	12	14	14	24
RHYSSODIDÆ.....	1	1	2	2	1	1
CUCUJIDÆ.....	8	16	9	19	10	21
CRYPTOPHAGIDÆ.....	3	9	8	13	3	10
MYCETOPHAGIDÆ.....	6	7	5	11	6	7
DERMESTIDÆ.....	8	18	8	13	8	23
HISTERIDÆ.....	16	63	15	46	17	77
NITIDULIDÆ.....	14	25	22	48	19	35
LATHRIDIIDÆ.....	10	30	5	8*	11	43
TROGOSITIDÆ.....	5	6	6	10	5	10
MONOTOMIDÆ.....	4	6	3	6	5	7
DERODONTIDÆ.....	.....	.....	1	1	.....	.....
BYRRHIDÆ.....	4	8	3	4	5	12
GEORYSSIDÆ.....	.....	.....	.....	.....	1	1
PARNIDÆ.....	6	9	6	9	8	15
HETEROCERIDÆ.....	1	4	1	6	1	5
DASYLLIDÆ.....	7	10	9	15	12	18
RHIPIDERIDÆ.....	1	1	.....	.....	1	1
ELATERIDÆ.....	29	78	38	127	40	162
THROSCIDÆ.....	2	4	3	3	3	6
BUPRESTIDÆ.....	18	81	15	48	18	94

FAMILY.	SOUTHERN CALIFORNIA.		SOUTHWESTERN PENNSYLVANIA.		CALIFORNIA.	
	Genera.	Species.	Genera.	Species.	Genera.	Species.
LAMPYRIDÆ.....	14	21	24	61	16	43
MALACHIDÆ.....	20	93	4	9	20	162
CLERIDÆ.....	10	26	14	27	12	30
PTINIDÆ.....	26	51	23	41	32	72
CUPESIDÆ.....	1	1	1	2	1	1
LYMEXYLIDÆ.....	.....	.....	1	1	.....	.....
CIIDÆ.....	3	6	3	7	5	9
SPHINDIDÆ.....	1	1	2	2	1	1
LUCANIDÆ.....	1	1	6	7	3	9
SCARABÆIDÆ.....	33	92	37	112	38	135
SPONDYLIDÆ.....	2	2	1	1	2	2
CERAMBYCIDÆ.....	62	85	74	163	73	152
CHRYSOMELIDÆ.....	50	124	65	152	61	171
BRUCHIDÆ.....	2	18	1	6	2	20
TENEBRIONIDÆ.....	68	187	30	52	75	247
ÆGIALITIDÆ.....	.....	.....	.....	.....	1	2
CISTELIDÆ.....	5	14	8	23	5	23
OTHNIIDÆ.....	1	1	.....	.....	1	1
LAGRIIDÆ.....	.....	.....	2	2	.....	.....
MONOMMIDÆ.....	1	1	.....	.....	1	1
MELANDRYIDÆ.....	4	5	20	28	6	10
PYTHIDÆ.....	1	2	5	6	4	6
ÆDEMERIDÆ.....	5	10	3	4	9	18
CEPHALOIDÆ.....	.....	.....	1	1	2	3
MORDELLIDÆ.....	4	20	5	36	4	24
ANTHICIDÆ.....	9	55	7	28	9	81
PYROCHROIDÆ.....	1	1	2	5	1	1
MELOIDÆ.....	10	46	5	10	12	56
RHIPIPHORIDÆ.....	2	3	1	1	2	3
STYLOPIDÆ.....	.....	.....	.....	.....	.....	.....
RHINOMACERIDÆ.....	.....	.....	.....	.....	2	4
RHYNCHITIDÆ.....	3	6	4	5	3	8
ATTELABIDÆ.....	.....	.....	1	3	.....	.....
BYRSOPIDÆ.....	.....	.....	.....	.....	.....	.....
OTIORHYNCHIDÆ.....	20	30	9	10	38	66
CURCULIONIDÆ.....	42	133	59	143	59	218
BRENTHIDÆ.....	.....	.....	1	1	.....	.....
CALANDRIDÆ.....	10	24	8	16	13	29
SCOLYTIDÆ.....	16	26	17	28	19	45
ANTHRIBIDÆ.....	2	4	6	8	3	6

Totals { Southern California, Families 70, Genera 826, Species 2,197  
 { Southwestern Pennsylvania, " 71, " 877, " 2,153  
 { California, " 74, " 1,031, " 3,408



## DESCRIPTIONS OF NEW SPECIES.

1. *Notiophilus obscurus*, sp. nov.

Black, bronzed, sides of elytra broadly but obscurely yellow toward the apex; base of antennæ, tibiæ and tarsi more or less pale. Head slightly wider than the thorax, front 8-10 striate. Striæ of elytra scarcely impressed, not effaced at apex, slightly approximated by pairs, the external interspaces faintly alutaceous but shining; there are normally three discal tuberculate punctures or foveæ, one before the middle and two subapical. Prothorax beneath very sparsely punctate. Length 5-5.5 mm.

*Habitat*:—San Bernardino Mountains, at an altitude of from 6,000 to 7,000 feet.

The above characters are about the only ones that are of general use for comparative purposes in this genus, and therefore a more detailed description is unnecessary. *N. obscurus* seems among our species to be most nearly related to a form occurring in Vermont and the White Mountains of New Hampshire, and which passes as *N. sylvaticus* in collections; this latter differs however by having the sides of the elytra yellow throughout, the striæ impressed and more strongly punctate, the intervals narrower, and the front less numerously striate. The White Mountain species is quite surely not the true *N. sylvaticus*, which is a west coast species (Alaska, Vancouver, Washington) with smaller head and less strongly sculptured elytra, and it is quite as certain that it is very close to the European *N. biguttatus*. The latter agrees very nearly with *N. obscurus* in the color of the elytra, but is otherwise scarcely distinguishable from the New England species.

2. *Dyschirius unipunctatus*, sp. nov.

Small, moderately stout, black with distinct bronze-green lustre, legs and antennæ rufous or brown, the latter somewhat darker apically.

Epistoma bisinuate, the middle tooth much smaller than the outer ones, but always distinct though somewhat varying in prominence; front not very strongly transversely impressed opposite the anterior margin of the eyes, the impression interrupted at middle by a short fine carina; vertex smooth. Prothorax transversely globose, very slightly wider posteriorly. Elytra just perceptibly wider than the prothorax, a little more elongate than in *D. globulosus*; base distinctly margined, striæ fine, very slightly abbreviated at base, gradually finer posteriorly, and almost completely effaced at apex; punctures of striæ fine and somewhat distant, becoming obsolete not far behind the middle; third interspace with a single setigerous puncture near the base. Terminal process of the anterior tibiæ only slightly curved, and a little shorter than the subapical interior spur; outer margin with a moderate subapical tooth and a second fainter one above it.

Length, 2.5 mm.

*Habitat*:—Pomona, San Bernardino, Riverside.

Rather smaller than any other species in our fauna and distinguishable from all by the unipunctate third interval of the elytra.

Some years ago in a letter from a friend, Mr. F. Blanchard of Tyngsboro, Mass., the writer's attention was called to the presence of two small teeth in the clypeal emargination of certain of those species of *Dyschirius* in which the elytra are pale at tip. Mr. Blanchard at that time expressed a doubt as to the propriety of referring all eastern specimens to *D. hæmorrhoidalis*, and a recent comparison of my material with the types of *D. terminatus* and *D. analis*, and presumably of authentic specimens of *D. hæmorrhoidalis*, leads me to believe that we may with reasonable certainty add two species to the three now listed. All these except *D. terminatus* have the clypeal emargination bidentate, and are, in fact, so closely allied in all respects except those mentioned in the table below that a more detailed description of the new forms is unnecessary. There are certain differences in the form of the prothorax, which, however, are not so easily appreciable as the characters

mentioned, and being moreover subject to some individual variation are difficult to define.

Striæ of elytra effaced at apex.

Clypeal emargination truncate at middle; front rather deeply and sharply transversely impressed, vertex smooth. [New York, Kansas, California (Le Conte)].....*D. terminatus* LEC.

Clypeal emargination bidentate at middle.

Strial punctures coarse, front and vertex smooth, front rather deeply transversely impressed; size larger. [Kansas, Ohio (Le Conte)].....*D. hæmorrhoidalis* LEC.

Strial punctures comparatively fine, front and vertex rugose. front scarcely or feebly transversely impressed, size smaller. (Texas—Luling and El Paso).....*D. duplicatus*, sp. nov.

Striæ of elytra very feeble but traceable at apex; strial punctures fine, front and vertex rugose, front scarcely or feebly transversely impressed; clypeal emargination bidentate. (Massachusetts, eastern New York).....*D. affinis*, sp. nov.

Striæ distinct at apex, and coarsely punctate; front smooth, not distinctly transversely impressed; clypeal emargination bidentate. (California—Colorado River).....*D. analis* LEC.

Of the above species, *D. duplicatus* and *D. affinis* are most closely related and may prove to be geographical races of the same species; both are quite distinct from *hæmorrhoidalis*.

### 3. *Schizogenius seticollis*, sp. nov.

This name is proposed for a species which has been quite widely distributed under the name *S. pluripunctatus*, from which and from all other species known it is distinguished by the prothorax having seven or eight marginal setæ instead of the usual two. The form is only moderately depressed, the color black, legs piceous or rufopiceous, the tibiæ and tarsi usually paler. The alternate elytral intervals beginning with the third are 10–12 setose.

Length, 4–5 mm.

Generally distributed west of the mountains and rather common at Pomona, Pasadena, Riverside, and other places in Southern California, and as far north as Sylvania (Ricksecker).

#### 4. *Schizogenius validus*, sp. nov.

Elongate, stouter than usual, polished black, antennæ and legs brown. Head and prothorax proportioned and sculptured above nearly as in *S. depressus*. Elytra parallel in basal three-fifths, rather strongly striate, striæ distinctly punctate, the punctures becoming smaller toward the apex, where they are nearly obsolete; third, fifth and seventh intervals 5-7 setigerous punctate. Legs about as usual; front tarsi strongly dilated (at least in the male), joints 2-4 being nearly or quite twice as wide as long, the fifth joint distinctly narrowed from base to apex; the claws one-half the length of the joint.

Length, 6.5 mm.

Several examples taken by Dr. H. G. Griffith on the Rio Verdi in central Arizona.

This is our largest species. It most nearly resembles in size and color *S. seticollis*, but is larger, with more distinctly punctate elytral striæ; the intervals with fewer, and the thorax with but the two usual setigerous punctures. The throat is distinctly less punctate than usual, and the front tarsi are more widely dilated than in any other species known to me.

#### 5. *Schizogenius litigiosus*, sp. nov.

This name is suggested for still another form, occurring in middle and northern California, which has heretofore passed as *S. pluripunctatus*. It is really very closely allied structurally to *S. depressus*, but seems to differ constantly in its black color, in the antennæ, which are a trifle more slender, and in the sides of the throat, which are a little more densely punctate. In a series the thorax appears a little less broad than in *S. depressus*. The differences aside from color are slight, but the writer feels quite confident that it is not the same as *S. depressus*, which is always brown, and more southern in habitat.

The true *S. pluripunctatus* is a very different thing, separable from all our other species by the impunctate elytral striæ, and by the very numerous setæ (15 or more) of the intervals.

Mr. Blanchard observes *in litt.* that the males of all species of the genus known to him have two setæ near the anterior margin of the last ventral, which are lacking in the females.

#### 6. *Trechus pomonæ*, sp. nov.

Very near *T. ovipennis*, from which it differs in its slightly more transverse prothorax, longer elytra with distinct indication of humeral angle, traces of striæ excessively faint, two dorsal punctures—basal and apical—and in the antennæ, which are a trifle less stout.

From *T. barbæ* it must differ in its more transverse thorax, distinct frontal impressions, two dorsal punctures (instead of three), and smaller size.

Three examples seen from Pomona and Pasadena.

#### 7. *Pterostichus inermis*, sp. nov.

Moderately elongate; piceous black, shining; legs dark rufous or rufopiceous. Frontal foveæ distinct. Prothorax wider than long, widest at about the apical third; sides rather narrowly margined, strongly arcuately narrowed behind, not in the least sinuate before the hind angles, which are obtuse and rounded; base three-fourths as wide as the apex, margined at the sides; apex feebly emarginate, anterior angles not prominent, rounded; basal impressions rather feeble, the outer entirely wanting. Elytra scarcely wider than the thorax, parallel, sides feebly rounded; humeri slightly dentate; striæ fine, impunctate; scutellar striæ short, joining the sutural stria. Legs moderate; front face of middle thighs bisetigerous; first joint of hind tarsi with lateral groove. Prosternum usually feebly margined at tip but somewhat variable. Last ventral with two setæ each side in both sexes. In the male the two middle setæ are much more widely separated from each other than from the lateral setæ while in the female the four are nearly equally spaced.

Length, 10-12.5 mm.

Three males and three females from the foot-hill cañons of the Sierras near Pomona; March, April, September.

*P. inermis* belongs to the first section of the genus (elytra without dorsal puncture), and is sufficiently distinct from all described forms to warrant its description in advance of a general revision of the genus. The obtuse hind angles of the thorax with the sides not sinuate before them is a rare character in the genus.

#### 8. *Cœlambus pedalis*, sp. nov.

Resembles *C. medialis* but is slightly larger, a little more densely punctate, and with the elytra more completely and uniformly infusate. Posterior femora with numerous coarse punctures which outwardly tend to coalesce into oblique rugæ. The claws of the front tarsi are similar in the male, but in this sex the middle tibiæ are quite strongly sinuate on the inner side toward the apex.

Five examples from Pomona and Riverside have been seen by the writer.

In *C. medialis* the hind thighs bear the usual line of fine setigerous punctures only.

#### 9. *Cœlambus femoratus*, sp. nov.

Superficially almost exactly like *C. nubilus*, but differing from it and all others by all the femora being sinuate beneath near the knee, the margins strongly laminate at apex. The tibiæ are pedunculate at base, then suddenly widened and nearly parallel, instead of gradually widening from base to apex as in all other species examined.

A single male example collected by Mr. Wickham at Albuquerque, New Mexico, was sent to the writer long ago as *C. nubilus*.

The western species of *Cœlambus* described by Le Conte are very perplexing and unless better characters than those given in the brief descriptions can be found, the number will have to be reduced. In two species, *C. masculinus* and *C. unguicularis*, Crotch observed in his description that the claws of the front tarsi in the male



are dissimilar. In these two species the disparity is very striking; but Crotch apparently failed to note its existence in numerous others. In *C. medialis*, *C. fraternus*, *C. femoratus* and *C. nubilus*, and perhaps others, the anterior (inner) claw is slightly broader and more strongly bent toward the base than its fellow, but the claws do not differ materially in length. In *C. fraternus* the front tarsi are much more strongly dilated in the male than in the same sex of *C. medialis*. *C. lutescens* is certainly very close to *C. medialis*, and I have not discovered any reliable means of separation; my specimens, however, of *C. lutescens* are all female, and it is possible the male may show good characters for separation.

10. *Ochthebius wickhami*, sp. nov.

Piceo-testaceous, head darker, legs pale; surface shining, feebly bronzed. Sides of thorax arcuate to middle, where they are slightly sinuate and abruptly, deeply emarginate, the angle prominent and acute; transparent border wide behind the sinuation, exceedingly narrow in front, but extending to the apical angles; disk sparsely punctate; median line deep, entire; anterior discal foveæ wanting, posterior deep; lateral grooves as in *O. discretus*. Elytra broad, sides arcuate; striæ feebly impressed, punctures not closely placed; intervals much wider than the striæ on the disk, where they are nearly flat; toward the sides the striæ are deeper and closer, the intervals more convex.

Length, 2 mm.

Taken in numbers at Winslow, Arizona, by Mr. Wickham.

The form of the thorax is very much like that of *O. discretus*.\* *O. wickhami* comes under "4" of Horn's table, and should precede *O. rectus* under the caption—sides of thorax arcuate in front, transparent border narrowly attaining the front angles. The author dedicates the species with much pleasure to its discoverer.

---

\* See Horn's figure, Trans. Am. Ent. Soc., Vol. XVII, Pl. II.

II. *Ochthebius costipennis*, sp. nov.

Piceous, not at all bronzed, legs brown. Outline and sculpture of thorax nearly as in *O. rectus*, except that the discal foveæ are connected by an impressed line. Elytra deeply striate, the punctures large and closely placed, intervals very narrow and strongly costiform throughout.

Length, slightly more than 1 mm.

One example taken by the author at Ventura.

No fuller description of this singular specimen is needed, as it agrees in all respects except size and sculpture, as noted, with *O. rectus*. I would not have ventured giving this specimen a name, relying solely on my own judgment, as it seemed possibly an aberrant *O. rectus*; but Mr. Frederick Blanchard, to whom the specimen was sent, returned it as "n. sp. near *O. discretus*;" and Mr. Liebeck, after carefully comparing it with Dr. Horn's material, wrote "There is nothing like it in the collection." I have seen scores of *O. rectus*, but have never seen a specimen approaching this one in size or character of elytral intervals. It should follow *O. rectus* in a cabinet arrangement.

12. *Tropisternus salsamentus*, sp. nov.

This name is proposed for a species related to *T. sublævis* and *T. californicus*. Though probably nearer to the former, it can best be described by a comparison with the more familiar *T. californicus*.

The form is a little more elongate than in *T. californicus*, punctuation more noticeable, but not dimming the surface lustre which is more or less greenish in certain lights; legs entirely yellow except a piceous pubescent area at base, and the middle and hind tarsi, which are brown. The finely punctate pubescent area occupies in *T. salsamentus* about the basal fourth of the hind femora, in *T. sublævis* the basal third, and in *T. californicus* is confined to the extreme base. The femora are more slender, more nearly parallel, and more sparsely punctate than in *T. californicus*. The mentum is not obviously more densely punctate in the male, and the tooth of the inner claw of the middle and hind tarsi (male) is usually blunt at tip instead of acute as in *T. californicus*. The outer claw of

same tarsi in the male is more or less angulate near the middle in *T. salsamentus*, never so in *T. californicus*. The raised line or crest at the tip of the last ventral segment is stronger in *T. salsamentus* and projects beyond the margin of the segment; less developed and terminating a little before the apex in *T. californicus*.

Specimens have been sent to Mr. Blanchard, who has kindly compared them with *T. sublaevis*. He reports that the latter species differs in being larger, broader behind, sides of thorax not sinuate posteriorly, mesosternal keel less strongly punctured, and the apical process of the last ventral less developed.

All the specimens seen (about a dozen) have been taken in April and July by Dr. Fényes and the writer from a small salt lake just back of the ocean beach at Redondo. This lake is much saltier than the ocean itself.

### 13. *Berosus salinus*, sp. nov.

Very similar to *B. infuscatus* in form and size; surface highly polished throughout in both sexes. Head black, with violaceous, green, and golden reflections; punctures not dense, vertical carina nearly obsolete. Thorax yellowish testaceous, nearly twice as wide as long; sides nearly straight and slightly convergent in front; angles rounded; punctures sparse, finer and coarser intermixed, a little closer at the sides. Elytra greenish testaceous, finely striate, striæ punctate; third interval with a confused line, fifth with a regular series of punctures; the others nearly or quite impunctate; sutural tips very slightly truncato-emarginate. Beneath piceous; abdomen not cristate; fifth ventral with a broad lobe at the bottom of the emargination; mesosternal crest acuminate; legs entirely pale, femora finely punctate and pubescent in rather less than basal half.

Length, 3.7-4.5 mm.

Redondo and Santa Barbara, in salt lake or pools.

The maculation is almost entirely absent, there being only a small divided thoracic spot, and a small diffuse shade on the second and third intervals of the elytra at one-third from the apex. The males have the first two joints of the front tarsi rather narrowly dilated and papillose beneath. By Horn's table *B. salinus* would be

associated with *B. styliferus*, but is remarkably distinct from any in our fauna by the peculiar elytral sculpture.

### Chætarthria.

Three species of this genus are more or less plentiful at almost any time of the year, by washing the sandy margins of streams throughout Southern California,—viz.: a pale form, *C. pallida* Lec.; a slightly larger, piceous black species with pale legs and antennæ, *C. nigrella* Lec.; and a smaller black species for which the name *C. minor* is now proposed. It may always be distinguished from *C. nigrella* by its distinctly smaller size (1.2–1.5 mm. for *C. minor* and 1.6–2.2 mm. for *C. nigrella*) and is moreover generally darker in color throughout, this being especially noticeable of the legs and antennæ. The form is also a little less broad in *C. minor*, the third joint from the last of the antennæ is smaller and more transverse, and the antennal club is less elongate.

A somewhat careful examination of our four species shows that well defined specific characters are not easy to find. It appears not to have been noticed that there is a sexual difference in the form of the front tibiæ. In the females the tibiæ are gradually broader to tip, while in the males there is a distinct angulation on the inner margin, beyond which the sides are parallel. In the eastern *C. atra* the angulation is near the knee, while in the Californian *C. nigrella* it is at or even a little beyond the middle; otherwise these two species are very close. In *C. pallida* the angle is less marked than in the two preceding, but its position is nearly as in *C. nigrella*. In *C. minor* the angle is not very obvious and is situated at about the basal third.

**14. *Philhydrus conjunctus*, sp. nov.**

Elongate-oval, less convex than *P. carinatus*, piceous black, the sides of the thorax paler. Punctuation as in *P. carinatus*, but with the lines of coarser punctures on the elytra less well defined. Prosternum not carinate, mesosternal lamina well developed, the lower edge horizontal, the anterior edge a little sinuate and slightly oblique, the free angle mucronate. Abdominal apex not emarginate. Thighs piceous, tibiae and tarsi paler. Claws of anterior tarsus in the male unequally toothed, the tooth of the front or inner claw large, lobe-like, and slightly everted; that of the other claw smaller but strong and subrectangular. Claws of middle and hind tarsi similar, strongly angulate at base.

Length, 4-4.5 mm.

Described from two males and one female taken at Lake Tahoe.

*P. conjunctus* much resembles *P. carinatus*, but is smaller and less convex, and by the palpal characters belongs to a different section of the genus. The claws are toothed somewhat after the manner of *P. hamiltoni*, but this latter species and its allies differ in color and in having the claws of the posterior tarsi sinuate in the male.

No mention seems to have been made of the fact that several species in this genus show a very small emargination at the apex of the last ventral segment, which is, so far as the writer has examined, very constant and entirely independent of sex. This emargination is present in *P. carinatus*, *P. cinctus*, *P. californicus*, *P. ochraceus*, and *P. nebulosus* (in the last two, much alike and deeper than in the others), while *P. perplexus*, *P. diffusus*, *P. hamiltoni* and *P. reflexipennis* do not show a trace of it. Of *P. fucatus*, *P. cuspidatus*, and *P. consors* nothing can be said, as no specimens are at hand.

The small, pale species, *P. ochraceus* and *P. nebulosus*, agree with the larger species of the same color, *P. diffusus*, etc., in the form of the posterior claws in the



male, a character seemingly of more importance in this genus than the carination of the prosternum and the form of the mesosternal lamina; they should therefore be placed together instead of at opposite ends of the genus as at present.

### 15. *Creniphilus elegans*, sp. nov.

Oblong-oval, slightly broader behind, not very convex, black, shining, without aeneous lustre, sides of thorax narrowly pale, elytra with a rather abrupt pale border which is wider behind. Punctuation distinct, but fine and rather sparse, a little stronger on the elytra. Antennæ seven-jointed, testaceous; club piceous: Femora piceous; trochanters, knees, tibiæ, and tarsi testaceous. Hind femora almost impunctate and without trace of strigosity; all the tarsi subequal in length to their respective tibiæ; claws slender, simple. Prosternum carinate, the carina obtusely prominent in front; mesosternal keel long, strongly protuberant. Abdomen densely punctate, subopaque.

Length, 2-2.8 mm.

Described from five examples taken by the writer in a salt lake at Redondo. Specimens collected by Mr. Hubbard in the salt springs at Salton, on the Colorado Desert, have recently been seen.

The difficulty of counting the intermediate joints of the antennæ in this genus is well known; nevertheless I am confident of the correctness of the statement above made; and if this is so, *C. elegans* agrees in antennal structure with *C. degener* only; and may, it seems, best precede that species in Horn's table. There is no possibility of confusing it with any other species of *Creniphilus* occurring in the same region. Of these, *C. infuscatus* has the hind femora punctate and pubescent; *C. rufiventris* has both pro- and mesosternum simple; and *C. subcupreus* is shorter, more convex, more coarsely punctate, and with shorter front tarsi.

In *C. subcupreus* the front claw joint is stouter and toothed beneath in the male, and apparently more conspicuously so in Californian than in eastern specimens.



**16. *Agathidium virile*, sp. nov.**

Moderately contractile, piceous black, apex of elytra and margins of thorax a little paler, legs brown. Head finely, sparsely punctate. Prothorax less than twice as wide as long, hind angles broadly rounded, front angles less broadly rounded, surface still more finely, obsoletely punctulate. Elytra longer than wide, distinctly but rather sparsely punctate, sutural stria deeply impressed except toward the base; side margin when viewed in profile subangularly rounded near the base. Metasternum alutaceous, obsoletely punctulate, without oblique elevated ridge. Mesosternum continuous with the metasternum, carinate in front. There is the usual difference in the tarsi of the sexes, and in addition the posterior femora are broader and obtusely angulate near the apex in the males, and in the same sex there is near the middle of the front margin of the metasternum a transversely oval fovea bearing a tuft of erect hairs.

Length (extended), 3 mm.

Riverside, Pomona and Pasadena.

This species may at once be known by the sexual characters of the male. It should be placed between *A. californicus* and *A. revolvens* in Horn's table.

**17. *Lomechusa angusta*, sp. nov.**

Red-brown, parallel, much smaller and narrower than either *L. cava* or *L. montana*. Upper surface, except the abdomen, minutely reticulate and sparsely, finely punctate, the head and prothorax somewhat dull, the elytra more shining; abdomen both above and beneath polished and nearly impunctate. Pubescence very fine, short and sparse; represented on the abdomen by longer erect setæ, which on the dorsal surface are arranged in a more or less regular transverse line on each segment, but are longer and more numerous beneath. Front concave; eyes situated at more than their own length from the hind margin of the head. Antennæ a little longer than half the body, the intermediate joints about twice as long as wide; terminal joint just visibly longer than the two preceding united. Prothorax nearly twice as wide as long, broadly emarginate in front throughout the width of the head, the front angles broadly rounding into the sides, which are feebly sinuate behind the middle, and thence slightly divergent to the hind angles, the latter nearly right and very narrowly rounded; base rather strongly arcuate at middle, sinuate laterally, side margins broadly reflexed. Elytra slightly wider than the prothorax, proportioned as in our other species. Abdomen and legs also as usual in the genus.

Length, 3.25 mm.

Pasadena and Redondo.

One example from each of the above localities is all that has been seen. The Pasadena specimen was taken with ants in a large oak gall, in February, by Dr. Fényes; and the Redondo specimen was picked up by Mr. Daggett on the beach in early spring. As indicated in the description, *L. angusta* is readily distinguished from both *L. cava* and *L. montana* by its small size and narrow form. In both the former also the last antennal joint is slightly but visibly shorter than the two preceding united, while in *L. angusta* it is slightly longer than the two preceding. *L. cava* and *L. montana* resemble each other much more closely, the size and form being nearly the same. In *L. cava* the sides of the prothorax are distinctly sinuate, the width being as great in front of the sinuation as it is posteriorly; while in *L. montana* the sides are not evidently sinuate, but diverge to a point just before the hind angles. In *L. angusta* the prothorax is widest at the hind angles. The intermediate joints of the antennæ in *L. cava* are decidedly more elongate than in *L. montana* or *L. angusta*.

#### 18. *Heterothops carbonatus*, sp. nov.

This species agrees tolerably well with Horn's description of *H. fumigatus*, except in the following particulars: Average size a little smaller; color deep black throughout except the legs, which may be more or less piceous; head more broadly oval, eyes much larger, distant about half their length from the nuchal constriction; last ventral of male less deeply emarginate.

Length, 3-3.5 mm.

Common along the margins of streams and in the vegetable mould which accumulates in damp situations throughout maritime Southern California.

While the color seems to be a perfectly reliable means of separation, the much larger eyes will in any event at once distinguish *H. carbonatus* from *H. fumigatus*; in

the latter species the eyes are distant very nearly their full length from the nuchal constriction.

*H. pusio* is much smaller, brownish in color, with a differently punctate head, and differently proportioned antennal joints.

*H. occidentis* is also much smaller, very slender, entirely testaceous, with exceedingly small eyes, and the last ventral of the male deeply cleft.

### 19. *Baptolinus punctiventris*, sp. nov.

Slender, rufous, elytra blackish, the basal margin, or at least the tips of the humeral umbones, less dark. Head, exclusive of mandibles, quadrate; hind angles broadly rounded, surface feebly alutaceous, excessively minutely, sparsely punctulate, with setigerous punctures as follows: a puncture over the insertion of the antennæ, a larger one above the eye, two smaller ones on the front, and a series of about eight at the hind angles and along the posterior margin; beneath with a few moderately fine punctures. Antennæ not reaching the middle of the thorax, proportioned as usual. Thorax as wide as, or very slightly wider than the head, parallel, longer than wide, angles rounded; dorsal series of two punctures, one near the middle, the other near the front margin and toward the side; otherwise impunctate except for the usual marginal setigerous punctures. Elytra barely equal in length to the thorax, glabrous except for the marginal setæ, and a series of about five shorter setæ in the sutural stria; surface very finely, indistinctly punctulate. Abdomen rather sparsely, but unusually coarsely muricately punctate, especially beneath. Last ventral of male with a broad and shallow but distinct triangular emargination.

Length, 5-6 mm.

Five specimens from Lake Tahoe and Vancouver Island. Mr. Schwarz writes that he has taken *B. punctiventris* in the mountains of Utah. It is rather common under bark, at Lake Tahoe.

### 20. *Actobius formosus*, sp. nov.

Form slender, piceous black; prothorax, legs, apical margin of elytra, posterior margin of ventral segments, and basal three (sometimes two) joints of antennæ, yellow. Antennæ slender, about as long as the head

and prothorax, tenth joint nearly square, preceding scarcely as broad as long. Head slightly longer than wide, sides behind the eyes nearly parallel, surface coarsely punctate, with a median smooth space. Prothorax one-half longer than wide, scarcely as wide as the head, distinctly narrowed behind; sides more sparsely and a little less coarsely punctate than the head. Elytra wider than the prothorax, not or but very little longer, longer than wide, usually with distinct greenish lustre; finely, sparsely punctate. Abdomen finely, moderately, densely punctate, moderately shining, beneath a little more strongly punctate than above.

In the male the front tarsi are rather strongly dilated, the last ventral with a small triangular emargination, which is broader than deep, and a triangular impression.

In the female the front tarsi are moderately dilated, the last ventral entire.

Length, 3.5-6.5 mm.

Pasadena and Pomona. Found along the margins of streams; March to October.

*A. formosus* is allied to *A. pæderoides* and *A. elegantulus*, but is larger than either and of a different color, which remains constant in the ten examples studied. The abdominal punctuation is much closer than in *A. elegantulus*, but less dense than in *A. pæderoides*. It differs from the latter species also in its longer thorax, which is more narrowed behind.

### Orus.

The species of this genus inhabit the true Pacific Coast region of California, and if secondary sexual characters are reliable, there are several species to add to the two described by Casey. The following three species are very close to *O. punctatus*, and while there are discoverable small differences in details, these are difficult to express in description, and however laboriously described it is practically certain that the collector would find it quite impossible to identify his specimens without reference to the male sexual characters.

21. *Orus parallelus* Casey.

The species is described from the female and is unknown to the writer. The thorax is described as being nearly as long as the elytra and rather densely punctate; characters which separate it from the following new forms, all of which agree nearly with *O. punctatus* in these particulars.

22. *Orus punctatus* Casey.

*Male*.:—Posterior margin of fifth ventral with a median lobe which varies somewhat in development, the lobe and posterior half of the segment distinctly longitudinally impressed; sixth segment broadly, deeply emarginate, the sides of the emargination concave, the apex rather narrowly rounded; base of segment in front of the emargination rather feebly impressed, the impression more or less covered by the lobe of the fifth segment.

23. *Orus fraternus*, sp. nov.

*Male*.:—Fifth ventral truncate or very broadly, feebly emarginate, feebly impressed; sixth ventral broadly, deeply emarginate, nearly as in *O. punctatus*, the segment feebly impressed before the emargination. As indicated by the series studied, the color is here never so black as in *O. punctatus*, being rather piceous or brownish piceous; but this difference is not so marked as to enable the positive separation of the females.

Seven examples from Pomona and San Bernardino.

24. *Orus montanus*, sp. nov.

*Male*.:—Fifth ventral broadly, feebly emarginate and very slightly impressed; sixth, triangularly emarginate, the emargination much less deep than in *O. punctatus*, the sides not at all concave, the apical angle a little less than right, and but slightly rounded. The size is a little larger and the form quite distinctly more robust than in any of the other species here mentioned, the thorax being scarcely at all narrower than the head, as is usual. The color is nearly black.

Two examples, a male and a female, were taken some years ago in the San Bernardino Mountains, at an elevation of 5,000 feet.



25. *Orus femoratus*, sp. nov.

*Male*.—Fifth ventral with a short, broadly rounded median lobe longitudinally impressed, the impression longitudinally divided at base by a short median raised line, and becoming posteriorly a spoon-shaped depression which is limited at the sides by acutely elevated folds or ridges; sixth ventral with a deep oblong-elliptical emargination which is widest at the middle of its depth and obtusely rounded in front, the base of the segment in front of the emargination with a smooth triangular impression having abrupt side margins. Hind femora much stouter than usual, and fully as broad as the anterior; lower edge acutely, sinuately compresso-carinate in basal half.

Represented by a single male collected by the author in Marin County.

In all the above species antennal joints two-four are distinctly, progressively shorter and not subequal, as is described of *O. parallelus*.

26. *Leptorus californicus*, sp. nov.

Slender, parallel, head and thorax rufotestaceous; elytra, metasternum, and abdomen, fuscous, the former with the suture narrowly, indefinitely paler, antennæ and legs pale; surface subalutaceous and dull except the thorax, which is feebly shining. Antennæ proportioned as usual. Head, as limited anteriorly by a line tangent to the anterior margin of the eyes, nearly square; sides behind the eyes parallel, twice the length of the eyes, hind angles narrowly rounded; base truncate, without median sinuation; surface densely, very finely punctate. Thorax narrower than the head, a little longer than wide, sides parallel, front angles strongly obtuse and rounded, base elliptically rounded; surface excessively finely, and more sparsely punctate than the head. Elytra about one-fourth longer than wide, distinctly wider and slightly longer than the thorax; punctuation fine and dense, but distinctly coarser than on the head. Abdomen a little narrower at base, surface very densely, finely punctate; sixth ventral in the male rather deeply, triangularly emarginate, the emargination as wide as deep and scarcely rounded at the bottom.

Length, 1.9-2.4 mm.

Described from two males taken at Palm Springs, on the western border of the Colorado Desert.

In general facies much like *L. texanus* Casey, but easily distinguishable by the male abdominal characters.



In *L. texanus*, which occurs at Yuma, the sixth ventral has a deeper emargination, which is rounded at bottom and with nearly parallel sides.

27. *Leptorus longipennis*, sp. nov.

Very slender, piceous, head and thorax dark rufous, legs and antennæ paler; surface more shining than in *L. californicus*. Sides of thorax slightly convergent toward the base; elytra fully one-third longer than the thorax. Punctuation much as in *L. californicus*, that of the head a little coarser above and distinctly so beneath.

Length, 2.2 mm.

Described from a single female specimen taken at Pomona.

This is the first species of the genus known to occur west of the mountains. It is at once separable from any species known to me by the unusual disparity in the length of the thorax and elytra, and by the decidedly stronger punctuation of the lower surface of the head.

28. *Scopæus armiger*, sp. nov.

In most features of size, form, and sculpture, and especially in the male sexual characters, this species is very similar to the eastern *S. dentiger*.\* It is a little more slender, the hind angles of the head less broadly rounded, the prothorax distinctly narrower than the elytra, the latter comparatively more elongate than in *S. dentiger*; the legs are brown. The male characters are substantially the same, with the following exceptions: The emargination of the sixth ventral is a little less broadly rounded at its apex, the inner sinuate margin of the posterior femora is a little more strongly serrate, and the hind tibiæ are not serrate internally.

The only specimen seen by me is a male taken at Pomona.

29. *Stilicus occiduus*, sp. nov.

Form and size the same as that of *S. angularis*, color more nearly piceous. Head above rather finely and densely punctate, not rugose; below, rufopiceous, polished, rather sparsely punctate; sides behind the eyes a

---

\* Trans. Am. Ent. Soc. 1880, p. 179.

little convergent, hind angles rounded, base truncate. Thorax densely punctate, a narrow, smooth, slightly elevated median line. Elytra more finely and sparsely punctate, tips narrowly pale; abdomen very finely, moderately densely punctate.

Length, 4 mm.

Described from one male and two females taken at Los Angeles and in Kern County.

*S. occiduus* is allied to *S. quadriceps* and *S. opaculus* by the punctuation of the upper surface of the head; but it is at once distinguished from these two species by the sparse punctuation of the lower surface, which is nearly as in *S. angularis*. In the male the fifth ventral segment is toothed at the middle of the hind margin, and the sixth is triangularly emarginate.

### 30. *Pseudopsis detrita*, sp. nov.

Piceous brown, legs and antennæ paler, surface dull, sparsely pubescent. Head three-fourths as wide as the prothorax, very finely punctulate, scarcely evidently strigose; eyes moderate, tempora subequal in length to the eyes. Antennæ shorter than the head and prothorax; intermediate joints submoniliform, outer joints gradually more transverse, the terminal nearly as long as the two preceding united. Prothorax transverse, a little narrower than the elytra, widest just behind the middle, sides nearly evenly, moderately strongly rounded, not sinuate before the angles, which are obtuse, the anterior less distinct; apex truncate; base arcuate, a little sinuate within the angles; surface finely not closely punctate, the disk with four obtuse and slightly elevated longitudinal lines; the two intermediate obsolete toward the apex. Elytra scarcely as wide as long, a little wider behind, sparsely but more distinctly punctate than the thorax; each with two vague discal costæ and a more distinct submarginal one. Abdomen more shining, distinctly punctate, the margin widely reflexed; first four dorsal segments equal, and each with a broadly, anteriorly arcuate impressed line from side to side; fifth segment as long as the two preceding united. Ventral segments moderately closely and distinctly punctate; the sixth in the male with a broad and deep subrectangular emargination, the inner angles broadly rounded.

Length, 2-2.2 mm.

A rather common species in vegetable detritus along the streams of the western slopes of the southern

Sierras, occurring with *P. obliterata* and *P. minuta*. It is rather close to *P. obliterata*, but the latter species is on the average a trifle larger, the head and thorax are distinctly longitudinally strigose, the tempora are longer, the antennæ a little stouter, and the impressed lines of the dorsal surface of the abdomen are more strongly arcuate. In *P. obliterata* the thorax is almost always noticeably paler than the head and elytra, but never distinctly so in *P. detrita* in my experience.

### 31. *Pseudopsis minuta*, sp. nov.

Yellowish brown or darker, head piceous, antennæ, legs, and tip of abdomen paler. Punctuation very fine anteriorly, more distinct on the abdomen; pubescence short, sparse, and indistinct. Head a little narrower than the prothorax, with three acute and entire longitudinal carinæ, one median, the others lateral; tempora about equal to the diameter of the eye. Prothorax a little narrower than the elytra, transverse, sides parallel, feebly arcuate, and at the middle indistinctly subangulate; disk with four equidistant and nearly straight longitudinal carinæ. Elytra as long as wide, each acutely tricostate; the two inner costæ curved inwardly at the apex. First four dorsal segments of the abdomen strongly arcuately impressed at base, their posterior margins bearing a series of widely spaced stout scales.

Length, 1.7-1.9 mm.

Found in the cañons of the southern Sierras.

### 32. *Bledius deceptivus*, sp. nov.

Rather robust, piceous black, elytra brownish black, legs and antennæ, except the outer joints, rufotestaceous. Head, apical portion of prothorax, and abdomen, finely reticulate; surface shining throughout, except the head, which is but feebly so; pubescence pale, short and sparse, longer on the abdomen, especially beneath. Head, including the eyes, slightly narrower than the prothorax; antennal prominences moderate, polished; epistomal tubercles nearly obsolete; frontal suture arcuate, sharply defined; vertex evenly convex, the median fovea small; surface behind the frontal suture rather coarsely but feebly punctate, the punctures lacking along the median line, otherwise separated by about their own diameters. Second joint of antennæ a little longer than the third, joints six to ten gradually wider and slightly transverse. Prothorax a little wider than long, sides parallel and nearly straight in slightly more than apical half, then broadly rounded and convergent to base, the basal angles rounde

and scarcely defined; the surface rather coarsely, moderately, deeply punctate, punctures a little closer at the sides, but separated on an average by their own diameters; impressed line not very deep but sharply defined throughout. Elytra a little wider, and at the suture about one-fifth longer than the prothorax, slightly wider posteriorly; punctures finer than on the prothorax, separated by about their own diameters. Abdomen above finely, sparsely punctate, beneath more closely, less finely punctate. Prosternum finely, longitudinally carinate before the coxæ; episterna narrowed in front and nearly flat. No sexual characters noted.

Length, 4.8 mm.

One example from Kern County.

The thorax is acutely margined, the front coxal cavities short and closed, thus making it a member of the *semiferrugineus* group of Le Conte. It is evidently near *B. opacifrons* Lec., but the distinctly, coarsely punctate head and other differences in detail mark it as quite surely distinct. It is a little smaller, but superficially much like *B. armatus*, which occurs in the same region; the latter, however, has the prosternal sutures obsolete, the prosternum not carinate in front, the sides of the prothorax parallel for a longer distance, and the punctuation sparser throughout.

### 33. *Bledius relictus*, sp. nov.

The description of the preceding species applies to this very nearly except in the following particulars: The elytra are distinctly shorter, being but slightly longer than the prothorax, and are rufocastaneous, the base and suture more or less blackish. The antennæ have the outer joints more decidedly transverse, and the prothoracic episterna are here more narrowed anteriorly, the width at the front angles being less than one-half that at the coxal fissures, while in *B. deceptivus* it is fully one-half as great.

Length, 4.7-5 mm.

Two examples taken at Pomona in April.

### 34. *Bledius clarus*, sp. nov.

Slender, convex, black; prothorax, elytra, legs and antennæ rufotestaceous; the abdomen sometimes slightly rufescent beneath. Head finely reticulate, moderately shining; thorax, elytra and abdomen polished; the last-named very feebly, the thorax more evidently and more finely reticulate.

Punctuation throughout very fine and sparse, the punctures of the surface of the abdomen being especially distant on the dorsal surface but more numerous beneath. Head a little narrower than the prothorax; front evenly convex, transverse suture very fine but distinct, occipital fovea wanting. Antennæ moderately strongly incrassate, the second joint equal in length to the two following, or very nearly so, the sixth to the tenth transverse, eleventh about as long as the two preceding united. Prothorax wider than long, sides nearly straight and parallel in about the apical two-thirds, then rather abruptly rounded and slightly sinuately convergent to the basal angles, which are obtuse but moderately well defined; front angles rather broadly rounded; dorsal line fine. Elytra but little wider and not much longer than the prothorax. Prosternal episterna parallel, almost perfectly flat. Mentum more or less impressed at base.

Length, 2.5-2.8 mm.

Five examples, collected at Pomona, Riverside, and near San Bernardino, have been seen by the writer.

*B. clarus* is a very pretty species which shows little variation.

### 35. *Bledius rusticus*, sp. nov.

Moderately slender, piceous black, prothorax and elytra brownish yellow, legs and base of antennæ testaceous. Head and prothorax densely reticulate granulose, not shining; elytra polished; abdomen finely reticulate, moderately shining; punctuation rather sparse and fine throughout. Head slightly narrower than the prothorax; antennal tubercles moderate; front convex, transverse line faint, not at all impressed; no occipital fovea. Antennæ short, rather strongly widened externally, second joint nearly as long as the two following, the sixth to the tenth strongly transverse. Prothorax a little wider than long, sides feebly rounded and nearly parallel in basal two-thirds, then broadly rounded and sinuately convergent to the basal angles, which are obtuse and rounded; dorsal line fine. Elytra very slightly wider at base than the prothorax and about one-fourth longer, a little wider behind; abdomen gradually wider posteriorly. Prosternum not carinate; mentum broadly concave.

Length, 2.4-3.7 mm.

Three examples taken at Pomona.

The prosternal sutures are parallel with the margin, the hypomera parallel and slightly concave; coxal fissures open. It is therefore to be referred to the *annularis* group, and should probably stand near *B. luteipennis* Lec. The latter is said to have the thorax not



wider than long, and the mentum with a small, deep impression near the hind margin.

### 36. *Eustilbus notabilis*, sp. nov.

Oval, moderately convex, very slightly attenuate behind; head and thorax black, the latter a little paler at the side margin, elytra rufous, under surface, legs and antennæ rufous or rufotestaceous. Upper surface polished and impunctate throughout, the usual lines of punctures on the elytra being scarcely visible. The surface of the elytra is very finely, transversely strigulose, more noticeably so toward the apex, where the surface lustre is slightly dulled. The sutural stria obsolete at about the basal third; eighth antennal joint slightly transverse; free edge of prosternal process bearing four or five setæ, which are as long as the width of the process; first joint of hind tarsi rather more than half the length of the second; spurs of hind tibiæ a little unequal, distinctly longer than the terminal fringe of spinules, but shorter than the basal joint of the tarsus. There are no sexual differences observable in the seven examples studied.

Length, 1-1.3 mm.

Specimens taken at Pomona.

This is a minute species, easily distinguishable by its color from any other in our fauna.

### 37. *Exochomus histrio*, sp. nov.

Moderately convex, shining, surface sparsely, very finely punctulate, and finely, feebly reticulate; black throughout except a short parallel humeral stripe and a subapical spot on each elytron, pale. Claws simple.

Length, 2.3-3 mm.

Four examples collected near Pomona, with which are associated for the present two entirely black specimens from Santa Barbara.

*E. histrio* is an exact counterpart of Casey's *E. californicus*, but the size is smaller (the series of eight specimens of the latter species in the writer's collection ranges from 3 to 4 mm. in length), and the claws are not toothed at base.

The fact that in some of our species of *Exochomus* the claws are toothed at base while in others they are quite



simple seems to have been entirely overlooked. This character enables us to more conveniently tabulate the species, and in fact is the only certain means of separating the two above mentioned.

The series with simple claws includes, besides *E. histrio*, *E. septentrionis* Weise, *E. hoegei* Gorh, and *E. ovoideus* Csy. Weise's name is used here provisionally for a large, strongly punctured species occurring in New England and adjacent states. It agrees with Weise's description in size and most nearly in habitat (*E. septentrionis* was described from the Hudson Bay Region), and there can be little doubt, it seems, as to the correctness of the reference. No mention is made of *E. septentrionis* in Casey's recent Revision of the Coccinellidæ; this is doubtless an oversight. *E. desertorum* Csy. must be considered a synonym of *E. ovoideus* Csy.

### 38. *Smilia reversa*, sp. nov.

Broadly oval; black, elytra brownish. Thorax polished throughout; punctuation closer and more distinct than in any of our other species; impressed line of front angles distinct, joining the basal marginal line in a continuous curve along the side margin, which is distinctly explanate. Elytra very finely, sparsely punctulate; sutural line visible behind the middle, but scarcely impressed.

Length, 1.25 mm.

Five examples, beaten from conifers, at Lake Tahoe and in the San Bernardino Mountains, at an elevation of from 5,000 to 7,000 feet.

This is our largest species and is conspicuous from the fact that the thoracic punctuation, while somewhat variable, is in general closer and coarser than that of the elytra, whereas it is the reverse of this in all our other species (I do not know *S. coccidivora*). The joining of the apical and basal marginal lines of the

prothorax has not been observed by me in any other species. *S. reversa* seems most nearly allied to *S. marginata*.

### 39. *Hyperaspis excelsa*, sp. nov.

Broadly oval, strongly convex; shining, rather finely punctate; black, elytra with humeral marginal stripe joining a large transverse discal spot, red; a smaller apical spot sometimes present. Male with head, sides of thorax narrowly, front margin still more narrowly, and front legs, yellow.

Length, 3.75 mm.

Southern California (Pomona, Pasadena, San Bernardino Mountains); rare.

Marked much like *H. teniata*, but very much larger and with somewhat different sexual color marks in the male.

### 40. *Hyperaspis spiculinota*, sp. nov.

Elongate-oval, subdepressed; shining, except the head, which is finely reticulate and somewhat dull; black, with the sides of the thorax, median and apical marginal, and elongate discal spot, yellow. Legs dull yellowish, the femora more or less piceous. Head yellow in the male.

Pasadena and Pomona; not common.

The markings of this species are of the *H. undulata* type, but the humeral spot is never present, and the discal spot is very narrow, elongate, subtriangular, pointed behind, and truncate in front. The marginal spots are rather narrowly separated but are not connected in any specimens which I have seen. The oblong, depressed form, and shape of discal spot are characteristic.

There are in my cabinet three or four other probably undescribed species from Southern California, but they are not sufficiently strongly characterized, or not present in sufficient numbers, to warrant their description in advance of a monographic treatment of the genus.

**41. *Scymnus tædatus*, sp. nov.**

Oval, thoracic and elytral margins not continuous; dark brown; head, thorax and tip of abdomen yellowish; the thorax with faint piceous clouds at the middle and before the scutellum; elytra each with a small spot before the middle and nearer the suture, another a little larger and posterior to it, and more or less connected with an indefinite apical space, pale. Head evidently punctate; thorax lightly but distinctly punctate at middle, more finely laterally; sides nearly straight in basal two-thirds. Elytra less finely but not closely punctate. Prosternal elevated lines distinct; mesocoxal arcs joining the lateral suture a little before the middle; metacoxal arc incomplete, joining the first ventral suture. Abdomen finely punctate, and with six distinct segments.

Length, nearly 1.5 mm.

Found in the Sierras of Southern California. Several specimens beaten from *Pinus ponderosa*.

By the metacoxal lines this species belongs to Dr. Horn's Group A, and may follow *S. bigemmeus*. There are no obvious sexual differences.

**42. *Scymnus megacephalus*, sp. nov.**

Oblong, obtusely rounded before and behind, moderately convex, uniformly rufotestaceous throughout. Pubescence rather short, flavocinereous, not very conspicuous but slightly holosericeous in aspect owing to the arrangement of the hairs. Head very large, the eyes almost entirely exposed and separated on the front by distinctly more than twice their own width; punctuation rather fine, the punctures separated by from one to two times their own diameters. Prothorax less than twice as wide as long, the sides evenly, moderately arcuate, the apex evidently though not very much narrower than the base; surface punctate similarly to the head, the disposition and size of the punctures nearly uniform throughout; base not sinuate each side of the middle, the marginal line strictly basal before the scutellum. Elytra oblong, slightly wider than the prothorax, the sides very feebly arcuate, and parallel to a point far behind the middle; punctuation somewhat sparser than on the head and prothorax. Scutellum equilateral, larger than usual. Prosternal ridges strong and entire; mesosternum closely, rather coarsely punctate; abdomen less coarsely and closely punctate than the metasternum. Postcoxal arcs of the metasternum incomplete externally, though nearly reaching the suture; those of the first ventral obliterated externally, the lines reaching about to the apical fourth of the segment and scarcely visibly recurved, the plates punctate like the rest of the segment. Femora stout; claws

cleft, the interior portion not very much shorter than the outer in the front pair but more evidently so in the others. Abdomen with six distinct segments.

Length, 2.25 mm.

Described from a single example taken at Pasadena, November 2, 1899.

By the tables given by Horn and Casey, *S. megacephalus* falls near *S. difficilis*, *S. phelpsi*, and *S. nebulosus*, from all of which the more parallel form and large head will at once distinguish it. The sex of the type is not determinable with certainty, but it is probably a female.

#### 43. *Scymnus mimus*, sp. nov.

Resembles *S. pallens* exactly, and is only separable with certainty by the male characters. These consist of a subtriangular impunctate area surrounded by denser suberect pubescence at the middle of the posterior margin of the first ventral, and the strongly impressed sixth ventral. In *S. pallens* the first ventral of the male is unmodified, and the last is but feebly impresso-emarginate. In *S. mimus* the abdominal punctuation is distinctly finer and sparser, and there are but two pale segments, while in *S. pallens* the last three are usually pale.

There are two males in my collection, from Southern California (Riverside and Pomona); the female has not yet been identified.

#### 44. *Scymnus dentipes*, sp. nov.

Broadly oval; black; head, sides of thorax in front, and rather narrow apical space on the elytra, reddish yellow. Head and thorax rather sparsely, elytra more coarsely and closely punctate. Beneath black, abdomen paler at apex, legs dull reddish yellow, the femora often more or less clouded, especially towards the base. In the male the first ventral is distinctly longitudinally impressed at middle, both the impression and its margin punctulate and bearing numerous fine erect hairs; the last ventral is moderately impresso-emarginate, and the middle thighs are obtusely dentate beneath, near the knees.

In size, and in fact in all respects except the secondary sexual characters, this species is scarcely distinguishable from *S. fraternus*, and is possibly confused

with it in collections. For its discovery we are indebted to Mr. Frederick Blanchard of Tyngsboro, Mass., who finds it not uncommon on pitch pines in his vicinity.

#### 45. *Languria californica*, sp. nov.

Elongate, parallel, greenish black, head and prothorax red. Antennal club five-jointed, head moderately punctate. Prothorax longer than wide, sides feebly rounded, a little more strongly so in front, subsinuate before the hind angles; surface strongly, rather numerously punctate, basal striae short, distinct. Elytra rather deeply punctate striate, intervals somewhat convex, each with a more or less regular line of excessively fine and indistinct punctures bearing very fine short hairs. Beneath rather sparsely but strongly punctate throughout.

Length, 6.5 mm.

One example taken at Redondo, in April, on *Astragalus crotalaræ*.

*L. convexicollis*, the only other Californian species of the genus, is much larger, with more convex thorax, six-jointed antennal club, and generally feebler sculpture. *L. californica* may be placed between *L. convexicollis* and *L. læta*.

#### 46. *Hister simplicipes*, sp. nov.

Not very robust, black, polished; impunctate above, except along the basal margin of the thorax; elytra dark red at sides and in a subapical transverse line which is interrupted at suture. Thorax not fimbriate, internal lateral stria complete, external scarcely reaching the middle. Elytra with three entire dorsal striae; fourth short, apical; fifth represented by two or three small punctures near the apex; sutural stria extending in front of the middle; oblique humeral fine, subhumeral wanting. Propygidium moderately coarsely, densely punctate; pygidium densely punctate at base, nearly smooth at apex, which is strongly inflexed. Mesosternum moderately emarginate; anterior tibiae without spines or teeth except the apical, which is not very strongly bifid; hind tibiae bispinulose.

Length, 4 mm.

A single example in the writer's collection is from San Diego County.

*H. simplicipes* is less broad, less convex, and with less red than *H. sellatus*, which occurs in the same region. It is more like *H. militaris*, but very distinct from it and all our other species—except *H. lævipes*—by the non-denticulate outer margin of the front tibiæ. In *H. lævipes* the thoracic margin is fimbriate, and the sculpture different. The characters are such as to exclude *H. simplicipes* from any of the groups as limited in Dr. Horn's synopsis, but it seems best to place it following *H. lævipes*.

There is, in the writer's collection, a single example from San Bernardino County of what may prove to be a distinct species, but which is so close to the species above described, that further specimens are necessary to establish its status. It differs from *H. simplicipes* in being entirely black, with the fourth dorsal stria of the elytra interruptedly continued nearly to the base. There is a small group of well defined punctures between the lateral striæ of the thorax, and near the front angles, which is only feebly evident in *H. simplicipes*.

#### 47. *Saprinus liticolus*, sp. nov.

Oblong-oval, not very convex, black, shining, feebly but not very finely punctate. Supraorbital line obsolete. Thorax less than twice as wide as long, sides convergent and nearly straight to apical fourth; surface sparsely, rather finely punctate laterally, a line of coarser punctures along the base, disk with very minute and widely spaced punctures. Elytra about one-half longer than the thorax, rather strongly punctate in apical half, the punctures a little elongate and denser apically, but separated on the average by their own diameters. External subhumeral wanting; internal subhumeral either wanting or short and apical; oblique humeral distinct; four dorsals deeply impressed, punctate, subequal in length, and extending but little beyond the middle; fourth dorsal arched at base and joining the sutural, which does not quite reach the apex. Pygidia densely but not very coarsely punctate. Prosternum not strongly



convex, with numerous coarse, linear punctures, striæ strongly divergent in front, foveæ deep. Anterior tibiæ about octo-serrulato-spinose.

Length, 2.25-3 mm.

Not rare on the sea beach at Redondo.

*S. liticolus* belongs to Horn's Group VI, and would fall next to *S. vescus* in the tabular arrangement given; the latter species, however, differs in several details, notably the longer, unequal dorsal elytral striæ, and at the time of Horn's writing it was known only from the unique type, which was taken in Texas.

It is probable that *S. liticolus* is confined to the coast line. The only two species occurring in the same situations with which it could be confounded are *S. laridus* and *S. scissus*, both of which are smaller and have a narrower, more convex prosternum, which is scarcely punctate. *S. scissus* is, moreover, of broader form, and *S. laridus* is nearly equally punctate throughout.

#### 48. *Saprinus consobrinus*, sp. nov.

Closely related to *S. fraternus*, from which it differs in the following details: Head with a single irregular chevron. Prothorax with the sides slightly more convergent, the punctuation scarcely at all strigose at any point. Elytral punctuation very dense, not strigose; scutellar mirror very small, twice as wide as long; dorsal striæ not very distinct; internal humeral entire, but twice interrupted; external humeral short, distinct; oblique humeral very short, occupying the umbone, which is more evidently impunctate than in *S. fraternus*. Anterior tibiæ about 8-denticulate, the upper three teeth very small.

Length, 3 mm.

Specimens taken at Yuma.

*S. fraternus*, *S. consobrinus*, *S. laxatus*, *S. mancus* and *S. bigemmeus* form a natural section of Horn's Group III, distinguished by the evident sutural stria, and by having the elytra punctured more or less densely over the entire surface except a rather small scutellar mirror, which is well defined in all except *S. mancus*. *S. bigemmeus* is separated by the strongly aciculate strigose

elytra; it is the smallest of the group and has never been found far from the California seacoast. *S. frater-nus* and *S. consobrinus* have the sutural stria entire, while in *S. mancus* and *S. laxatus* it is indistinct or wanting behind the middle. *S. mancus* is a northern species; *S. laxatus* is larger and less densely sculptured, and is known only from Florida.

#### 49. *Acritus volitans*, sp. nov.

Broadly oval, moderately convex, rufopiceous, legs and antennæ a little paler. Head finely and sparsely, but rather strongly, punctulate. Second antennal joint twice as long as the third, the latter nearly as long as the fourth and fifth together, eighth a little wider than the preceding. Thorax strongly narrowed from the base, the sides broadly arcuate, punctuation fine and rather sparse, nearly uniform throughout; a transverse basal row of larger punctures extending to the middle of the elytra and curved forward in front of the scutellum. Elytra a little more coarsely punctate, the punctures feebly aciculate behind the middle, uniformly distributed, separated by from two to three times their own diameters; marginal stria of the inflexed flanks of the elytra fairly distinct, deeper toward the base; sutural and discal striæ entirely wanting. Pygidium finely, rather sparsely punctate. Prosternum about one-half longer than wide, with a few very feeble scattered punctures; the striæ equally divergent before and behind; meso-metasternal plate very finely, sparsely punctulate at middle, more coarsely at sides and posteriorly; metasternal suture obliterated, its position defined by a transverse row of elongate punctures somewhat similar to those on the prothorax; marginal stria of mesosternum rather broadly interrupted at middle. Anterior tibiæ but slightly broader than the middle.

Length, 1 mm.

Ten examples taken at Pomona and Pasadena from May to September. All occurred flying in early evening, or at electric lights.

By the characters given in Horn's table, this species should stand between *A. exiguus* and *A. floridæ*.

#### 50. *Stenelmis nubifer*, sp. nov.

Elongate, parallel, finely pubescent. Head and prothorax black without lustre, the apices of the front angles rufescent; elytra dull yellow with a broad diffuse median cloud; beneath fuscous or fuscoferruginous;

antennæ and legs testaceous, the femora sometimes darker. Head rather coarsely punctate; antennæ reaching almost to the hind angles of the prothorax. Prothorax about as long as wide, a little narrowed in front, sides moderately rounded just behind the middle, and also slightly so before the hind angles, which are a little acute; surface moderately closely granulate-punctate, disk with two small impressions at the middle of the base, a deeper longitudinal one each side extending to the middle, and an elongate median fovea. The base is slightly longitudinally channeled before the scutellum and between the small basal impressions, the channel is smooth at the bottom posteriorly. There is also in some examples an oblique impression extending from the lateral impressions inward and forward toward the median fovea. Elytra but little wider than the prothorax, nearly parallel, sides slightly sinuate; the rows of punctures feebly impressed; intervals nearly flat, the fourth a little prominent at the base, the sixth finely carinate nearly to the tip. Beneath coarsely, rather densely scabrous-punctate, except the abdomen, which is less roughly sculptured, especially posteriorly.

Length, 2-2.4 mm.

Described from nine examples taken in a small mountain stream near Pasadena.

This is the first species of the genus to be reported from the Pacific Coast; it is not closely related to any of the eastern species.

### 51. *Eurypogon confusus*, sp. nov.

Broader and more depressed than *E. niger*, brownish testaceous throughout, shining; pubescence yellowish, inclined backward on the elytra and forward on the thorax. Head sparsely, finely punctate. Antennæ slender, reaching slightly beyond the middle of the elytra, second joint a little more than half as long as the third, the latter nearly as long as the fourth; joints four to eleven subequal. Prothorax twice as wide as long, sides nearly parallel for a short distance at base, then rounded and rather strongly convergent to apical angles; punctuation fine, the punctures quite uniformly distributed and separated by from two to three times their own diameters; disk with two small impressions, about as distant from each other as from the side margin, and a somewhat vague impression extending from the anterior angles obliquely inward toward the discal impressions. Elytra a little wider than the thorax, sides parallel in basal three-fifths, surface faintly substriate, intervals confusedly punctate, the punctures scarcely distinguishable from those of the striæ.

Prosternum and sides of metasternum closely, not finely, punctate; abdomen less closely, middle of metasternum sparsely, more finely punctured.

Length, 4 mm.

Described from a single specimen taken at Pasadena by Dr. Fényes.

*E. confusus* is quite different from *E. californicus* (which is much like *E. niger* in general facies) because of the more transverse prothorax, which is more rounded on the sides, and the confused elytral punctuation.

In notes taken at the time of making comparison with the type of *E. californicus* in the Horn Collection, allusion was made to a difference in the form of the basal lobe of the prothorax, but the difference was not then described and cannot now be recalled.

## 52. *Horistonotus flavidus*, sp. nov.

Pale yellowish testaceous, sparsely fulvo-pubescent, shining; of the same form as *H. simplex*, but a little smaller and distinctly more slender. Head densely punctulate with numerous coarser punctures intermixed. Antennæ feebly serrate, very slightly longer than the head and prothorax. Prothorax subquadrate, hind angles not divergent, sides straight and nearly parallel for three-fourths their length; surface moderately convex, with the usual double punctuation, the coarser punctures rather fine and scarcely smaller posteriorly. Elytra but little wider than the prothorax; sides parallel and very feebly arcuate to apical two-fifths, apex elliptically rounded; striæ moderate, rather coarsely punctate at base, more finely toward the apex; intervals rather sparsely, confusedly punctulate. Sides of prothorax beneath coarsely punctate, the punctures separated on an average by little more than their own diameters; metasternum less coarsely, abdomen finely, sparsely punctate, the minute punctures present over the entire surface. Claws as in *H. simplex*.

Length, 6 mm.

Several specimens were received by Dr. Fényes from Palm Springs. The specimens were probably taken in June, at electric light.

This species may follow *H. simplex*, to which it is most closely related. It differs in its more slender form,

paler color, more shining surface, less widely separated eyes, and coarser punctures of the prosternal flanks. In all my examples of *H. simplex* the last joint of the maxillary palpi is externally sinuate toward the apex; this is not at all evident in *H. flavidus*.

### 53. *Agrilus gibbicollis*, sp. nov.

Robust, olivaceous bronze, feebly shining, sides of prothorax with whitish pubescence and efflorescence, elytra with entire pubescent vitta. Antennæ short, scarcely passing the upper margin of the eye when directed upwards, serrate from the fifth joint, seventh and following joints as wide as long. Head convex, front densely clothed with short yellowish brown hair, apparently not impressed. Prothorax nearly as long at middle as wide; sides rather broadly arcuate in front, posteriorly oblique and distinctly sinuate before the hind angles, which are a little prominent and slightly acute; disk strongly convex in front, gibbous when viewed in profile; surface punctate and concentrically strigose, more strongly so in front; hind angles moderately carinate; lateral margin sinuate when viewed from the side. Scutellum not carinate. Elytra a little sinuate at sides; the abdomen more strongly so, its sides visible from above behind the middle; tips rounded and serrulate. Prosternal lobe broadly rounded, subtruncate at middle; prosternal process nearly parallel, the visible apex truncate; sides of body beneath, and vertical portions of ventral segments with whitish pubescence and efflorescence. Pygidium without projecting carina. Claws moderately toothed at middle.

Length, 7 mm.

Described from a single female example taken by Dr. Fényes, near San Bernardino, in June. Another specimen seen in the Horn Collection was there placed with *A. blandus*. The gibbous prothorax with carinate hind angles, and non-carinate scutellum, easily distinguish *A. gibbicollis* from *A. blandus*, to which it bears only a superficial resemblance. By Horn's table, it would fall with *A. abductus* and *A. palmarcollis*, in both of which the scutellum is carinate and the thorax normally convex.



54. *Agrilus illectus*, sp. nov.

Moderately elongate, subcylindrical, cupreo-aeneous, dull above, more shining beneath; pubescence sparse and inconspicuous, a little closer in a faint subsutural vitta, and more markedly along the sides of the sterna and vertical portions of the ventral segments. Antennæ piceous, attaining the middle of the prothorax, serrate from the fifth joint, the outer joints as wide as or a little wider than long. Head broadly, strongly longitudinally impressed, punctate and strigose. Prothorax a little wider than long, sides feebly arcuate when viewed from above, sinuate when viewed laterally; hind angles right, distinctly carinate; surface quite strongly strigose except along the side margins; dorsal channel distinct but not strong, lateral impressions well marked. Scutellum indistinctly carinate. Elytra moderately sinuate; tips rounded, feebly or scarcely serrulate; disk without trace of costæ; surface imbricate. Prosternal lobe truncate in front, a faint sinus at the middle; intercoxal process broad, truncate at tip. Abdomen rather sparsely punctate, reticulate at sides. Pygidium with a fine carina which does not nearly reach the tip. Claws with a broad tooth.

Length, 5-7.5 mm.

Described from one male and three females taken at Pomona and Pasadena by sweeping late in June.

The sexual differences are feeble, consisting of a slight flattening of the first two ventral segments, and the longer, denser pubescence along the median line of the sterna in the male. *A. illectus* would fall near *A. impexus* by Horn's table, but it is not closely related to any species.

55. *Collops argutus*, sp. nov.

Upper surface shining throughout, clothed with erect black and shorter grayish hairs, the latter sparse and inconspicuous. Head black, labrum and epistoma yellow, antennæ black except the first joint and inner side of second joint, which are pale. Head distinctly but not coarsely punctate, the punctures separated on an average by their own diameters. Prothorax transversely oval, fully two-thirds wider than long, entirely rufous, very finely sparsely punctulate, the punctures a trifle closer at the sides. Elytra densely moderately closely punctate, rufous, each with two large black or bluish spots. The two basal spots involve the humeri, and are confluent anteriorly at the suture; the posterior spots reach the margin



but not the suture, and frequently join the basal spots. In one example the elytra are entirely dark except the middle of the lateral margin. Legs black except the front trochanters and the tips of the front coxæ. Abdomen black, at times rufous at the middle toward the base.

Length, 4.3-5 mm.

Many specimens have been seen from various localities in the State from San Diego to Sonoma County.

The coloration of the legs and antennæ is very constant in the series studied, and will probably prove a reliable means of separating the species from *C. 4-maculatus*, and others which are similarly colored above. The prothorax is here wider than in *C. 4-maculatus*, but less transverse than in *C. laticollis*, which has moreover a different style of elytral punctuation, and front legs and abdomen entirely pale.

*C. argutus* is one of a small number of species having the first antennal joint in the male elongate and distinctly excavate or sinuate on its inner face, as seen from above; and furthermore the claw-like appendage of the second joint in the male is here much longer than usual.

## 56. *Malachius inornatus*, sp. nov.

Rather robust, dark blue-green throughout; thorax somewhat shining, elytra dull; clothed sparsely with very short, recumbent pale pubescence. Head very finely punctulate, moderately shining, a longitudinal impression within the antennæ, and a more or less evident transverse impression, which is at times reduced to a shallow median fovea. Antennæ of the male moderately strongly serrate, reaching the middle of the elytra; those of the female feebly serrate and a little shorter. Prothorax transversely oval, broadly transversely impressed at base, finely, sparsely punctate, feebly alutaceous. Elytra subparallel (male), wider posteriorly (female), finely scabrous and punctulate, tip appendiculate in the male, broadly rounded in the female. Under surface and legs slightly darker in tint; last ventral of male deeply incised.

Length, 4-5 mm.

Four examples from Pomona.

The form of the elytral appendix is not easy to describe. The sutural angle is impressed and extended on a lower plane into an acute process; inferiorly the elytron is produced in the form of a broad, somewhat contorted plate which is deeply sinuate externally near the apex, the outer angle of the sinuation acute; parallel to and above the inner margin of the plate is an irregular process which is attached at base, rounded at tip, its upper margin bidentate near the base. The entire appendix is black without lustre, and is sparsely punctate and pubescent. The epistoma is more or less pale.

*M. inornatus* is nearest *M. auritus* and *M. thevenetii*, but in the former the thorax is red at sides, and it and the base of the elytra bear numerous fine, erect hairs in addition to the usual recumbent pubescence; while in *M. thevenetii* the male antennæ are pectinate, and in both the elytra are tipped with red in the female.

#### 57. *Malachius pristinus*, sp. nov.

Very small, slender, parallel, dull black, not shining, finely punctate; head very finely, thorax more distinctly alutaceous; pubescence short, pale, semierect, inclined forward on the prothorax, and outward on the elytra; mouth, under side of three or four basal joints of antennæ, and sutural angles of elytra, yellow; legs a little paler than the upper surface, knees of front and middle pair yellowish. Antennæ scarcely reaching the middle of the elytra, very feebly serrate, joints five to ten fully twice as long as wide. Head feebly impressed. Prothorax transverse, sides nearly parallel, all the angles rather broadly rounded; base and apex arcuate, the latter less strongly. Tips of elytra inflexed, each bearing on the vertical face and close to the suture a slender erect process which extends upward to the level of the elytra, and is curved forward at tip; the inferior plate is broad and reflexed at tip.

Length, 2.5 mm.

Found in the Sierras of Los Angeles County.

The small size, dull surface, and peculiar elytral appendix are peculiar. A single male only has been seen.

**58. *Malachius acutipennis*, sp. nov.**

Moderately slender; front margin of labrum and epistoma, and under side of two basal joints of antennæ, yellow; head black, prothorax entirely reddish yellow, or with a small median black spot extending from base to middle; elytra black with reddish bronze lustre, sutural angles reddish yellow. Antennæ strongly serrate and reaching the middle of the elytra in the male, a little shorter and much less strongly serrate in the female. Head and prothorax polished, impunctate; prothorax transversely oval. Elytra nearly parallel, feebly scabrous and punctulate, appendiculate, the sutural angle spiniform in the male; wider behind, the sutural angle broadly rounded in the female. Legs and lower surface, except prothorax, black, without well defined metallic lustre; hind tibiæ pale in the male.

Length, 4-5 mm.

Specimens taken at Pomona.

The form of the elytra and the appendix in the male is not very different from that in *M. spinipennis*, to which this species is evidently related; the color of the elytra, however, is bluish in *M. spinipennis*, the prothorax is broadly black at middle from base to apex, and the male antennæ are pectinate.

**59. *Malachius directus*, sp. nov.**

Not very slender; greenish black; prothorax with sides broadly yellow; apex of first, and entire second joint of antennæ, yellow on the under side. Head and prothorax polished, subimpunctate; elytra scabrous, less shining. Antennæ in the male reaching or slightly passing the middle of the elytra, rather strongly pectinate from the fifth joint; the third joint triangular, as wide as long; fourth with a shorter branch than those of the following joints. In the female the antennæ scarcely pass the humeri, and are moderately serrate, the widest joints a little longer than wide. Prothorax transversely oval. Elytra appendiculate in the male, the appendix consisting merely of a single, somewhat crumpled plate, which is but slightly inferior; the tip strongly rounded and more or less reflexed.

Length, 3.5 mm.

Found in the Sierras of Los Angeles and San Bernardino counties.

In the specimens which are to be regarded as types, the elytra are not tipped with yellow; other specimens

which are, however, quite surely the same thing, have the tips very narrowly yellow. These latter resemble *M. mixtus* quite closely, but this latter species has the elytra non-appendiculate in the male.

#### 60. *Malachius nigrinus*, sp. nov.

Slender, shining, black, without evident metallic lustre, the mouth, labrum, and epistoma, under side of the first three antennal joints, and a very narrow margin at the hind angles of the prothorax, yellow. Surface very finely, sparsely punctulate, with short sparse pale pubescence, and a few longer, erect dark hairs. Antennæ slightly passing the middle of the elytra and moderately pectinate in the male; scarcely reaching the middle of the elytra, and rather strongly serrate in the female. Head with a small median and the usual lateral impressions. Prothorax transverse; when viewed from above, the apex is nearly truncate, the sides nearly straight and parallel for a short distance, then broadly rounding into the base, which is rather strongly arcuate. Elytra parallel or nearly so in both sexes, the apices not appendiculate in the male, but separately rounded and not evidently produced in either sex.

Length, 3-3.2 mm.

Taken in the Sierras near San Bernardino and at Lake Tahoe.

There is no variation in the eight examples seen. The small size, narrow form, and almost entirely black color may be relied upon to distinguish this species from all others having the antennæ of the male pectinate, and the elytra non-appendiculate. The length of the antennal branches is approximately equal to the diameter of the eye as seen from the front.

#### 61. *Malachius prolixicornis*, sp. nov.

Slender, black, feebly bronzed, surface shining throughout; prothorax very narrowly margined with yellow except (in some specimens) for a short distance at the middle of the apex; mouth-parts, labrum and epistoma yellow; antennæ more or less pale toward the base; legs pale, femora clouded except at base. Antennæ in the male strongly pectinate from the fifth joint, reaching or even passing the tips of the elytra; in

the female feebly serrate, not reaching the middle of the elytra. Head with the usual frontal impressions, surface polished, finely, sparsely punctulate. Prothorax slightly longer than wide, widest in front of the middle, sides sinuate before the hind angles, apex a little wider than the base, all the angles rounded; surface polished and sparsely punctulate; pubescence fine, semirecumbent, intermixed with longer, erect hairs. Elytra parallel (male), or a little wider behind (female); a little less shining than the prothorax, but with similar pubescence; punctuation sparse and fine; tips obliquely, separately prolonged in the male, broadly rounded in the female, narrowly yellow in both sexes.

Length, 3.3-3.6 mm.

Described from three males and one female from Pasadena, kindly given the writer by Dr. Fényes, who took them with others by beating.

The form of the prothorax, and the extravagantly long antennæ with branches half as long as the elytral width, render this the most remarkable species of the genus yet discovered in our territory.

Following is Horn's table enlarged to include the new species here described. This table is perhaps not the best that could be devised, but as a thorough revision of the Malachiini is much needed, it has not been thought advisable to attempt a thorough investigation of the genus at this time. It is quite certain that there are other undescribed species of *Malachius* in collections.

#### TABLE OF SPECIES OF MALACHIUS.

Antennæ of male serrate.

Elytra not appendiculate in the male.

Antennæ with the second joint similar in length and form to the third; size large (Atlantic Region).....*M. æneus*.

Antennæ with second joint much smaller than the third; greenish black, with small yellow spot at sutural angle in both sexes; size small (California).....*M. biguttulus*.

Elytra appendiculate in the male.

Tips of elytra not spiniform in the male, the appendix visible from above.

Thorax with the sides broadly reddish yellow; both it and the elytra, especially toward the base, with numerous erect hairs; elytral tips reddish in the female only....*M. auritus*.



- Thorax entirely dark; erect hairs nearly or quite wanting.
- Entirely dark blue-green; antennæ strongly serrate; size large (4-4.75 mm.).....*M. inornatus*.
- Dull black, sutural tips yellow, antennæ feebly serrate; size small (2.5 mm.).....*M. pristinus*.
- Tips of elytra spiniform in the male, the appendix not visible from above.....*M. acutipennis*.
- Antennæ pectinate in the male.
- Elytra appendiculate in the male.
- Appendices not visible from above.....*M. spinipennis*.
- Appendices visible from above.
- Elytra of male either entirely yellow or tipped with pale rufous; appendix slender; hind tibiæ yellowish testaceous.....*M. mirandus*.
- Elytra of male entirely blue, appendix long but not very slender; legs blue-black.....*M. thevenetii*.
- Smaller, surface lustre greenish; appendix broad, rounded at tip.....*M. directus*.
- Elytra not appendiculate in the male.
- Prothorax wider than long, antennæ (male) shorter than the body.
- Elytra bluish with broad oblique pale vitta from humerus to sutural angle.....*M. macer*.
- Elytra bluish tipped with yellow in both sexes....*M. mixtus*.
- Elytra entirely black, sides of thorax broadly reddish yellow; form rather stout.....*M. ulkei*.
- Elytra black with slight greenish lustre, tipped with yellow in both sexes; thorax with sides very narrowly pale; form slender.....*M. montanus*.
- Almost entirely black without metallic lustre; form slender.  
*M. nigrinus*.
- Prothorax longer than wide, antennæ (male) fully as long as the entire body.....*M. prolixicornis*.

## 62. *Listrus definitus*, sp. nov.

Slender, parallel; black, shining; pubescence black and cinereous; the black hairs forming a rather wide median thoracic vitta, which is a little dilated posteriorly; at the apical third of the elytra is a wide transverse fascia, which is a little prolonged anteriorly along the suture; and between the fascia and the base is a longitudinal series of three spots on each elytron, with occasionally faint indications of one or two smaller spots near the fascia. Head rather sparsely punctate; antennæ reaching (female) or slightly passing (male) the hind margin of the prothorax, joints two to five pale, all the joints longer than wide. Prothorax very slightly longer



than wide, sides moderately, arcuately divergent and rounding into the base; basal angles not evident, apical angles very obtuse, rounded; punctuation moderately strong, closer than on the head, a little coarser but not denser at the sides. Elytra more coarsely punctate. Legs entirely pale.

Length, 2.3-2.5 mm.

Described from a small series taken at Pasadena, in April, by Dr. Fényes.

In the male the tibiæ are armed as usual, the fifth ventral segment is truncate and just perceptibly sinuate at tip, the elytra rounded as usual at the apex. In the female the fifth ventral is evenly rounded at the tip, and the elytra are separately, acuminate produced at the apex. The prothorax is longer than wide,—a character entirely unique among our species,—and the peculiar elytral sexual characters mark this as the most interesting species of the genus yet discovered. It should stand at the end of the genus.

The species of the genus are rather closely allied and difficult to determine with the existing literature. It seems not to have been noticed, or at least recorded, that the males of all species have the tibiæ on all the feet mucronate at tip. The mucros of the front and hind tibiæ are long and acute, though moderately stout, while that of the middle tibia is short and blunt. I have not been able to detect any such armature in the females. The males are often more slender than the females, and in some species, at least, the antennæ in this sex are distinctly longer.

### 63. *Dasytes musculus*, sp. nov.

Moderately elongate and convex, feebly shining; black throughout except the mandibles, which are more or less pale, and the tibiæ and tarsi, which are inclined to become picescent. Pubescence coarse, cinereous, rather dense, short, closely recumbent, uniformly distributed. Head rather finely, not closely punctate; frontal impressions nearly obsolete, eyes not

very prominent. Antennæ scarcely reaching the hind margin of the prothorax, feebly incrassate, joints four to ten as wide as or (the outer ones) a little wider than long. Prothorax nearly as wide as the elytra across the humeri; sides rather strongly rounded behind the middle, convergent and feebly arcuate anteriorly, rounded posteriorly into the base, the basal angles scarcely defined; margin serrulate and with close set, even fringe of recurved hairs; disk moderately strongly, not closely, punctate, the lateral rugose area not limited by a distinctly impressed line, but well defined by the break in the vestiture. Elytra fully one-half longer than wide, distinctly wider posteriorly, apex obtusely rounded; disk finely and not closely punctate. Beneath with fine, sparser, appressed pubescence; the anterior portion of the body more evidently punctate, the abdomen very finely so.

Length, 2.2-2.6 mm.

Described from eight examples taken at Pomona, Pasadena and Riverside.

*D. musculus* agrees with *D. lineellus* in the coarse, closely appressed vestiture without trace of erect hairs, but differs in its smaller size, less robust form, and uniform disposition of the pubescence.

#### 64. *Dasytes clementæ*, sp. nov.

Moderately elongate and convex, polished, black with strong greenish aeneous lustre, legs entirely rufotestaceous; palpi and antennæ blackish, the latter with the first three or four joints pale, the basal joint sometimes more or less obscure. Pubescence entirely luteo-cinereous, rather long, suberect, consisting, especially on the elytra, of somewhat uniformly mixed, shorter, more inclined and recurved, and longer, a little less inclined, hairs. Head narrower than the prothorax, sparsely, moderately finely punctate, and with the usual frontal impressions. Antennæ (male) distinctly longer than the head and prothorax, joints all longer than wide; in the female scarcely longer than the head and thorax, the intermediate joints barely longer than wide, the eleventh as long as the two preceding in both sexes. Prothorax about one-half wider than long, widest and rather strongly rounded behind the middle, quite strongly convergent in front, a little less so behind; hind angles obtuse but distinct, front angles rounded, side margins serrulate; disk finely, sparsely punctate, with an abruptly limited rugose area. Elytra less than half wider than the prothorax, slightly wider behind the middle; apex broadly rounded, finely distantly serrulate; disk rather sparsely but moderately coarsely punctate. Beneath finely, sparsely punctate and pubescent.

Length, 2.5-3 mm.

Described from specimens collected on San Clemente Island.

*D. clementæ* is a near ally of *D. dissimilis*, but is quite distinct from the latter species because of its stouter form, more strongly aeneous lustre, longer, less uniform pubescence, and entirely pale legs.

### 65. *Dasytastes insularis*, sp. nov.

Black, elytra slightly bronzed; head, prothorax, legs, and antennæ, except the three outer joints, rufous. Pubescence much inclined and without trace of erect hairs; punctuation rather fine and sparse throughout, a little coarser toward the base of the elytra than elsewhere. Head a little convex, front scarcely impressed, eyes moderately prominent. Antennæ subequal in length to the head and prothorax, joints four to ten as broad as long, the outer submoniliform; eighth a trifle smaller than the seventh and more evidently smaller than the ninth; eleventh as long as the two preceding combined. The eighth joint is usually quite pale like all that precede, but is sometimes a little obscure. Prothorax unusually convex, about one-third wider than long, widest near the middle, base and apex subequal; sides rather strongly rounded, and distinctly sinuate before the hind angles, which are obtuse but defined; margin serrulate and with an even fringe of recurved hairs; pubescence dark. Elytra about two-thirds longer than wide, widest behind the middle, strongly transversely impressed from side to side behind the base, the sides sinuate behind the humeri; apex obtusely rounded; pubescence luteo-cinereous in the transverse impression and at the apex, but otherwise dark. Beneath finely punctate and sparsely pubescent.

Length, 1.8-2 mm.

Specimens taken on Santa Catalina Island.

*D. insularis* is a very singular species, which however must be placed in *Dasytastes* because of its feeble, unguar appendages and the lack of lateral rugose area on the pronotum. It is without much doubt confined to the island. It should follow *D. bicolor* in Casey's table.

### 66. *Elasmocerus californicus*, sp. nov.

This Californian species, which passes as *terminatus* in some collections, and as "n. sp." in others, seems

surely distinct from the eastern species, and the above name is therefore proposed for it.

The chief differences noted are as follows:—

*Elasmocerus terminatus.*

Antennæ 10-jointed, the last joint in the female fully as long as the preceding joints combined.

Punctuation, especially of the elytra, closer and much coarser.

Prothorax yellow, with a larger or smaller discal spot.

Elytra with lateral margin pale in front, the transverse median pale spot feeble, or with only a slight inward prolongation of the yellow margin.

Abdomen entirely yellow in the male, the sixth segment black in the female.

Legs black.

*Elasmocerus californicus.*

Antennæ 8-jointed (male), 9-jointed (female); the last joint in the female scarcely as long as the preceding joints combined.

Punctuation comparatively fine.

Prothorax black.

Elytra with a transverse median pale spot which usually reaches nearly to the suture, but is rarely marginal; the margin of the elytra not pale in front of the spot.

Abdomen more or less yellow along the middle, the sides black in both sexes.

Legs bicolored.

In the male the three or four antennal joints following the second are so short and closely connate that they are difficult to count. In the single male of *E. californicus* here described the number of joints is surely eight, but other examples will, perhaps, show the number to be nine, as in the female. Specimens from both middle and Southern California have been seen by the writer.

### Hemiptychus.

The species of this genus are moderately numerous in the United States, but they offer so little in the way of differential characters that their identification is a matter of much difficulty. The following species seem in one or two respects so different from any known to us that the very brief diagnoses here given should be sufficient for their recognition.

**67. *Hemiptychus integer*, sp. nov.**

Dark brown, moderately elongate, oval, a little more suddenly rounded in front; sides of prothorax more convergent and less curved than usual. Surface moderately shining, finely pubescent, the pubescence not concealing the surface sculpture, which is fine, the coarser punctures present as usual, but inconspicuous. Sides of elytra with two complete marginal striæ, and a third within these two, extending from the humerus to the middle. Superior outline when viewed in profile not at all gibbous, but equally curved before and behind. Ventral sutures arcuate anteriorly at the middle.

Length, 2.3-2.8 mm.

Collected at Lake Tahoe.

The entire lateral striæ, with the inner accessory one, are peculiar to this species.

**68. *Hemiptychus luteotectus*, sp. nov.**

Elongate-elliptical, rufocastaneous, densely clothed with short, recumbent luteous hair, which nearly conceals the surface sculpture; this however, when visible, is seen to be very fine, with the coarser punctures sparse and finer than usual. The lateral striæ of the elytra are two as usual, and extend from the apex to a little in advance of the middle. In well preserved specimens the vestiture is denser, in narrow lines on the disk of the elytra, giving a feebly vittate appearance. The form as seen from above is equally rounded at the extremities; when viewed in profile it is a little more strongly arched in front. The length is four-fifths greater than the width.

Length, 2.9-4 mm.

Taken at Riverside, Palm Springs and Yuma.

**69. *Hemiptychus palliatus*, sp. nov.**

Similar to the preceding, but distinctly smaller and stouter, being barely one-half longer than wide, and more densely clothed with yellowish white recumbent hair, which completely conceals the surface. The form when viewed in profile is more strongly convex, with the declivity much longer and more gradual posteriorly than in front.

Length, 1.9-2 mm.

Collected at Yuma.



### 70. *Polycaon megalops*, sp. nov.

Allied to *P. confertus*, with which alone it can be compared because of its ten-jointed antennæ. It differs most conspicuously in the very large and prominent eyes, which are separated by three times their own diameter when viewed from the front, while in *P. confertus* they are distant fully six times their own diameter. Further differences are as follows: prothorax distinctly wider than long, not canaliculate, the transverse impression less distinct, the granulation less dense; elytra two and one-half times as long as the head and prothorax (scarcely more than twice as long in *P. confertus*), rather sparsely punctate, much as in the male of *P. confertus*, but not at all granulate towards the sides and apex; second and third tarsal joints with only a group of projecting hairs, instead of the densely matted brushes which in *P. confertus* assume the appearance of membranous lobes.

Two examples have been seen, both apparently females; one, in the writer's collection, is 5 mm. in length, the other, in the collection of Dr. Van Dyke, is 7 mm. long. The larger specimen barely reaches the stature of the smallest *P. confertus*, which varies in length from 7.5 to 10.5 mm.

Specimens taken at Pomona, Los Angeles County.

### 71. *Aphodius ungulatus*, sp. nov.

Oblong, scarcely broader posteriorly, antennæ testaceous, club not darker; head piceous, margin paler; sides fimbriate posteriorly; surface not distinctly tuberculate, punctures moderate, a little finer and sparser at the middle; clypeus emarginate and distinctly though not sharply angulate at sides. Thorax piceous with paler side margins, about one-half wider than long, sides parallel, slightly arcuately narrowed in front, margins finely and rather sparsely fimbriate with short hairs; hind angles obtuse; disk finely, rather sparsely punctate at middle, more closely and coarsely at sides, with larger punctures intermixed; marginal basal line distinct at sides but tending to become obsolete at middle. Elytra dull yellow, immaculate, but with a lateral, not very sharply defined, fuscous shade reaching from the humerus to beyond the middle; striæ fine, with fine punctures which are well separated, interspaces nearly flat, with two more or less regular series of punctures which are about equal to those of the striæ. Beneath brownish, abdomen paler; surface alutaceous, abdomen and sterna, except at sides, sparsely punctate. Mesosternum not



distinctly carinate. Anterior tibiæ smooth in front, externally tridentate, above crenate, first tarsal joint shorter than the second; first joint of hind tarsi as long as the next two.

Length, 5-5.75 mm.

Occurs at Pomona and Pasadena, though rarely, during December and January.

The claws are very slender and distinctly longer than in any other species observed. There are slight differences in the length and stoutness of the spurs of the front tibiæ in the five examples from which the description is drawn, but it is not certain that these differences are sexual in nature. *A. unguulatus* must be placed next to *A. serval*, from which it may be easily distinguished by the unspotted elytra, shorter first joint of hind tarsi, and longer claws.

## 72. *Amphicoma cooperi* Horn.

This cannot be placed as a synonym of *A. ursina*, as is announced by Horn in his review of the species of this genus.\* *A. ursina* differs from all our remaining species of the genus in the eyes, which are more widely separated on the vertex than the width of the front between the antennæ, and in the hind tibiæ, which have but a single spur, characters which are quite independent of sex, and which appear to have escaped previous observation. *A. cooperi* is in accord with the other species in these particulars, and if it is to be placed with either of the described species it must be with *A. canina*. It seems best for the present, or until a careful study can be made with ample material, to hold the species as distinct, separating it from *A. canina* by its smaller size and non-maculate elytra.

---

\* Trans. Am. Ent. Soc., Vol. X, 1882, p. 119.

73. *Idæmea californica*, sp. nov.

Slender, cylindrical, brownish testaceous, pubescent; thighs a little paler at base. Head wider than the thorax, eyes broadly contiguous on the vertex. Antennæ slightly longer than the body in the female, nearly twice as long as the body in the male, clothed externally with rather dense, very short, erect hairs, which become sparser and longer toward the base. Prothorax cylindrical, a little dilated at the middle; surface finely, closely, but somewhat unevenly punctate. Elytra about three-fourths wider than the prothorax, gradually feebly narrowed behind, closely and a little more coarsely punctate than the thorax.

Length, 13-16 mm.

Beaten from live-oak, and taken at electric light at Pomona and Pasadena, during April and June. There is a specimen from Yuba County in the Horn Collection.

*I. californica* is very similar to *I. fulleri* in a general way, but the latter is somewhat stouter and more coarsely punctured, the eyes are well separated on the vertex, and the antennæ in the male are described as being but little longer than the body.

74. *Eusattus difficilis* Lec.

This species was based upon a series, part of which was taken at San Diego and part at Vallecito. An examination of the Le Conte material shows that two species were confused under this name. The San Diego specimens, which may be considered the true *E. difficilis*, differ from every other species known to me in possessing a well developed scutellum. The Vallecito specimens are apparently identical with a species which was described by Le Conte as *E. convexus*, his type coming from near Long's Peak, Colorado. This species he describes as closely resembling *E. difficilis*, and subsequently it was placed in synonymy. *E. convexus* is a rather common species, occurring from El Paso, Texas, westward to the desert regions of California, while

*E. difficilis*, so far as the observation of the writer goes, is confined to the coast region about San Diego. It is obvious from the above that the name *convexus* must be restored.

### 75. *Notoxus caudatus*, sp. nov.

Moderately elongate, piceous brown, legs somewhat paler, the thoracic horn rufous; the elytra are best described as brownish black, each with two testaceous spots, one transversely oval about two-fifths from the apex, the other antemedian, larger, subtriangular, extending forward laterally to the humeral umbone; or the elytra may be described as testaceous, with a triangular scutellar area, a median band, and the apex broadly dark; these dark areas confluent both along the suture and side margin; the apex of the elytra is in some specimens narrowly, indefinitely, and indistinctly paler. Pubescence cinereous, fuscous in great part on the darker portion of the elytra, intermixed with sparse, longer, semierect hairs. Head nearly flat, rather closely but not deeply punctate; eyes about twice as long as the tempora. Prothorax transversely globose, the horn broad, dilated, the margin crenulate at sides except in front, crest rather strongly elevated, distinctly margined, the margin feebly or scarcely crenulate; surface of prothorax sparsely, finely punctate at middle, more closely at sides. Elytra widest just behind the middle, the apex rather suddenly obtusely rounded and nearly similar in the sexes. In the males there is a small, rounded, impressed area immediately before the middle of the apex of each elytron; this area is clothed with short, dense, closely recumbent cinereous hair; in the same sex the pygidium is strongly produced to an acute point, the sides of the process are sinuate before the apex; and the fifth ventral is broadly, arcuately emarginate, the limiting angles of the emargination being sharply defined and a little reflexed. In the female the elytra and tip of abdomen are unmodified.

Length, 3.1–3.5 mm.

Described from a series of eight specimens collected at Santa Fé, New Mexico, by Dr. A. Fényes.

*N. caudatus* is perhaps most like *N. nuperus* in general facies, but the extraordinary sexual characters of the male are nowhere approached in our species. There is considerable variation in the extent of the pale spots of the elytra, more especially the anterior ones, which are in some specimens almost lacking.

76. *Xylophilus brunnescens*, sp. nov.

Elongate-oblong, moderately convex, rufotestaceous, the elytra becoming darker posteriorly, head and under surface brownish, legs and antennæ paler; integuments shining; pubescence pale brown, moderate in length, not dense. Head much wider than the prothorax, sparsely, finely punctate; front concave between the eyes, declivous anteriorly, the epistomal suture fine and feeble. Eyes moderately large, separated on the front by about three-fourths their own width, the inner margin faintly emarginate; tempora parallel and about one-third the length of the eyes. Antennæ fully two-thirds the length of the body, scarcely thicker apically; first and third joints subequal, second joint half as long, joints four to ten subequal, a little shorter than the third, last joint a little longer and wider than the tenth, pointed. Prothorax subquadrate, sides parallel and impressed at the middle and again at the front angles, disk with a transverse, subbasal impression; surface more closely and a little less finely punctate than the head. Elytra nearly twice as wide as the prothorax, punctuation coarse, the punctures separated by rather less than their own diameters toward the base, but somewhat finer and less close apically. Beneath shining, the sides of the metasternum with sparse, coarse punctures, the abdomen finely, very remotely punctate. First abdominal suture nearly obliterated but traceable at sides. The first and second segments are subequal, the third a little shorter, fourth equal to the second, fifth one-half longer than the fourth, the apex evenly rounded, the disk unmodified. Front and middle thighs slender, the posterior stout and densely pilose beneath; first joint of hind tarsi about three times as long as the remaining joints united, and about three-fourths as long as the tibia. Front tibiæ with a short, slender spur at the inner apical angle.

Length, 2.3 mm.

Described from a single male taken in the San Bernardino Mountains, in July.

By Casey's table *X. brunnescens* must be referred to his genus *Vanonus*. It differs from any of the species there mentioned in its color and the stout hind thighs, and apparently from all except *X. wickhami* in the well developed tempora. It should be placed before *X. wickhami*. The spur of the anterior tibia will very likely prove to be a male character.

**77. *Xylophilus nucleus*, sp. nov.**

Elongate-oval, subdepressed, brown, prothorax red-brown, legs and antennæ testaceous; surface moderately shining, pubescence fine, sparse, and recumbent. Head finely, closely punctate, front convex, epistomal suture obsolete; eyes large, not evidently pilose or emarginate, extending to the base of the head, distant on the front by a little more than their own width. Antennæ about half the length of the body, not distinctly thickened externally; second joint but little shorter than the first or third, these subequal, the following joints very slightly decreasing in length to the eleventh, which is subequal to the two preceding united. Prothorax small, much narrower than the head, transverse, scarcely narrowed in front, disk deeply, transversely impressed posteriorly; surface closely, rather finely punctate. Elytra elongate-oval, twice as wide at middle as the prothorax, disk obliquely impressed each side near the base, surface finely punctate, the punctures separated by from two to three times their own diameters. Beneath sparsely, finely punctate; middle coxæ separated by their own width; ventral segments decreasing slightly in length toward the apex, the last segment a little longer; suture between the first two segments completely obliterated at the middle third, but very distinct laterally. All the legs very slender, the hind femora unmodified; basal joints of hind tibiæ equal to the remainder.

Length, 1-1.4 mm.

Two examples taken in the San Bernardino Mountains.

**78. *Xylophilus constrictus*, sp. nov.**

Oblong-oval, subdepressed, piceous or piceo-testaceous, antennæ and legs paler. Head and thorax finely punctate; elytra more strongly and closely so. Head wide, front slightly convex, epistomal suture distinct; eyes not obviously emarginate or pilose, separated on the front by rather more than twice their own width; tempora scarcely evident. Antennæ rather slender, reaching the middle of the elytra; first joint short; second smaller, subglobose, one-half as long as the third; joints three to ten gradually decreasing in length, the eleventh as long as the third. Prothorax narrower than the head, transverse, deeply constricted behind the apex, and again, but less strongly so, along the basal margin, the constrictions in both cases extending entirely across the disk. Elytra oblong-oval, sides broadly arcuate and subparallel, twice as wide at middle as the prothorax. Middle coxæ narrowly but distinctly separated, ventral segments slightly decreasing in length, the last a little longer than the preceding; the suture between the first two segments distinct throughout. Legs very slender; basal joint of hind tarsus slightly shorter than the remainder.

Length, 1.5 mm.



Specimens from the Sierra Madre Mountains (Los Angeles County) and Lake Tahoe.

Described from two examples, in which no sexual characters were observed. The peculiar structure of the prothorax will at once distinguish the present species from any previously described. Neither *nucleus* nor *constrictus* can properly be referred to any of the numerous genera lately proposed by Casey, and for a variety of reasons it seems best for the present to use the old name *Xylophilus*.

### 79. *Rhynchites æratoides*, sp. nov.

This species is almost the exact counterpart of *R. æratus*, from which it seems constantly to differ in the more closely punctate head, broader front, more prominent eyes, and the presence in the male of a longitudinal median impressed line on the first three ventral segments.

*R. æratus* is somewhat variable in the cephalic characters mentioned, and while the writer has seen no specimens that approach the Californian form very closely in this respect, there would be some hesitancy in separating them were it not for the peculiar abdominal characters of the males above mentioned, the detection of which is due to the observation of Mr. Frederick Blanchard, to whom examples were sent for comparison with the eastern specimens in his cabinet.

*R. æratoides* occurs at Redondo (on *Eriogonum cinereum*), Pomona and Pasadena, from March to June.

### 80. *Cleonus pacificus*, sp. nov.

Robust, parallel, clothed with moderately dense, recumbent, squamiform, cinereous pubescence; the disk of the thorax, except two dorsal vittæ, and the sutural interval of the elytra, glabrous. Beak stout, somewhat dilated at tip, sides and tip glabrous, carinate above in basal two-thirds, not very coarsely or deeply punctate, frontal impression feeble. Prothorax a trifle wider than long, sides parallel in basal half, thence moderately rounded and convergent, without apical constriction; basal lobe



angulate; disk with moderate basal depression, obtusely carinate anteriorly; punctuation rather strong, sparse, and unevenly distributed. Elytra a little more than twice as long as wide, about one-third wider than the prothorax; humeri obliquely rounded; apices divergent but not produced; the series of rather fine punctures unimpressed except the sutural; third and sixth intervals moderately elevated at base. Beneath speckled with subdenuded punctures. Legs moderate, lobes of third joint of hind tarsi completely spongy pubescent beneath.

Length, 13 mm.

A single specimen was taken on *Astragalus crotalaræ*, at Redondo, in April.

According to Casey's table the species should come between *C. inornatus* and *C. canescens*.

### 81. *Cleonus erysimi*, sp. nov.

Elongate, subparallel, densely clothed with rather fine whitish recumbent squamiform hairs which nearly conceal the sculpture; the disk of the pronotum, except two anteriorly convergent dorsal vittæ, and the sutural and eighth elytral intervals, glabrous. Beak nearly three-fourths as long as the prothorax, scarcely broader at tip, densely, finely punctate with coarser punctures intermixed, more or less distinctly carinate, frontal impressions well marked. Prothorax slightly longer than wide, sides straight and nearly parallel almost to apex, where they are feebly narrowed and slightly constricted; basal lobe moderately prominent, rounded or truncate at apex; basal excavation not deep; disk obtusely prominent anteriorly along the median line, but not distinctly carinate; coarser punctures sparse at middle, more numerous at sides. Elytra but little wider than the prothorax, apices divergent but not produced, punctures of striæ not coarse, the striæ not impressed; third and sixth intervals feebly elevated at base. Lower surface and legs about as usual.

Length, 7.5-11.5 mm.

Many specimens have been seen, occurring only on the flowers of *Erysimum asperum*, on the sand-hills overlooking the ocean at Redondo; March and April.

The sutural interval is pubescent at base and throughout its outer fourth, and the eighth is thinly clothed posteriorly. The ocular lobes are nearly wanting. The third joint of hind tarsi is completely spongy pubescent

beneath. *C. erysimi* would be associated with *C. quadrilineatus* Chev., but this is said to have the thorax distinctly carinate and (according to Le Conte's description) the third joint of the hind tarsi is not broader than the second and not spongy beneath.

## 82. *Anthonomus apertus*, sp. nov.

Short, stout, black, moderately shining, without aeneous lustre, clothed sparsely with white pubescence, which is more conspicuous beneath. Beak somewhat variable, rather long, shining and more or less punctate at the apex, dull and more or less longitudinally strigose and sulcate behind the insertion of the antennæ; frontal fovea distinct, punctiform. Antennæ testaceous, club piceous; funicle seven-jointed, slender, its second joint slightly longer than the third, outer joints scarcely as wide as long. Prothorax about twice as wide as long, sides moderately rounded, more strongly in front, apex feebly constricted, punctuation strong and rather close. Elytra short, abruptly more than one-third wider than the prothorax, sides parallel to beyond the middle; striæ moderate, punctures rather fine, not very closely placed; intervals wide, feebly convex, slightly transversely rugose; scutellum densely clothed with white pubescence. Femora all armed with a distinct spiniform tooth; front and middle tibiæ slightly curved inward at tip; tarsi piceo-testaceous, the first joint of anterior pair about one-half longer than the second; claws armed with a moderately long tooth.

Length, 1.75-2.25 mm.

Several examples were taken at Riverside and Los Angeles.

*A. apertus* belongs to the *scutellatus* group, and is most closely related to *A. ebeninus*, from which it differs greatly in size and also in several minor details. It resembles *A. morulus* very closely, a specimen sent to Dr. Horn being so determined, but aside from the group characters, it is more shining, with more slender antennæ, and longer femoral teeth.

## 83. *Anthonomus helianthi*, sp. nov.

Pitchy brown, beak red-brown, legs paler; densely clothed with elongate greyish white scales which become faintly yellowish on the middle of the prothorax and on the basal portion of the third (sometimes also the second

and fourth) elytral interspace, and still more faintly on the fifth interspace behind the middle. Beak a little longer than the head and prothorax in the male, still longer in the female, finely carinate behind the insertion of the antennæ. Antennæ slender, inserted beyond the middle in both sexes; funicle seven-jointed, second joint of funicle two-thirds as long as the first and equal to the third and fourth together. Prothorax slightly wider than long, narrowed from base to apex, sides nearly straight in basal half, thence more rounded and feebly constricted before the apex; surface densely punctured, the sculpture, however, entirely concealed by the vestiture. Elytra distinctly wider at base than the prothorax, humeri rounded, sides parallel to the middle, striæ apparently finely impressed. Legs moderate, front and middle femora armed with a very small, acute tooth; the hind tibiæ in the male straight externally, dilated internally at the apex, the dilated portion with parallel sides. Claws with a long, slender tooth which is in contact with its fellow at the tip.

Length, 2.75-3 mm.

The species is described from two males and one female beaten from sunflowers, April 20, at Pomona.

It belongs to the subgenus *Cnemocyllus*, and should be placed before *A. subvittatus*, at the head of the group.

#### 84. *Anthonomus tahoënsis*, sp. nov.

Elongate-oval, black, legs and beak sometimes entirely or in part reddish brown; antennæ rufous, club darker; clothed rather sparsely above and beneath with uniformly cinereous scales; the scales vary a little in form, being somewhat broader on the parapleuræ, but they are as a rule subparallel and about twice as long as wide, not in the least overlapping and scarcely in mutual contact, so that the color of the derm is not concealed. Head finely sulcate between the eyes, the latter moderately convex, their hind margins slightly free; beak rather long, evenly arcuate, rather densely but not very coarsely punctate in the male, smoother in the female. Antennæ not very slender, second joint of funicle a little longer than the third, the latter subequal to the fourth. Prothorax nearly as long as wide, sides broadly arcuate and quite strongly convergent, feebly constricted at apex; surface rather coarsely and closely punctate, strongly shining. Elytra abruptly fully one-third wider at base than the prothorax, sides subparallel in basal two-thirds, thence gradually narrowed, apex parabolically rounded. Striæ fine and rather finely punctate, intervals wide, nearly flat, and almost smooth. Last ventral segment scarcely longer than the preceding in either sex, not relatively longer in the male. Legs slender; front thighs not more stout, and armed with a slender, acutetooth; hind

tibiæ of male a little bent at tip, the inner margin quite deeply sinuate; in the female the hind tibiæ are scarcely bent and much less deeply sinuate internally, but are visibly stouter than the others and very decidedly stouter than the corresponding ones in the male. The apical mucro is curved downward. Claws with long, slender, approximate teeth.

Length, 2.75 mm.

Described from a single pair given the writer by Dr. Fényes, who took them at Tallac (Lake Tahoe) early in July.

*A. tahoënsis* is most closely related structurally to *A. subvittatus*, but differs from it and all other members of the subgenus in its sparse vestiture.

### 85. *Anthonomus stolatus*, sp. nov.

Elongate-oblong, pitchy brown; legs, beak (except extreme tip) and antennæ, reddish brown; extremely densely clothed above and beneath with broad, rounded, overlapping scales; the scales become narrower on the head and along the middle of the abdomen, and are very small and especially densely matted on the scutellum; the prevailing color is pale yellow, varied with yellowish brown on the upper surface, the prothorax showing three pale vittæ, and the elytra exhibiting indistinctly the usual pale markings, viz., a post-scutellar sutural line, a line on the fourth interval behind the middle, and another on the sixth interval beginning at the base; these markings at best are faint, and some examples show scarcely a trace of them. Beak as long as the head and thorax (in the male) or somewhat longer (in the female), parallel, moderately strongly, evenly arcuate, glabrous except at base, punctate and carinate at sides, more finely punctate and scarcely carinate above. Head densely clothed with scales, apparently with an elongate puncture or short sulcus between the eyes, the latter moderately convex, their posterior margins not free. Antennæ rather slender, third funicular joint two-thirds to three-fourths as long as the second and a little longer than the fourth. Prothorax distinctly wider than long, sides evenly, arcuately convergent and moderately constricted before the apex, which is about three-fifths as wide as the base; surface coarsely, densely, deeply punctate, the sculpture however entirely concealed by the vestiture. Elytra slightly wider at base than the prothorax, sides parallel to apical third, striæ of moderate closely placed punctures, intervals feebly convex, and nearly smooth, shining. Last ventral segment longer than the preceding and somewhat longer in the male than in the female. Legs as usual; the front thighs feebly toothed; claws with a moderately long, acute tooth, which is approximate to its fellow.

Length, 2.3-2.6 mm.

The description is drawn from seven examples, all of which are from San Diego or the vicinity.

The hind tibiae are apparently very slightly more strongly sinuate along the inner margin in the males, but this difference is hardly enough in itself to suggest the placing of this species in the subgenus *Cnemocyllus*; the affinities in other respects are, however, obviously in that direction, and *A. stolatus* should find a place near *decipiens*, from which, and all others of the subgenus (except possibly *A. ligatus*), it differs in the almost entire lack of sexual difference in the posterior tibiae; from *A. ligatus* it may be separated by color and also the longer unguar teeth.

### 86. *Epimechus arenicolor*, sp. nov.

Elongate-oblong, color of derm varying from testaceous to rufous, the thorax and abdomen usually darker and the legs and antennae paler than the elytra; moderately densely clothed with broadly oval, pale, dull yellowish scales, which are nearly uniform in color throughout and as a rule do not overlap, but become denser or even strongly imbricate in the position of the usual paler vittae in *Cnemocyllus*. Head clothed with sparser, more elongate scales; eyes feebly convex, hind margin not free; front narrow, the eyes separated by a distance distinctly less than their own width as seen from the front; beak moderately long and evenly arcuate, punctate in longitudinal series, not distinctly carinate above at base; antennal funicle, slender, second joint one-half longer than the third. Prothorax evidently transverse, sides subparallel at base, gradually arcuately narrowed in front, apex very slightly constricted; surface deeply, rather densely punctate. Elytra slightly wider at base than the prothorax, sides parallel to apical third, apex rather narrowly parabolically rounded; striae punctures nearly as wide as the intervals, the latter feebly convex and slightly rugose. Last ventral segment (in the male) longer than the two preceding united. Femora rather stout, the anterior feebly toothed; hind tibiae (in the male) distinctly but not strongly curved at apex; claws simple.

Length, 2-2.5 mm.

All specimens seen are from Arizona—Phoenix (Stromberg) and Palomas.



This species must by the simple claws be placed in *Epimechus*, but in every other respect it is closely allied to the members of the subgenus *Cnemocyllus*. The unguual teeth in *Anthonomus* vary greatly in development, and it is to be doubted whether the genus *Epimechus*, which is apparently founded on this character alone, will prove tenable. A similar condition of affairs obtains in *Apion*, which it has not been thought proper to divide although there is more to warrant a separation than in the present instance.

*E. arenicolor* by its comparatively dense scaly vestiture must be associated with *E. nevadicus* and *E. æmulus* (see following description); it is, however, less densely clothed than either of these, and of quite a different color; in the extreme length of the last ventral of the male (females have not been seen) it differs conspicuously from all allied forms; this segment in other species of *Epimechus*, in fact, being scarcely at all longer than the fourth. The claws in *E. nevadicus*, *E. arenicolor*, and *E. æmulus* are distinctly smaller than in the non-squamose species of the genus *E. adspersus*, *E. mimicus*, etc.

### 87. *Epimechus æmulus*, sp. nov.

Elongate-oval, densely clothed with whitish and grayish brown scales, giving the species an appearance closely resembling *E. nevadicus*, with which it agrees in most points of structure. The white scales are, however, more conspicuous on the elytra, and are chiefly confined to the alternate intervals, beginning with the second. There is a transverse fascia of the darker scales in the usual position behind the middle, beginning on the third interval and extending to the sides of the elytra; this darker fascia is bordered both before and behind by a conspicuous spot of white scales on alternate intervals, and on the fourth and sixth these white scales extend to the base of the elytra. The lateral pale vittæ of the thorax are nearly lacking in the type. The funicle of the antennæ is seven-jointed, the second joint much longer than the third, the latter equal to the fourth. Last ventral equal to the preceding.

Length, 3.1 mm.



Described from one example of doubtful sex taken at or near San Diego.

It is distinguishable at once from *E. nevadicus*, its nearest ally, by its larger size and seven-jointed funicle.

### 88. *Ceutorhynchus pervestitus*, sp. nov.

Oblong-oval, densely clothed throughout with broad, overlapping scales of whitish, pale brown, and dark brown intermingled. Antennæ inserted at the middle of the beak; piceous; funicle seven-jointed. Beak moderately elongate, punctate and striate, smoother and more shining at tip, squamose at base; front concave, vertex finely carinate; eyes concealed in repose. Prothorax about one-third wider than long, sides subparallel from base to middle, then rather suddenly rounded into the deep apical constriction; apical margin elevated, notched at middle; lateral and dorsal tubercles prominent, those on either side apparently joined by a transverse elevation; dorsal channel deep. In addition to the vestiture above mentioned, there are sparsely placed blackish, erect, piliform scales, which are more noticeable at the sides and in a patch on either side of the dorsal channel at base. Elytra about one-third wider than the prothorax, sides nearly straight, a little narrowed behind; each interspace with a row of blackish, erect, linear scales like those on the prothorax; the alternate intervals on the disk, beginning with the second, tessellate with patches of these scales; the scales proceeding from the stria punctures are long, bristle-like, and closely recumbent. Femora moderate, distinctly toothed; last joint of tarsi equal to the two preceding; claws with a rather small but acute subbasal tooth.

Length 2.75 mm.

Described from a single example taken at Bakersfield.

The type is a male, having the last ventral segment vaguely concave, and the middle and hind tibiæ unguiculate. The vestiture is so dense as to entirely conceal the color and sculpture of the integuments, but they are probably much the same as in allied forms. This species must be referred to the *subpubescens* group, from all the members of which it differs greatly in appearance.

### 89. *Baris heterodoxa*, sp. nov.

Oblong-oval, moderately robust, black without distinct aeneous lustre, surface of pronotum polished between the punctures, that of the elytra

shining but minutely sculptured; setæ nearly lacking on the pronotum, the elytra with scattered but somewhat numerous subsquamiform, white, recumbent hairs which are about as long as the width of the interspaces. Head finely, sparsely punctate; beak as long as the prothorax, moderately stout, strongly, evenly arcuate, closely punctate. Antennal funicle rather slender; the second funicular joint about as long as wide, following joints gradually more transverse but only slightly increasing in width, the eighth less than half as wide as the club. Prothorax distinctly transverse, sides nearly parallel and feebly arcuate from base to apical fourth, then strongly rounded and constricted at the apex; surface densely punctate except toward the middle of the disk, where the punctures are separated by nearly their own width; a distinct fusiform, impunctate median line from the apical third nearly to the base. Elytra a little wider than the prothorax and scarcely twice as long; sides very slightly convergent behind, rather broadly but subparabolically rounded at apex. Striæ shallow, with distant, moderately well defined punctures; intervals about three times as wide as the striæ, all very numerous, confusedly punctate, the punctures as a rule not more than one-fourth the width of the intervals. Abdomen shining, moderately coarsely, but not densely punctate. Prosternum densely, coarsely punctate, the intercoxal process rather less than half the coxal width.

Length, 3.7 mm.

One specimen taken in Los Angeles County.

In the vestiture and sculpture of the elytra this species is perhaps the most extraordinary in our fauna. The second and third intervals are just visibly wider than the others, but the punctuation is equally close and confused from base to apex on all. *B. heterodoxa* seems to fall between *B. vespertina* and *B. oblongula* in Casey's table, but is evidently not closely related to either.

#### 90. *Baris monticola*, sp. nov.

Elongate-oval, moderately convex; black throughout, with very faint aeneous lustre; surface rather strongly shining. Head finely, sparsely punctate; beak three-fourths as long as the prothorax, strongly curved, rather stout, strongly and closely punctate at sides, more finely and less closely above, especially toward the apex. Antennæ about as usual, the basal joint of the club constituting a little less than half its mass, polished but pubescent toward the apex. Prothorax very little wider than long, sides moderately convergent and nearly straight almost to the apex,

then strongly rounded and with a very short apical constriction; surface strongly and rather coarsely punctate, the punctures as a rule separated by less than their own diameters; they are somewhat finer and less close at the middle of the disk anteriorly, and at the extreme sides are more or less coalescent, forming longitudinal rugæ; disk without median smooth line or with but faint traces of it. Elytra a little less than twice as long as the prothorax, slightly wider, the sides feebly convergent posteriorly and broadly elliptically rounded at apex; striæ rather fine, moderately deep, obsoletely punctate; intervals flat, about two and one-half times as wide as the striæ; the second and third fully three times as wide as the striæ; second, third, and ninth with more or less confused punctuation, the other intervals with a single series which is, however, somewhat confused or irregular at base; the punctures as a rule not more than one-third as wide as the intervals, and distant from each other by about their own diameter. Beneath rather coarsely and closely punctate; prosternum nearly flat, the front coxæ separated by rather more than half the coxal width.

Length, 3.4-4 mm.

Described from a good series taken by Mr. F. S. Daggett in Bear Valley (San Bernardino Mountains), at an elevation of about 6,400 feet.

*B. monticola* would fall next to *B. aprica* according to Casey's table, but it is very distinctly different in its finely punctate elytral intervals, among other characters. The prothorax is scarcely more strongly declivous in front, as is described of the section in which the elytra are less than twice as long as the prothorax, and to which *B. monticola* is therefore referred. The setæ of the upper surface are very short and inconspicuous.

### 91. *Sphenophorus tardus*, sp. nov.

Moderately robust, black, feebly shining. Beak one-half the length of the prothorax, distinctly arcuate, moderately compressed; punctuation sparse and minute except at base, the usual puncture between the eyes, before the puncture a short, fine, impressed line. Prothorax longer than wide, sides evenly arcuate, a little convergent anteriorly, apex constricted; surface without grooves or elevations, densely, moderately coarsely punctate, with a narrow, median, smooth space, or line, which reaches neither base nor apex. Elytra not at all, or scarcely wider, and but little longer,

than the prothorax; striæ moderately deep and rather finely punctate on the disk, more coarsely so at sides, those of the seventh and eighth striæ being wider than the striæ and thus encroaching on the intervals; intervals flat, alternating in width, the narrower ones uniseriately, the wider ones biseriately, but all more or less irregularly punctate, the punctures rather coarse and very closely placed. Under surface more shining, rather coarsely but less densely punctate than above. Femora moderately punctate.

Length, 6.5-9 mm.

Numerous specimens have been seen, taken at or near San Bernardino.

The species may stand at the end of Division A of Group V of Horn's Synopsis.

# INDEX TO GENERA OF COLEOPTERA OF SOUTHERN CALIFORNIA.

Names of new species in **heavy faced** type.

ABREUS .....	97	Allobates.....	169
Acanthocinus.....	151	Alphitobius .....	173
Acanthoscelis .....	198	Alphitophagus.....	174
Acmæodera.....	119	Amannus .....	146
Acmæops.....	149	Amara.....	45
Acneus.....	106	Amartus .....	98
Acritus.....	97	Amnesia.....	188
<b>volitans</b> .....	238	Amotus .....	187
Actenodes.....	118	Amphicerus .....	134
Actidium.....	80	Amphichroum.....	78
Actium.....	63	Amphicoma .....	139
Actobius.....	69	cooperi.....	255
<b>formosus</b> .....	221	Amphicyrta .....	103
Acylopus.....	82	Amphidora .....	169
Acylophorus.....	67	Amphizoa .....	51
Adaleres.....	188	AMPHIZOIDÆ.....	51
Adalia.....	84	Anapleus.....	95
Adasytes.....	127	Anaspis .....	179
Adelocera .....	107	Anatis .....	84
Adoxus.....	154	Anchastus .....	109
Adranes .....	61	Anchomma .....	88
Ægialia.....	136	Anchyteis .....	106
Ægialites.....	175	Ancyrophorus .....	77
ÆGIALITIDÆ.....	175	Androlyperus.....	157
Æletes.....	97	Aneflus.....	145
Ænigmaticum .....	82	Anelastes.....	107
Agabinus.....	53	Anemia .....	174
Agabus.....	53	Anepsiota .....	65
Agasphærops.....	187	Anepsius .....	163
Agathidium .....	60	Anillus .....	43
<b>virile</b> .....	219	Anisodactylus.....	51
Aglenus .....	89	Anisosticta .....	83
Agonoderus .....	50	Anisotarsus.....	51
Agrilus.....	120	Anisotoma .....	59
<b>gibbicollis</b> .....	241	Anobium.....	132
<b>illectus</b> .....	242	Anomala .....	141
Agriotes.....	111	Anorus .....	106
Agronus.....	188	Anthaxia .....	116
Agyrtes.....	59	ANTHICIDÆ.....	180
Alæphus.....	170	Anthicus .....	181
Alaudes .....	172	Anthobium .....	79
Alaus.....	108	Anthonomus.....	195
Aleochara .....	66	<b>apertus</b> .....	262
Alindria.....	102	<b>helianthi</b> .....	262
Allonyx .....	128	<b>stolatus</b> .....	264
Allopogon.....	106	<b>tahoensis</b> .....	263

Anthophilax.....	149	Baris.....	198
Anthrenus.....	93	<b>heterodoxa</b> .....	267
ANTHRIBIDÆ.....	203	<b>monticola</b> .....	268
Aphanotus.....	173	Batrissus.....	63
Aphodius.....	137	Batyle.....	146
<b>ungulatus</b> .....	254	Belonuchus.....	68
Aphorista.....	87	Bembidium.....	42
Aphricus.....	114	Berginus.....	92
Apion.....	190	Berosus.....	56
Aplastus.....	114	<b>salinus</b> .....	215
Apocellus.....	77	Bidessus.....	52
Apocrypha.....	174	Biotus.....	61
Apristus.....	48	Bius.....	170
Aræoschizus.....	163	Blapstinus.....	171
Aragnomus.....	189	Blechrus.....	48
Aramigus.....	190	Bledius.....	75
Argoporis.....	169	<b>clarus</b> .....	228
Articerus.....	61	<b>deceptivus</b> .....	227
Artochia.....	77	<b>relictus</b> .....	228
Asaphes.....	113	<b>rusticus</b> .....	229
Asclera.....	178	Boletobius.....	74
Asemobius.....	77	Bolitochara.....	66
Aseum.....	143	Bostrychus.....	134
Asida.....	164	Bothriophorus.....	104
Asydates.....	125	Brachylobus.....	50
Atænius.....	137	Brachynus.....	49
Atheta.....	64	Brachypterus.....	98
Athoneus.....	98	Brachytarsus.....	203
Athous.....	112	Bradycellus.....	50
Atimia.....	147	Bradycinetus.....	138
Atomaria.....	91	Brathinus.....	78
Attagenus.....	92	Brontes.....	91
Attalus.....	125	Brothylus.....	144
Auchmobius.....	151	BRUCHIDÆ.....	160
Aulacoscelis.....	151	Bruchus.....	160
Aulanothroscus.....	115	Bryaxis.....	63
Auletes.....	186	Bryobiota.....	67
Aulicus.....	129	Bryoporus.....	74
Aulonium.....	89	BUPRESTIDÆ.....	115
Autalia.....	66	Buprestis.....	116
Axinopalpus.....	48	BYRRHIDÆ.....	103
		Byturus.....	92
		CACTOPHAGUS.....	200
BABIA.....	152	Cafius.....	70
Bacanius.....	97	Calandra.....	200
Bactridium.....	103	CALANDRIDÆ.....	200
Badister.....	46	Calathus.....	46
Bagous.....	194	Callida.....	48
Balaninus.....	199	Callidium.....	144
Bamona.....	67	Calligrapha.....	155
Baptolinus.....	71	Callinus.....	145
<b>punctiventris</b> .....	221	Callimoxys.....	145
Barilepton.....	199	Calloides.....	146
Barinus.....	199		



Calochromus .....	121
Calodera .....	65
Caloderma .....	72
Calopteron .....	121
Calopus .....	178
Calosoma .....	40
Calospasta .....	185
Cantharis .....	185
Canthon .....	136
CARABIDÆ .....	39
Carabus .....	40
Carcinops .....	95
Cardiophorus .....	108
Carebara .....	177
Carphoborus .....	203
Carpophilus .....	98
Cartodere .....	101
Cassida .....	160
Cathartus .....	90
Catorama .....	132
Cenophengis .....	122
Centrinogyna .....	199
Centrinus .....	199
Centrioptera .....	164
Centrodera .....	148
Centronopus .....	170
CEPHALOIDÆ .....	179
Cephaloon .....	179
Cephaloscymnus .....	87
CERAMBYCIDÆ .....	142
Cercus .....	98
Cercyon .....	58
Cerenopus .....	169
Cerophytum .....	115
Cerylon .....	89
Ceutorhynchus .....	198
<b>pervestitus</b> .....	267
Chætarthria .....	56, 216
<b>minor</b> .....	216
Chætocnema .....	158
Chætophlœus .....	203
Chalcolepidius .....	108
Chalcophora .....	115
Charissa .....	130
Chilocorus .....	85
Chilometopon .....	163
Chlœnius .....	49
Chlamys .....	153
Choleva .....	59
Chrysanthia .....	178
Chrysobothris .....	117
Chrysochus .....	154
Chrysomela .....	155
CHRYSOMELIDÆ .....	151
Chrysophana .....	120

Cibdelis .....	170
Cicindela .....	37
CICINDELIDÆ .....	37
CIIDÆ .....	135
Cilea .....	74
Cionistes .....	195
Cis .....	135
Cistela .....	177
CISTELIDÆ .....	175
Clambus .....	60
Cleonus .....	192
<b>erysimi</b> .....	261
<b>pacificus</b> .....	260
CLERIDÆ .....	128
Clerus .....	129
Clivina .....	41
Clytus .....	146
Cnemeplatia .....	172
Cnemidotus .....	52
Cnemodus .....	163
Coccinella .....	83
COCCINELLIDÆ .....	83
Cœlambus .....	53
<b>femoratus</b> .....	212
<b>pedalis</b> .....	212
Cœlogaster .....	198
Cœnocara .....	133
Cœlocnemis .....	169
Cœlomorpha .....	166
Cœlus .....	166
Cœnopœus .....	150
Cœnonycha .....	140
Colaspidea .....	154
Colaspis .....	154
Colastus .....	98
Collops .....	123
<b>argutus</b> .....	242
Colon .....	59
Colpodota .....	64
COLYRIDÆ .....	88
Colymbetes .....	54
Conibius .....	171
Coninomus .....	100
Coniontis .....	165
Connophron .....	60
Cononotus .....	178
Conosoma .....	74
Conotelus .....	98
Conotrachelus .....	197
Copidita .....	178
Copris .....	136
Coptocycla .....	160
Coptotomus .....	53
Copturus .....	197
Corphyra .....	180

Corticaria .....	101	Dasytastes.....	128
CORYLOPHIDÆ.....	82	<b>insularis</b> .....	251
Corymbites .....	113	Dasytellus .....	127
Corynetes .....	130	Dasytes.....	127
Coscinoptera .....	152	<b>clementæ</b> .....	250
Cossonus .....	201	<b>musculus</b> .....	249
Cotalpa.....	141	Decarthron.....	62
Coxelus.....	88	Deleaster.....	77
Craniotus.....	161	Dendrobium.....	145
Cratidus.....	169	Dendroctonus .....	203
Cregya.....	130	Deporax.....	187
Cremastochilus.....	142	Deretaphrus.....	89
Creniphilus .....	57	Dermestes.....	92
<b>elegans</b> .....	218	DERMESTIDÆ.....	92
Creophilus.....	68	Derobrachys.....	142
Crepidodera .....	158	Deronectes .....	53
Crossidius.....	146	Desmocerus.....	147
Crypbalus .....	202	Desmopachria.....	52
Cryptarcha .....	99	Desmoris.....	193
Cryptobium.....	72	Diabrotica.....	156
Cryptocephalus .....	153	Diachus.....	153
Cryptoglossa.....	164	Diaphorus.....	47
Cryptognatha.....	85	Dibolia.....	159
Cryptohypnus.....	109	Dicentrus .....	143
Cryptophagus.....	91	Dicerca.....	115
CRYPTOPHAGIDÆ.....	91	Dichelonycha.....	139
Cryptorhopalum.....	93	Dinapate.....	134
Ctenisis .....	61	Dinocleus.....	192
Ctenobium.....	132	Dinoderus .....	134
CUCUJIDÆ.....	89	Diodyrhynchus.....	186
Cucujus.....	90	Diplocelus.....	92
Cupes.....	135	Diplotaxis.....	140
CUPESIDÆ.....	135	Dirotognathus.....	189
CURCULIONIDÆ.....	190	Discoderus.....	50
Cybister.....	54	Disonycha .....	157
Cybocephalus.....	99	Ditemnus .....	123
Cychrus.....	39	Ditoma.....	88
Cyclocephala .....	141	Ditylus .....	178
Cycloneda.....	84	Doliema.....	170
Cyllene.....	146	Dolopius.....	111
Cymatodera.....	129	Donacia .....	151
Cymbiodyta .....	57	Dorytomus.....	193
Cymindis.....	49	Drachylis.....	179
Cynæus.....	173	Drasterius.....	110
Cyphon.....	106	Dromæolus.....	107
Cyrtusa.....	60	Dromius.....	48
Cystodemus .....	183	Dryocetes .....	202
Cytilus.....	104	Dryops .....	105
		Dryophthorus.....	200
DACNE.....	88	Dyschirius.....	41
Dacoderus.....	163	<b>affinis</b> .....	209
Dactylosternum .....	58	<b>duplicatus</b> .....	209
DASCYLLIDÆ.....	105	<b>unipunctatus</b> .....	207
Dascyllus.....	106	Dyslobus .....	188
Dasycerus.....	100	Dystaxia .....	118

Dysticheus.....	189	Euphoria.....	141
DYTISCIDÆ.....	52	Euphorticus.....	47
Dytiscus.....	54	Euplectus.....	63
ECHIDNOGLOSSA.....	64	Eupsophus.....	170
Echocerus.....	172	Eurelymis.....	128
Edrotes.....	161	Europs.....	103
Ega.....	48	Eurygenius.....	180
Elaphidion.....	144	Eurymetopon.....	162
Elaphrus.....	40	Eurypogon.....	105
Elasmocerus.....	128	<b>confusus</b> .....	239
<b>californicus</b> .....	251, 252	Euryscopa.....	152
Elassoptes.....	201	Eusattus.....	166
Elater.....	109	difficilis.....	256
ELATERIDÆ.....	107	Eustilbus.....	82
Elatrinus.....	110	<b>notabilis</b> .....	230
Eleates.....	174	Eustrophus.....	177
Elissa.....	189	Euthysanius.....	114
Elleschus.....	196	Exema.....	153
Ellychnia.....	121	Exochomus.....	85
Elmis.....	105	<b>histrio</b> .....	230
Eleodes.....	167	FALAGRIA.....	64
Embaphion.....	168	Formicomus.....	181
Emmenastus.....	162	Fuchsina.....	102
Emphyastes.....	192	GALERITA.....	47
Endalus.....	194	Galerucella.....	156
Endeodes.....	124	Gastroidea.....	155
ENDOMYCHIDÆ.....	87	Geoderces.....	189
Enicmus.....	100	Geodercodes.....	189
Enneathron.....	135	Geodromicus.....	77
Epicauta.....	184	GEORYSSIDÆ.....	104
Epierus.....	95	Georyssus.....	104
Epimechus.....	196	Geotrupes.....	138
<b>remulus</b> .....	266	Glyptina.....	158
<b>arenicolor</b> .....	265	Glyptoscelimorpha.....	118
Epitragus.....	162	Glyptoscelis.....	154
Epitrix.....	158	Gnathium.....	184
Epurea.....	98	Gnathocerus.....	172
Erchomus.....	74	Gnathotrichus.....	201
Eretes.....	54	Gonioctena.....	155
Ergates.....	142	Gonops.....	203
Eriopis.....	83	Grynocharis.....	103
Ernobius.....	131	Gyascutus.....	115
Eros.....	121	Gymnopyge.....	139
EROTYLIDÆ.....	87	Gyretes.....	55
Eschatocrepis.....	128	GYRINIDÆ.....	54
Esthesopus.....	109	Gyrinus.....	55
Euceratocerus.....	133	Gyrorycha.....	67
Eucinetus.....	106	Gyrotus.....	198
Eucnemis.....	107	HABROCERUS.....	74
Eucrossus.....	144	Hadrobregmus.....	132
Eucyllus.....	189	Hadrotes.....	68
Eudasytes.....	125	HALIPLIDÆ.....	51
Eulabis.....	168	Haliplus.....	51
Eupagoderes.....	187		

Haltica .....	157	Hymenorus .....	176
Haplidus .....	144	Hyperaspidium .....	86
Haploderus .....	76	Hyperaspis .....	85
Harmonia .....	84	<b>excelsa</b> .....	232
Harpalus .....	50	<b>spiculinata</b> .....	232
Hedobia .....	131	Hyperplatys .....	150
Helochares .....	57	Hypnoldeus .....	109
Helodes .....	106	Hypophloeus .....	174
Helophorus .....	55	Hyporhagus .....	177
Helops .....	175	Hypothenemus .....	202
Hemiglyptus .....	158		
Hemiptychus .....	132, 252	IDOMEA .....	151
<b>integer</b> .....	253	<b>californica</b> .....	256
<b>luteotectus</b> .....	253	Ilybiosoma .....	53
<b>palliatu</b> s .....	253	Iphthimus .....	169
Henoticus .....	91	Ipochus .....	150
Hesperobæus .....	103	Ips .....	99
Heterius .....	95	Ischnonemis .....	146
HETEROCERIDÆ .....	105	Isoglossa .....	66
Heterocerus .....	105	Isomira .....	176
Heterothops .....	67		
<b>carbonatus</b> .....	220	LACCOBIUS .....	57
Hippodamia .....	83	Lacconotus .....	177
Hippomelas .....	115	Laccophilus .....	52
HISTERIDÆ .....	93	Lachnophorus .....	47
Hister .....	94	Lachnosterna .....	141
<b>simplicipes</b> .....	235	Læmophloeus .....	90
Hololepta .....	93	Lamprorhiza .....	121
Holoparamæcus .....	99	LAMPYRIDÆ .....	121
Holopleura .....	145	Languria .....	87
Homalium .....	78	<b>californica</b> .....	235
Hoplandria .....	64	Lapethus .....	89
Hoplia .....	139	Lara .....	104
Horistonotus .....	108	Lasconotus .....	88
<b>flavidus</b> .....	240	LATHRIDIDÆ .....	99
Hybodera .....	145	Lathridius .....	109
Hydaticus .....	54	Lathrimæum .....	78
Hydnobius .....	59	Lathrobium .....	72
Hydnocera .....	130	Lathropus .....	91
Hydræna .....	56	Lebasiella .....	130
Hydrobius .....	57	Lebia .....	48
Hydrocharis .....	56	Lema .....	152
Hydrochus .....	55	Leptacinus .....	70
HYDROPHILIDÆ .....	55	Leptalia .....	148
Hydrophilus .....	56	Leptinotarsa .....	155
Hydroporus .....	53	Leptorus .....	73
Hydroscapha .....	81	<b>californicus</b> .....	224
HYDROSCAPHIDÆ .....	81	<b>longipennis</b> .....	225
Hydrotrupes .....	53	Leptoschema .....	112
Hygrotus .....	52	Leptotrix .....	158
Hydrovatus .....	52	Leptura .....	149
Hylastes .....	203	Leptusa .....	67
Hylesinus .....	203	Lepyrus .....	191
Hylotrupes .....	143	Lesteva .....	77
Hylurgops .....	203	Ligyrus .....	141

Limnebius .....	57	Megilla.....	83
Limnichus.....	104	Megobrium.....	145
Limnobaris.....	199	Melanactes.....	114
Limonius.....	111	MELANDRYIDÆ.....	177
Lina.....	155	Melanophila.....	116
Liodes.....	59	Melanophthalma.....	101
Lispinus.....	79	Melanotus.....	111
Lissonotus.....	145	Melasis.....	107
Listrochelus.....	141	Meligethes.....	99
Listronotus.....	192	Meloe.....	183
Listrus.....	127	MELOIDÆ.....	183
<b>definitus</b> .....	248	Meristhus.....	108
Litargus.....	92	Merotemnus.....	173
Lithocharis.....	73	Metachroma.....	154
Lixus.....	192	Metaclisa.....	173
Lomechusa.....	65	Metaleptus.....	146
<b>angusta</b> .....	219	Metamasius.....	200
Longitarsus.....	158	Methia.....	151
Loricera.....	40	Metophthalmus.....	100
LUCANIDÆ.....	135	Metopotoma.....	200
Ludius.....	111	Metrius.....	41
Luperodes.....	157	Micracis.....	202
Lyctus.....	134	Microglossa.....	66
Lypsimena.....	151	Microlipus.....	124
		Micromastus.....	197
MACRONYCHUS.....	105	Micropeplus.....	80
Macropogon .....	105	Microphotus.....	121
Macrops.....	192	Microphopa.....	159
Macrorhoptus.....	195	Microschattia.....	164
Macrorhyncholus.....	201	Miloderes.....	188
Magdalinops.....	195	Mimetes.....	188
Magdalis.....	194	Minyomerus.....	187
MALACHIDÆ.....	123	Molorchus.....	145
Malachius.....	124	Monarthrum.....	201
<b>acutipennis</b> .....	245, 248	Monilema.....	150
<b>directus</b> .....	245, 248	Monohammus.....	150
<b>inornatus</b> .....	243, 248	MONOMMIDÆ.....	177
<b>nigrinus</b> .....	246, 248	Monotoma.....	103
<b>pristinus</b> .....	244, 248	MONOTOMIDÆ.....	103
<b>prolixicornis</b> .....	246, 248	Monoxia.....	156
Malacopterus.....	144	Mordella.....	179
Mallodon.....	142	MORDELLIDÆ.....	179
Malocorhinus.....	157	Mordellistena.....	179
Malthodes.....	123	Norius.....	63
Maseochara.....	66	Motschulskium.....	80
Mastinocerus.....	122	Mycetina.....	87
Matheteus.....	121	Mycetochares.....	176
Mecynotarsus.....	183	MYCETOPHRAGIDÆ.....	91
Mecysmus.....	170	Mycetophagus.....	91
Medon.....	72	Mycetoporus.....	74
Megapenthes.....	110	Mycterus.....	178
Megasternum.....	58	Mylacus.....	189
Megarthritis.....	79	Myllæna.....	67
Megataphrus.....	88	Myochrous.....	154
Megaleates.....	174	Myodites.....	186

Myrmecixenis.....	92	Oodes.....	50
Myrmecochara.....	66	Ophistomis.....	150
Myrmedonia.....	65	Orchestes.....	196
Mysia.....	84	Orimodema.....	188
NACERDES.....	178	Orphilus.....	93
Nemia.....	83	Orobanus.....	79
Narpus.....	105	Oropus.....	63
Narthecius.....	90	Orsodacna.....	152
Nausibius.....	90	Orsonyx.....	140
Nebria.....	40	Orthocis.....	135
Necrophilus.....	58	Orthoperus.....	82
Necrophorus.....	58	Orthoptochus.....	189
Necydalis.....	148	Orthoris.....	199
Nemognatha.....	184	Orus.....	73, 222
Neoclytus.....	147	<b>femoratus</b> .....	224
NITIDULIDÆ.....	98	<b>fraternus</b> .....	223
Nitidula.....	99	<b>montanus</b> .....	223
Nocheles.....	188	parallelus.....	223
Nomaspis.....	183	punctatus.....	223
Nomidius.....	188	Othius.....	71
Noserus.....	163	OTHNIDÆ.....	177
Nosodendron.....	103	Othnius.....	177
Nothus.....	177	Otidoccephalus.....	194
Notibius.....	171	OTIORHYNCHIDÆ.....	187
Notiophilus.....	40	Oxacis.....	179
<b>obscurus</b> .....	207	Oxoplus.....	146
Notoxus.....	182	Oxygonus.....	113
<b>caudatus</b> .....	257	Oxylemus.....	89
Nyctoporis.....	164	Oxypoda.....	66
OBERA.....	151	Oxyporus.....	75
Ochodæus.....	138	Oxytelus.....	76
Ochthebius.....	55	Ozognathus.....	131
<b>costipennis</b> .....	214	PACHYBRACHYS.....	153
<b>wickhami</b> .....	213	Pachyphanes.....	193
Octotemnus.....	135	Pachyplectrus.....	138
Odontæus.....	138	Pachyta.....	148
Odontosphindus.....	135	Pactopus.....	115
Odontota.....	159	Pæderus.....	73
ÆDEMERIDÆ.....	178	Palæoxenus.....	107
Ædionychis.....	157	Palaminus.....	73
Æme.....	144	Panormus.....	189
Olibrus.....	82	Papusus.....	60
Oligomerus.....	131	Paragoges.....	197
Oligopteris.....	72	Parandra.....	142
Oligota.....	67	Paraptochus.....	188
Olisthærus.....	74	PARNIDÆ.....	104
Omophron.....	39	Paromalus.....	95
Omosita.....	99	Patrobus.....	44
Omus.....	37	Pediacus.....	90
Oncerus.....	139	Pelenomus.....	198
Oniticellus.....	136	Peltis.....	103
Onthophilus.....	95	Pentaphyllus.....	174
Onychobaris.....	199	Pentaria.....	179
		Perlecompus.....	44



Perigona.....	47	Podapion.....	191
Perimegatoma.....	92	Pœcilobrium.....	145
Peritelinus.....	189	Pœcilonota.....	116
Peritelodes.....	189	Pogonocherus.....	151
Peritelopsis.....	189	Pogonus.....	44
Perothops.....	115	Polemios.....	123
Perthalykra.....	99	Polycaon.....	134
Petalium.....	132	<b>megalops</b> .....	254
PHALACRIDÆ.....	81	Polycesta.....	118
Phalacrus.....	81	Polyphylla.....	141
Phaleria.....	173	Polystoma.....	65
Phellopsis.....	163	Pontomalota.....	64
Philhydrus.....	57	Poreospasta.....	184
<b>conjunctus</b> .....	217	Priacma.....	135
Philonthus.....	68	Priognathus.....	178
Philophnga.....	49	Prionus.....	142
Phlœodes.....	163	Pristonychus.....	46
Phlœonemus.....	88	Pristoscelis.....	125
Phlœopora.....	65	Promecotarsus.....	194
Phlœosinus.....	203	Protinus.....	79
Phlœotrya.....	177	Prostomis.....	90
Phobetus.....	141	Psammodius.....	136
Phodaga.....	186	PSELAPHIDÆ.....	61
Phthora.....	172	Pselaptrichus.....	62
Phyconomus.....	103	Pselaptus.....	62
Phycocœtes.....	194	Psephenus.....	104
Phyllobrotica.....	156	Pseudebæus.....	125
Phyllotreta.....	159	Pseudobaris.....	199
Phyllotrox.....	194	Pseudomorphus.....	51
Phymaphora.....	87	Pseudopsis.....	75
Phymatinus.....	188	<b>detrita</b> .....	226
Phymatodes.....	143	<b>minuta</b> .....	227
Piazurus.....	197	Psoa.....	134
Pinacodera.....	49	Psylliodes.....	159
Pinodytes.....	59	Psyllobora.....	84
Pinophilus.....	73	Ptenidium.....	80
Pityobius.....	112	Pterostichus.....	44
Pityophagus.....	99	<b>inermis</b> .....	211
Pityophthorus.....	201	Pterotus.....	121
Placonycha.....	106	Ptilinus.....	133
Plagiodera.....	155	Ptilium.....	80
Plastocerus.....	114	PTINIDÆ.....	130
Platycholens.....	59	Ptinodes.....	132
Platydema.....	174	Ptinus.....	130
Platynus.....	46	Ptomaphagus.....	59
Platystethus.....	76	Ptosima.....	120
Plectrodes.....	140	Purpuricenus.....	146
Plegaderus.....	97	Pyrochroa.....	183
Pleocoma.....	138	PYROCHROIDÆ.....	183
Plesiocis.....	135	Pyropyga.....	121
Pleurophorus.....	137	Pyrotichus.....	148
Plochionus.....	49	PYTHIDÆ.....	178
Pocadius.....	99		
Podabrus.....	122	QUEDIUS.....	67

REICHENBACHIA .....	62	Scopæodera .....	73
Revelieria .....	101	Scopæus .....	73
Rhagium .....	148	<b>armiger</b> .....	225
Rhagodera .....	88	SCYDMÆNIDÆ .....	60
Rhantus .....	54	Scydmaenus .....	60
Rhexidius .....	63	Scymnillus .....	86
Rhigopsis .....	187	Scymnus .....	86
Rhinomacer .....	186	<b>dentipes</b> .....	234
RHINOMACERIDÆ .....	186	<b>megacephalus</b> .....	233
Rhinoplatia .....	179	<b>minus</b> .....	234
Rhinosimus .....	178	<b>tædatus</b> .....	233
RHIPICERIDÆ .....	107	Scyphophorus .....	200
RHIPIPHORIDÆ .....	186	Scythropus .....	190
Rhipiphorus .....	186	Serica .....	140
Rhizobius .....	87	Sericoderus .....	82
Rhizophagus .....	99	Sericosomus .....	113
Rhodobænus .....	200	Sibynes .....	197
Rhynchites .....	186	Silis .....	122
<b>æratoides</b> .....	260	Silpha .....	58
RHYNCHITIDÆ .....	186	SILPHIDÆ .....	58
Rhyncholus .....	201	Silusa .....	66
Rhypodes .....	189	Silvanus .....	89
Rhyssematus .....	197	Sinodendron .....	135
Rhyssenus .....	137	Sinoxylon .....	133
Rhyssodes .....	89	Sipalia .....	67
RHYSSODIDÆ .....	89	Sitodrepa .....	131
Romaleum .....	144	Sitones .....	190
Rosalia .....	145	Smicrips .....	99
SACIUM .....	82	Smicronyx .....	193
Sagola .....	63	Smicrus .....	80
Salpingus .....	178	Smilia .....	85
Sandalus .....	107	<b>reversa</b> .....	231
Saperda .....	151	Sognorus .....	61
Saprinus .....	96	Somatium .....	67
<b>consobrinus</b> .....	237	Sphæricus .....	130
<b>liticolus</b> .....	236	SPHÆRIIDÆ .....	81
Sarpedon .....	107	Sphærites .....	59
Saxinis .....	152	Sphærius .....	81
Scalenarthrus .....	62	Sphenophorus .....	200
SCAPHIDIIDÆ .....	81	<b>tardus</b> .....	269
Scaphisoma .....	81	SPHINDIDÆ .....	135
SCARABÆIDÆ .....	136	SPONDYLIDÆ .....	142
Scarites .....	41	Spondylis .....	142
Scelolyperus .....	156	Stamoderes .....	187
Schizillus .....	164	STAPHYLINIDÆ .....	64
Schizogenius .....	42	Staphylinus .....	68
<b>litigiosus</b> .....	210	Stenelmis .....	105
<b>seticollis</b> .....	209	<b>nubifer</b> .....	238
<b>validus</b> .....	210	Stenochidus .....	175
Schizonotus .....	194	Stenocolus .....	106
Schizopus .....	118	Stenolophus .....	50
Sciopithes .....	188	Stenopelmus .....	194
SCOLYTIDÆ .....	201	Stenopodius .....	160
Scolytus .....	202	Stenoptochus .....	189
		Stenosphenus .....	146

Stenotrichus.....	169	Tinotus .....	65
Stenus.....	71	Tomicus.....	202
Stereopalpus.....	180	Toxotropis.....	203
Stibia.....	161	Toxotus .....	148
Stilicus.....	73	Trachypachys .....	39
<b>occiduus</b> .....	225	Trachyscelida.....	156
Strangalia.....	149	Tragidion .....	146
Sunius.....	73	Trechus.....	44
Synaphæta.....	150	<b>pomonæ</b> .....	211
Synchita.....	88	Triarthron .....	59
Synertha.....	194	Tribalister.....	94
Syneta.....	152	Tribalus.....	94
Systema.....	158	Tribolium.....	172
		Trichobaris .....	199
TACHINUS .....	74	Trichochrous .....	125
Tachycellus.....	51	Trichodes .....	129
Tachyporus .....	74	Trichodesma .....	132
Tachypterus.....	195	TRICHOPTERYGIDÆ.....	80
Tachys.....	43	Trichopteryx.....	80
Tachyusa.....	65	Trichoton .....	171
Tanaops.....	124	Trigonogenius.....	130
Tanarthrus.....	182	Trigonoscuta.....	187
Tanyrhinus.....	78	Trigonurus .....	80
Taphrocerus.....	121	Triglyphus.....	190
Tarphiota.....	65	Triorophus .....	161
Tecnophilus.....	48	Trirhabda .....	155
Tegrodera .....	186	Tritoma .....	88
Telephorus.....	122	Trogloderus.....	168
Tenebrio .....	170	Trogodendron.....	129
TENEBRIONIDÆ .....	161	Trogoderma.....	93
Tenebrioides.....	102	Trogophkæus.....	76
Teretrius.....	97	Trogosita.....	102
Tetracha.....	37	TROGOSITIDÆ .....	102
Tetragonodera .....	48	Trogosma.....	143
Tetraopes.....	151	Tropisternus.....	56
Tetropium.....	143	<b>salsamentus</b> .....	214
Thalpius.....	47	Trox.....	138
Thanasimus.....	129	Trypopitys .....	132
Theca .....	132	Tychius.....	196
Thermonectes.....	54	Tychus .....	62
Thiasophila.....	66	Tyloderma.....	197
Thinobius.....	77	Typhlusechus .....	163
Thinopinus.....	68	Typhcea.....	92
Thinoxenus.....	189	Typophorus .....	154
Thinusa.....	66	Tyrus.....	61
Thricolema.....	152		
Thricolepis.....	189	ULOCHETES.....	148
Throscidæ.....	115	Uloma.....	173
Throscinus.....	105	Ulosonia.....	173
Throscus .....	115	Ulus.....	171
Thyce.....	141	Unamis.....	77
Thlea.....	77	Usechus .....	163
Timarcha.....	155		

VALDA.....	61	Xylophilus.....	183
Valgus.....	142	<b>brunnescens</b> .....	258
Vanonus.....	258	<b>constrictus</b> .....	259
Vectura.....	128	<b>nucleus</b> .....	259
Vellica.....	77	Xyloterus.....	202
Vrilletta.....	132	Xylotrechus.....	147
		Xystropus.....	175
XANTHOCHROA.....	178	YUCCABORUS.....	200
Xantholinus.....	70	ZABROTES.....	160
Xenomycetes.....	87	Zalobius.....	77
Xestobium.....	131	Zarhipis.....	122
Xyleborus.....	202	Zascelis.....	197
Xyletinus.....	132	Zeugophora.....	152
Xylobius.....	107	Zonitis.....	184
Xylocleptes.....	202	Zopherus.....	163
Xylocrius.....	144	Zuphium.....	47

## ERRATA.

- Page 61, ninth line, insert ( ), to read *S. (catalinus) angustus*.  
Page 108, second line from bottom, for *Horistonatus* read *Horistonotus*.  
Page 120, for *Crysophana* read *Chrysophana*.  
Page 132, under *Trypopitys*, for *temuilineata* read *tenuilineata*.  
Page 137, first line, for *cælatus* read *cælatus*.  
Page 143, for *Decentrus* read *Dicentrus*.  
Page 151, under *Methia*, omit ( ) about Linell.  
Page 199, under *Trichobaris*, for *metilloides* read *metelloides*.







# Occasional Papers

OF THE

CALIFORNIA

ACADEMY OF SCIENCES

---

VIII

---

SAN FRANCISCO

1901

PUBLICATION COMMITTEE

CHARLES H. GILBERT, *Chairman*

WILLIAM E. RITTER

WILLIAM A. SETCHELL

---

*EDITORS OF ZOOLOGICAL PUBLICATIONS*

CHARLES H. GILBERT

WILLIAM E. RITTER

OCCASIONAL PAPERS  
OF THE  
CALIFORNIA ACADEMY OF SCIENCES

VIII

---

LIST  
OF THE  
Coleoptera  
OF  
Southern California,  
WITH  
Notes on Habits and Distribution  
AND  
Descriptions of New Species

BY  
H. C. FALL

*Issued November 11, 1901*

SAN FRANCISCO  
CALIFORNIA ACADEMY OF SCIENCES  
1901















SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00773 9121